

ORIGINAL ARTICLE



Perceived Exertion Responses to Exercise Differ for Progressively Increasing and Decreasing Order of Intensity: A Crossover Design Study

^{1,2}Wootae Lim ^{*}

¹Department of Physical Therapy, Rehabilitation Science Faculty, Woosong University, Daejeon, Republic of Korea. ²Woosong Institute of Rehabilitation Science, Rehabilitation Science Faculty, Woosong University, Daejeon, Republic of Korea.

Submitted February 2, 2022; Accepted in final form March 17, 2022.

ABSTRACT

Background. The effects of exercise intensity on the rating of perceived exertion (RPE) have been widely studied. However, in most of these studies, the contraction intensities were provided in a randomized order. **Objectives.** This study aimed to examine the influence of intensity order on RPE. **Methods.** During the first period, three trials of isometric contractions were performed at three different target intensities in the following order: 25%, 50%, and 75% of maximal voluntary contraction (MVC) in group A and the reverse order: 75%, 50%, 25% of MVC in group B. Each group's intensity order provided during the first period was reversed during the second period. The RPE was measured at each target intensity. **Results.** There was a significant difference in the perceived intensity of 75% of MVC between groups during the first period. The perceived intensity of 75% of MVC in group A during the second period and group B during the first period was significantly lower than the target intensity. **Conclusion.** The order of exercise intensity affected the perceived exertion responses in this study. When establishing an exercise program consisting of varying intensities, practitioners should consider the order of intensity implementation.

KEYWORDS: *Exercise, Hamstring Muscles, Isometric Contraction, Physical Exertion.*

INTRODUCTION

Resistance exercise is essential for maintaining a healthy body, and intensity is considered essential during exercise. It can be challenging to improve muscle strength if the intensity is too low, and excessive high-intensity exercise can cause exercise-induced injuries. To establish an effective exercise program, it is vital to measure the exercise intensity. One-repetition maximum (1RM) is widely used to determine the appropriate dose for resistance exercise in clinical practice. However, equipment may be needed to measure 1RM accurately (1), and the computation of complex calculations is necessary to achieve

accurate prediction (2). In addition, since it is time-consuming, the applicability of 1RM measures may decrease in large groups (3). Thus, the rating of perceived exertion (RPE) has been proposed as an alternative method to evaluate exercise intensity (4).

RPE is used in the general population and target groups, including patients and athletes (5–7). RPE relies on the individual's subjective judgment and has reasonably high validity and reliability (8, 9). Many studies have shown that RPE correlates with exercise intensities and physiological properties such as maximum oxygen consumption, heart rate,

*. Corresponding Author:

Wootae Lim, Ph.D.

E-mail: wootaeclimpt@wsu.ac.kr

and respiration rate (10, 11). RPE may be used for aerobic exercise; however, it is an effective method to evaluate exercise intensity, especially for monitoring resistance training intensity. However, most prior studies on exercise intensity had considerable limitations. Some studies applied different intensities after randomization (12–15). Even at the same intensity, RPE responds differently depending on the muscle and exercise duration (16, 17), and its ratings tend to be higher during concentric exercise than during eccentric exercise (18).

Additionally, prior resistance exercise can affect the performance of the muscles subsequently used (19). Even if the exercise program consists of the same intensities, the RPE may differ depending on the order of intensities. The present study compared the RPE between two groups of three different intensities that were progressively increased or decreased.

Currently, two main types of RPE are widely used—the Borg scale (6–20) and the modified Borg category ratio (CR-10). The OMNI-Resistance Exercise Scale (OMNI-RES) is also used less frequently (9). The OMNI-RES uses an easy-to-understand visual scale, but its use during aerobic exercise is limited (20). The Borg 6–20 can be useful for both aerobic and resistance exercises, but because of its wide range from 6 to 20, it is difficult for people to intuitively match their exercise intensity to the Borg 6–20. It is easier to match the exercise intensity to the CR-10 as it only uses 0–10; however, this range can also be too narrow to capture subtle changes, which might be meaningful in clinical practice. In this study, the range of RPE was set as 0–100 (100 was anchored as maximal). This exactly matched the contraction intensity, ranging from 0–100 (21). Based on the maximum voluntary contraction (MVC), the target intensities were composed of three submaximal contraction intensities—75%, 50%, and 25% of MVC.

This study aimed to examine the influence of intensity order on RPEs between two groups. The groups performed isometric contractions at three different submaximal intensities that were progressively increased or decreased. The study followed a crossover design; hence, the order of intensity presentation during the first period was reversed during the second period.

MATERIALS AND METHODS

Participants. Thirty healthy adults were randomly divided into two groups (Table 1).

Participants with no prior surgeries on the hip, knee, and ankle joints in the past six months were included. The study was conducted after the Institutional Review Board of ooooooo University reviewed and approved the protocol, and informed consent was obtained from all study participants before any study-related procedures.

Table 1. Demographics of participants

	Group A (n=15)	Group B (n=15)	<i>p</i>
Age	21.9±1.3	22.0±1.6	0.801
Weight	59.7±10.9	62.7±13.8	0.506
Height	164.7±8.4	165.7±8.9	0.852

Procedures. When the participant lay on the treatment table in a supine position, the examiner attached a strap connected to a sling rope to the participant's ankle. The other end of the sling rope was connected to the sling system attached to the ceiling, and the angles of the rope and lower extremity were adjusted to 90 degrees (22). The examiner performed passive straight leg raises. Slowly, the examiner induced hamstring elongation before the participant felt any discomfort or pain. In the end range, each participant performed three trials of MVC. During MVC, the examiner motivated the participant to use maximum effort. The isometric contraction was performed for 5 s, and 10 s of rest was given between trials during each trial.

After performing three trials of MVC, a break of 3 min was provided before performing the isometric contractions at three different submaximal intensities. After setting the submaximal target intensity in 25%, 50%, and 75% of MVC, Group A performed isometric contractions at each assigned target intensity. Group B performed the contractions in the reverse order of group A: 75%, 50%, and 25% of MVC. Similar to MVC, three trials of isometric contractions were performed at each target intensity. Rest was given for 10 s between trials and 3 min between target intensities. The perceived intensity was recorded after three trials of isometric contractions at each target intensity. The examiner asked each participant to evaluate the intensity, the relative perceived exertion compared to the maximum intensity, in the range of 0%–100%, with 100% being the maximal intensity. After performing isometric contractions at three different submaximal intensities with a progressive increase or decrease depending on the groups,

which was the first period, a 10-min break was given before starting the second period. During the second period, the order was reversed. Group A performed the target intensities in the following order: 75%, 50%, and 25% of MVC, while group B performed the target intensities in the reverse

order: 25%, 50%, and 75% of MVC. During the second period, the perceived intensity was verbally requested and recorded after three trials of isometric contractions at each target intensity (Figure 1).




Figure1. Study flow chart: a crossover study design.

Statistical analysis. Normally distributed data were assessed using the Shapiro–Wilk test. Differences in age, weight, height, and perceived intensity between groups A and B were statistically compared using an independent t-test or the Mann–Whitney U-test. The Wilcoxon signed-rank test examined within-group differences (between the first and second periods). The carryover effect was also assessed using an independent t-test. The differences in perceived intensities among three submaximal intensities within each group were statistically compared using the Friedman test with the Wilcoxon signed-rank test as a post house. The target intensity was compared with the perceived intensity using the one-sample Wilcoxon signed-rank test. Data analysis was performed using IBM SPSS Statistics 25 (IBM Corp., Armonk, NY, USA). Statistical significance was set at $p < 0.05$. All values are reported as mean \pm standard deviation.

RESULTS

There were no significant differences in age, weight, and height between groups A and B (Table 1). At 75% of MVC, the perceived intensity was significantly different between groups A and B ($p = 0.017$) during the first period, but it was not significantly different during the second period (Table 2). No significant differences between groups A and B were observed at 50% and 25% of MVC during the first and second periods. There was a

statistically significant difference in perceived intensity among three different submaximal intensities in group A during the first and second periods (Table 3) and group B during the first and second periods (Table 4). There was no carryover effect on perceived intensities from the first to the second period. The perceived intensity of 75% of MVC in group A during the second period ($p = 0.011$) and in group B during the first period ($p = 0.003$) was significantly different from the target intensity (Figure 2).

DISCUSSION

RPE during isometric contraction with various intensities is widely studied. In most studies, the contraction intensities were provided in a randomized order. However, unlike in research settings, the clinical settings allow us to vary the order of intensities in a progressively increasing or decreasing manner, especially in resistance exercise programs. Hence, it is essential to study the effects of intensity ordering. RPE is not solely determined by the information received from the sensory receptor in the joint and muscle during muscle contraction (23–25). Even in the same program, RPE may be different if the intensity order is not the same. This study confirmed the RPE at 75%, 50%, and 25% of MVC. The order of these three different intensities was provided in a progressively increasing or decreasing manner. A crossover design was used to reduce potential confounding variables.

Table 2. The difference in perceived intensity between Group A and B

Target intensity (% MVC)	75%	50%	25%
First period			
Group A	73.1±5.6	46.5±7.3	24.7±3.1
Group B	67.7±6.6	50.2±6.5	25.0±6.8
<i>p</i>	0.017	0.354	0.800
Second period			
Group A	70.1±5.8	48.3±6.9	24.5±4.5
Group B	73.7±4.6*	49.5±4.7	23.1±4.8
<i>p</i>	0.118	0.883	0.317


* Significant difference compared to the one during the first period.

Table 3. The significant difference in perceived intensity among three different submaximal intensities in Group A

Target intensity (% MVC)	75%	50%	25%
First period			
75%	-	0.007	<0.001
50%	-	-	0.021
25%	-	-	-
Second period			
75%	-	0.009	<0.001
50%	-	-	0.012
25%	-	-	-

Table 4. The significant difference in perceived intensity among three different submaximal intensities in Group B

Target intensity (% MVC)	75%	50%	25%
First period			
75%	-	0.022	<0.001
50%	-	-	0.007
25%	-	-	-
Second period			
75%	-	0.010	<0.001
50%	-	-	0.010
25%	-	-	-

**Figure 2. Box and whisker plots with the minimum, maximum, first and third quartile, and median values of perceived intensity. *Significant difference compared to the target intensity.**

In the first period, the participants reported higher perceived intensity when the isometric contraction intensity was progressively increased than when the isometric contraction intensity was progressively decreased. However, the significantly higher perceived intensity was only noted at 75%. Between-group differences in perceived intensity were similarly observed in the second period, where the intensities were presented in reverse after crossover. Additionally, even when each group's first and second periods were compared, the higher perceived intensity was observed in 75% of MVC only when the intensity progressively increased. In conclusion, similar between- and within-group patterns were observed in this study. Thus, RPE values may differ even in the same exercise program when the program consists of various intensities in various orders. This difference is particularly noticeable during high-intensity exercises. High-intensity muscle contractions may differ in RPE depending on whether the contraction occurred in the last step of the progressive escalation or the first step of progressive de-escalation. The physical and/or psychological fatigue produced during high-intensity exercises may be affected by the order of intensity implementation.

If repeated muscle contractions at submaximal intensities are performed, and subsequent muscle contractions are additionally requested at different intensities, the muscle may experience physical fatigue before entering the final step, resulting in high-intensity muscle contractions. When exercise-induced physical fatigue occurred in previous studies, decreased physical effort and decreased motivation were observed during subsequent motor tasks (26). In the present study, repeated muscle contractions, performed in the first step of progressive escalation, resulted in physical fatigue and affected subsequent muscle contractions at high intensities. Specifically, greater physical exertion might be required to maintain consistent muscle activity in this study. In the Guo 2017 study, the fatigue group with a short rest interval during repetitive tasks showed a higher RPE than the control group (27). In the Pincivero 2000 study, RPE gradually increased when the contraction time was increased by 80% of MVC (17). In recent years, attention has been paid to physical and mental workloads (16, 28). Thus, many studies on the relationship between physical exercise load and cognitive performance have been conducted (29, 30). According to the

Marcora 2009 study, mental fatigue can impair motor performance and muscle capacity. This results in significantly higher RPE during physical exercise of the same intensity (31, 32). This is probably because mental factors affect cortical centers associated with perceived exertion (33). However, similar results were observed after a short rest during this study's first and second periods. It can be inferred that the effect on RPE, whether physical or physiological, is only temporary. If RPE is monitored in clinical practice, it is necessary to recognize that different RPEs can be measured at the same intensity, depending on the implemented exercise intensities. In particular, when the high-intensity muscle contraction is first applied in a gradually decreasing order, RPE may be somewhat lower than expected.

When comparing the perceived intensity at three different target intensities within the group, this was differentiated between target intensities. The perceived intensity measured at 75% of MVC was significantly higher than at 50% and 25% of MVC. The perceived intensity of 50% of MVC was also significantly higher than at 25% of MVC. In prior studies, participants could clearly distinguish the contraction force when the subsequent intensity level differed by 30%–50% (15). When comparing target intensity and perceived intensity in this study, the perceived intensity of 75% of MVC was significantly lower than the target intensity that de-escalated from high to low intensity in both groups. There were no significant differences between target intensity and perceived intensity in all other cases. In prior studies, underestimation was often reported. However, there is no clear consensus as to at what level of intensity underestimation occurs. In the quadriceps femoris muscle, the CR-10 scales were 1.2, 1.6, 2.9, 3.5, and 4.9 during 20%, 30%, 40%, 50%, and 60% of 1RM, respectively. In other words, RPE values were significantly lower than the target intensities, which were considered low- to moderate-intensity. However, there was no difference in the high intensity of 70%–90% (13). In another study, underestimation was reported at 10–90% (34) or 50%–90% (35). In most prior studies, intensity levels were assigned after randomization. It is, therefore, possible that the order of intensity was not uniformly randomized. In this study, when the level decreased from high to low intensity, the perceived intensity at high intensity was

significantly lower than the target intensity. In a previous study that examined pain after maximal and submaximal isometric contractions, the subject complained of significantly increased pain during high-intensity exercise than during low- to moderate-intensity exercise (36, 37).

Additionally, an increase in exercise intensity can decrease pleasure. Further, if the examiner imposes the exercise intensity and is not self-selected, this could decrease tolerance to higher intensities, which can be more easily linked to displeasure (38). Taken together, it may be more advantageous, in terms of emotional state and pain, to perform the exercise at high intensity first.

CONCLUSION

In conclusion, the order of intensity can affect the RPE value when various intensities are applied. Additionally, RPE is similar to target intensity when the intensity is gradually increased from low to high. However, when the intensity gradually decreases from high to low, the RPE at the high intensity is significantly lower than the

target intensity. Unlike in research settings, in clinical settings, intensity can be progressively increased or decreased depending on the purpose. When constructing an exercise program in which various intensities are used, practitioners should consider the order of intensity implementation.

APPLICABLE REMARKS

- The order of intensity can affect the perceived exertion responses when the intensity is gradually decreased from high to low.
- When establishing an exercise program consisting of varying intensities, practitioners should consider the order of intensity implementation.

CONFLICT OF INTERESTS

The author has no conflict of interest to disclose.

SUPPORT

This study was supported by 2020 Woosong University Academic Research Funding.

REFERENCES

1. Verdijk LB, van Loon L, Meijer K, Savelberg HHCM. One-repetition maximum strength test represents a valid means to assess leg strength in vivo in humans. *J Sports Sci.* 2009 Jan 1;27(1):59-68. [doi:10.1080/02640410802428089](#) [pmid:19031334](#)
2. LeSuer DA, McCormick JH, Mayhew JL, Wasserstein RL, Arnold MD. The Accuracy of Prediction Equations for Estimating 1-RM performance in the Bench Press, Squat, and Deadlift. *The Journal of Strength & Conditioning Research.* 1997 Nov;11(4):211-3. [doi:10.1519/00124278-199711000-00001](#)
3. Brzycki M. Strength Testing-Predicting a One-Rep Max from Reps-to-Fatigue. *Journal of Physical Education, Recreation & Dance.* 1993 January 1;64(1):88-90. [doi:10.1080/07303084.1993.10606684](#)
4. Eston R, Evans HJL. The Validity of Submaximal Ratings of Perceived Exertion to Predict One Repetition Maximum. *J Sports Sci Med.* 2009 December 1;8(4):567-73.
5. Dawes HN, Barker KL, Cockburn J, Roach N, Scott O, Wade D. Borg's rating of perceived exertion scales: do the verbal anchors mean the same for different clinical groups? *Arch Phys Med Rehabil.* 2005 May;86(5):912-6. [doi:10.1016/j.apmr.2004.10.043](#) [pmid:15895336](#)
6. Eston R. Use of ratings of perceived exertion in sports. *Int J Sports Physiol Perform.* 2012 Jun;7(2):175-82. [doi:10.1123/ijsspp.7.2.175](#) [pmid:22634967](#)
7. Pérez-Landaluce J, Fernández-García B, Rodríguez-Alonso M, García-Herrero F, García-Zapico P, Patterson AM, et al. Physiological differences and rating of perceived exertion (RPE) in professional, amateur and young cyclists. *J Sports Med Phys Fitness.* 2002 Dec;42(4):389-95.
8. Day ML, McGuigan MR, Brice G, Foster C. Monitoring exercise intensity during resistance training using the session RPE scale. *J Strength Cond Res.* 2004 May;18(2):353-8. [doi:10.1519/00124278-200405000-00027](#) [pmid:15142026](#)
9. Guidetti L, Sgadari A, Buzzachera CF, Broccatelli M, Utter AC, Goss FL, et al. Validation of the OMNI-cycle scale of perceived exertion in the elderly. *J Aging Phys Act.* 2011 Jul;19(3):214-24. [doi:10.1123/japa.19.3.214](#) [pmid:21727302](#)
10. Habibi E, Dehghan H, Moghiseh M, Hasanzadeh A. Study of the relationship between the aerobic capacity (VO₂ max) and the rating of perceived exertion based on the measurement of heart beat in the

- metal industries Esfahan. *J Educ Health Promot* [Internet]. 2014 Jun 23 [cited 2020 Feb 29];3. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4114002/>
11. Milanez VF, Pedro RE, Moreira A, Boullosa DA, Salle-Neto F, Nakamura FY. The role of aerobic fitness on session rating of perceived exertion in futsal players. *Int J Sports Physiol Perform*. 2011 Sep;6(3):358-66. doi:10.1123/ijsp.6.3.358 pmid:21911861
 12. Jones LA. Perceptual constancy and the perceived magnitude of muscle forces. *Exp Brain Res*. 2003 Jul;151(2):197-203. doi:10.1007/s00221-003-1434-4 pmid:12768260
 13. Pincivero DM, Coelho AJ, Campy RM. Perceived exertion and maximal quadriceps femoris muscle strength during dynamic knee extension exercise in young adult males and females. *Eur J Appl Physiol*. 2003 Apr;89(2):150-6. doi:10.1007/s00421-002-0768-0 pmid:12665978
 14. Row BS, Knutzen KM, Skogsberg NJ. Regulating explosive resistance training intensity using the rating of perceived exertion. *J Strength Cond Res*. 2012 Mar;26(3):664-71. doi:10.1519/JSC.0b013e31822ac367 pmid:22310518
 15. Sheard PW, Smith PM, Paine TJ. Athlete compliance to therapist requested contraction intensity during proprioceptive neuromuscular facilitation. *Man Ther*. 2009 Oct;14(5):539-43. doi:10.1016/j.math.2008.08.006 pmid:18996733
 16. Mehta RK, Agnew MJ. Subjective evaluation of physical and mental workload interactions across different muscle groups. *J Occup Environ Hyg*. 2015;12(1):62-8. doi:10.1080/15459624.2014.942455 pmid:25025738
 17. Pincivero DM, Gear WS. Quadriceps activation and perceived exertion during a high intensity, steady state contraction to failure. *Muscle Nerve*. 2000 Apr;23(4):514-20. doi:10.1002/(SICI)1097-4598(200004)23:43.0.CO;2-4
 18. Peñailillo L, Mackay K, Abbiss CR. Rating of Perceived Exertion During Concentric and Eccentric Cycling: Are We Measuring Effort or Exertion? *Int J Sports Physiol Perform*. 2018 April 1;13(4):517-23. doi:10.1123/ijsp.2017-0171 pmid:29035598
 19. Boat R, Atkins T, Davenport N, Cooper S. Prior self-control exertion and perceptions of pain and motivation during a physically effortful task. *Prog Brain Res*. 2018;240:19-34. doi:10.1016/bs.pbr.2018.08.007 pmid:30390831
 20. Morishita S, Tsubaki A, Nakamura M, Nashimoto S, Fu JB, Onishi H. Rating of perceived exertion on resistance training in elderly subjects. *Expert Rev Cardiovasc Ther*. 2019 Feb;17(2):135-42. doi:10.1080/14779072.2019.1561278 pmid:30569775
 21. Borg E, Borg G. A comparison of AME and CR100 for scaling perceived exertion. *Acta Psychol (Amst)*. 2002 Feb;109(2):157-75. doi:10.1016/S0001-6918(01)00055-5
 22. Lim W. Easy Method for Measuring Stretching Intensities in Real Clinical Settings and Effects of Different Stretching Intensities on Flexibility. *J Back Musculoskelet Rehabil*. 2019 July 23;32(4):579-85. doi:10.3233/BMR-181243 pmid:30530964
 23. Ament W, Verkerke GJ. Exercise and fatigue. *Sports Med*. 2009;39(5):389-422. doi:10.2165/00007256-200939050-00005 pmid:19402743
 24. Borg G. Psychophysical scaling with applications in physical work and the perception of exertion. *Scand J Work Environ Health*. 1990;16 Suppl 1:55-8. doi:10.5271/sjweh.1815 pmid:2345867
 25. Taylor JL, Gandevia SC. A comparison of central aspects of fatigue in submaximal and maximal voluntary contractions. *J Appl Physiol*. 2008 Feb;104(2):542-50. doi:10.1152/japplphysiol.01053.2007 pmid:18032577
 26. Barth JL, Holding DH, Stamford BA. Risk versus effort in the assessment of motor fatigue. *J Mot Behav*. 1976 Sep;8(3):189-94. doi:10.1080/00222895.1976.10735071 pmid:23964574
 27. Guo F, Sun Y-J, Zhang R-H. Perceived exertion during muscle fatigue as reflected in movement-related cortical potentials: an event-related potential study. *Neuroreport*. 2017 February 8;28(3):115-22. doi:10.1097/WNR.0000000000000732 pmid:28121808
 28. Vera J, Jiménez R, García JA, Cárdenas D. Simultaneous Physical and Mental Effort Alters Visual Function. *Optom Vis Sci*. 2017;94(8):797-806. doi:10.1097/OPX.0000000000001105 pmid:28708697
 29. Chang YK, Labban JD, Gapin JJ, Etnier JL. The effects of acute exercise on cognitive performance: a meta-analysis. *Brain Res*. 2012 May 9;1453:87-101. doi:10.1016/j.brainres.2012.02.068 pmid:22480735

30. Lambourne K, Tomporowski P. The effect of exercise-induced arousal on cognitive task performance: a meta-regression analysis. *Brain Res.* 2010 Jun 23;1341:12-24. [doi:10.1016/j.brainres.2010.03.091](#) [pmid:20381468](#)
31. Marcora S. Perception of effort during exercise is independent of afferent feedback from skeletal muscles, heart, and lungs. *J Appl Physiol.* 2009 Jun;106(6):2060-2. [doi:10.1152/jappphysiol.90378.2008](#) [pmid:18483166](#)
32. Mehta RK, Parasuraman R. Effects of mental fatigue on the development of physical fatigue: a neuroergonomic approach. *Hum Factors.* 2014 Jun;56(4):645-56. [doi:10.1177/0018720813507279](#) [pmid: 25029891](#)
33. Hallett M. Volitional control of movement: the physiology of free will. *Clin Neurophysiol.* 2007 Jun;118(6):1179-92. [doi:10.1016/j.clinph.2007.03.019](#) [pmid:17466580](#)
34. Pincivero DM, Coelho AJ, Campy RM, Salfetnikov Y, Bright A. The effects of voluntary contraction intensity and gender on perceived exertion during isokinetic quadriceps exercise. *Eur J Appl Physiol.* 2001 Mar;84(3):221-6. [doi:10.1007/s004210170008](#) [pmid:11320639](#)
35. Pincivero DM, Lephart SM, Moyna NM, Karunakara RG, Robertson RJ. Neuromuscular activation and RPE in the quadriceps at low and high isometric intensities. *Electromyogr Clin Neurophysiol.* 1999 Feb;39(1):43-8.
36. Lim W. Optimal Intensity of PNF Stretching: Maintaining the Efficacy of Stretching While Ensuring Its Safety. *J Phys Ther Sci.* 2018 Aug;30(8):1108-11. [doi:10.1589/jpts.30.1108](#) [pmid:30154610](#)
37. Lim W. Changes in Pain Following the Different Intensity of the Stretching and Types of Physical Stress. *Phys Ther Korea.* 2019 Nov;26(4):63-9. [doi:10.12674/ptk.2019.26.4.063](#)
38. Ekkekakis P, Parfitt G, Petruzzello SJ. The pleasure and displeasure people feel when they exercise at different intensities: decennial update and progress towards a tripartite rationale for exercise intensity prescription. *Sports Med.* 2011 August 1;41(8):641-71. [doi:10.2165/11590680-000000000-00000](#) [pmid:21780850](#)