



ORIGINAL ARTICLE

Association between Selected Body Composition, Blood Pressure, and Musculoskeletal Fitness in Nigerian Children

¹Sunday O. Onagbiye , ²Olumatoyin O. Toriola

¹The Africa Unit for Transdisciplinary Health Research (AUTHeR), North-West University, Potchefstroom Campus, South Africa. ²Prince Sultan University, Women Campus, Health and Physical Education Department, Riyadh, Kingdom of Saudi Arabia.

Submitted September 11, 2021; Accepted in final form November 28, 2021.

ABSTRACT

Background. Evidence shows that obesity negatively influences children's physical, physiological and musculoskeletal well-being. **Objectives.** This study examined the relationship between triponderal mass index (TMI), blood pressure (BP), and musculoskeletal fitness (MSF) in a cross-sectional sample of school children in Ado-Ekiti, Southwest Nigeria. **Methods.** Body weight, stature, systolic and diastolic blood pressures (SBP & DBP), and three components of musculoskeletal fitness, comprising sit and reach (SR), sit-up (SUP), standing broad jump (SBJ), and 20-meter shuttle runs, were assessed in 1229 (boys=483, 39.3%; girls=746, 60.7%) school children (age range: 9-13 years). **Results.** A significant positive correlation was found between TMI and SBP ($r=0.182$, $P<0.01$), and DBP ($r=0.090$, $P<0.01$), while TMI and SR ($r=-0.067$, $P<0.05$) and SBJ ($r=-0.246$, $P<0.01$) yielded a substantial inverse relationship. Children with severe underweight were significantly less probable to perform well in SBJ (OR= 0.96, CI= 0.94; 0.99) and 20-meter shuttle run (OR= 1.39, CI= 1.22; 1.61). Childhood overweight was associated with increased SBP (OR= 1.03, CI= 1.02; 1.05) and poor SBJ (OR= 0.98, CI= 0.97; 0.99). Obese children were significantly associated with increased SBP (OR= 1.04, CI= 1.02; 1.06), poor SR (OR= 0.92, CI= 0.87; 0.98) and SBJ (OR= 0.94, CI= 0.93; 0.96) scores compared to age- and sex-specific norms. **Conclusion.** Elevated BP and poor musculoskeletal fitness could adversely affect the well-being of undernourished, overweight, and obese Nigerian children. Findings implicate the need to implement dietary and physical activity intervention programs to promote desirable health-related quality of life in school-going children.

KEYWORDS: *Triponderal Mass Index, Musculoskeletal Fitness, Children, Sub-Saharan Africa.*

INTRODUCTION

Overweight and obesity in children are a public health challenge and should remain a national priority worldwide. The origin of cardiovascular disease (CVD) in adulthood, of which obesity remains one of its major risk factors, starts in childhood (1, 2). Obesity is strongly associated with major health risks such as hyperglycemia, asthma, arthritis, high BP, hyperlipidemia, and a faltering quality of life (3). Numerous studies have shown that there is a

connection between risk factors of CVD and poor musculoskeletal fitness in school children (1, 4). Several studies have also reported that any child with a highly active lifestyle and improved musculoskeletal fitness could prevent himself or herself from the future occurrence of any cardiovascular illness (5).

The two major anthropometric assessments most used to evaluate body fat in several age groups are body mass index (BMI) and tri-ponderal mass index

*. Corresponding Author:

Sunday O. Onagbiye, Ph.D.

E-mail: 24539678@nwu.ac.za

(TMI) (6, 7). However, BMI has been most extensively recognized and used as a body weightness valuation means among youths, while its association with physical or musculoskeletal fitness and performance has been widely investigated (8-11). However, it has been reported that BMI inherently has weaknesses in measuring fat mass percentage (7, 12). In contrast, recent studies have suggested that TMI could be a better prognosticator of health in children than BMI (6, 7, 12). Despite its usefulness in body conformation research, the connection between TMI, blood pressure, and musculoskeletal fitness in children has not been conclusively elucidated.

Manyeki et al. (13) reported that the relations between body mass index and musculoskeletal health in children in industrialized countries had been explored, but relatively tiny empirical evidence in this regard is obtainable in underdeveloped countries. In addition, studies examining the association between TMI, blood pressure, and musculoskeletal performance in children from sub-Saharan African countries, especially Nigeria, have rarely been conducted. Examining the relationship linking TMI, BP, and musculoskeletal fitness status in children is important from a public health standpoint. First, it could provide useful data for planning the health system due to excessive youth fatness (14). Second, the study's findings could facilitate early detection of children at risk of future cardiovascular and musculoskeletal dysfunction so that appropriate intervention could be instituted. Third, various health advantages were linked with musculoskeletal fitness. For example, it lowers risk factors for coronary diseases, lowers osteoporosis risk, improves elasticity, improves sugar forbearance, and help complete daily task without unnecessary fatigue (15). Finally, information on musculoskeletal fitness parameters, blood pressure, and TMI are usable markers of an individual's overall well-being, especially for youths. Hence, this study examined the association between TMI, blood pressure, and constituents of musculoskeletal fitness (flexibility: sit-and-reach test; muscular endurance: sit-up test; explosive leg power: standing broad jump, and shuttle run counts) in a sample of school children aged 9 to 13 years old in Ado-Ekiti, Southwestern Nigeria.

MATERIALS AND METHODS

Ethical Considerations. We sought authorization to carry out the Ekiti State Ministry

of Education study. It was followed by contacting the local and school authorities and the children's parents. An information leaflet containing the study's purpose, procedures, and where the study would take place was distributed to all the children and their guardians/parents or carers, who subsequently provided a written agreement form. Only children whose parents or guardians signed the consent forms were permitted to participate in the study. The children also gave verbal assent before the study commenced. The study was also executed following the ethical codes of the Helsinki pronouncement for investigations relating to human subjects (16).

The methodology of this study followed the approach used in Toriola et al. (8) study, where a cross-section research plan was utilized to gather data on TMI, blood pressure, and musculoskeletal health variables. These are flexibility (sit-and-reach test), muscular endurance (sit-up test), explosive leg power (standing broad jump), and shuttle run counts among 1229 (boys=483, 39.3% and girls=746, 60.7%) school children aged 9-13 years, haphazardly chosen from five municipal primary schools in Ado-Ekiti, Southwest, Nigeria. The children's ages were confirmed from their school register with the assistance of class teachers.

Measurements. Anthropometry. The practice of the International Society for the Advancement of Kinanthropometry was followed to assess height, body weight, and skinfolds. The skinfolds assessed include the triceps and subscapular (17). The participants' height was assessed to be close to 0.1 cm barefooted while the participants stood erect alongside a fixed stadiometer. We used a numerical weighing gauge named Tanita HD 309 manufactured by Creative Products, MI, USA, to assess body weight with participants flippantly dressed. Regular calibration of the weighing gauge close to 0.1 kg was performed after every 15 assessments. Based on the value derived from the participant's height and body weight, TMI was computed by apportioning weight (kg) by height cubed (m^3). Thus, TMI expressed as kg/m^3 poses a better assessment of plumpness in youths (12). The youngsters' TMI was afterward grouped as severe underweight, underweight, normal, overweight, and obese for age and gender using the age- and sex-specific TMI limit points (12). The waist-to-height ratio (WHR) was derived from the waist splitted by height in centimeters.

All data were collected by a group of trained field workers and documented using appropriate datasheets.

Blood Pressure. Blood pressure (BP) was assessed (in mm Hg - millimeters of mercury) with an automated BP sensor (Omron HEM - 705 CP devices, Tokyo, Japan). The regular procedures of the Blood Pressure Association (18) were applied for the children's BP assessment. We used the last two readings to calculate the average for systolic (SBP) and diastolic (DBP) measurements. The four BP categories as documented by the BP association are Low (90 over 60, i.e., 90/60 or less), Normal (More than 90 over 60 or 90/60) and less than 120 over 80, i.e., 120/80), pre-hypertension (More than 120 over 80 and less than 140 over 90 (120/80-140/90)), and hypertension (140 over 90, i.e., 140/90 or higher).

Musculoskeletal Fitness Assessment. We followed the protocol of the European test of physical fitness to assess health-related physical fitness variables (19). The tests were chosen based on their easiness of usability in a large number of participants. The test also provides dependable and useable health-related musculoskeletal and physical fitness assessments. Distinctively, the musculoskeletal fitness parameters that encompassed sit-and-reach, sit-up, standing broad jump, and 20-meter endurance shuttle-run (i.e., beep test) were performed as follows:

Sit-and-Reach Assessment. The SR tests were conducted to evaluate flexibility. We used a normal sit-and-reach wooden box. Participants progressively pushed the ruler on the topmost of the reach box by stretching their hands with no twitching and bent their trunks to lean forward as far as possible with the two hands upright and knees wholly stretched. This test was performed in a seated position with knees and feet positioned tightly alongside a flooring mat. During the test, the fingertips of the subject need to attain a similar expanse at the equivalent time with no leaping. The farthest gap touched in centimeters was noted after the test had been performed twice, i.e., an improved result was chosen and logged.

Sit-Up Assessment. Muscular strength was assessed using the sit-up (SUP) test. It was to examine the belly and hip flexor powers in 30 seconds of sequencing. The participants sat on a laid mat on the floor with their backs straight forward, and hands fastened at the rear of the neck. The participants' knees were also stretched

at an angle of 90° with cads and feet on the floor mat. The participant then laid with their back so that their shoulders contacted the mat on the floor and reverted to their inactive or sitting spot while their elbows were put forward to contact their knees. The assessments were completed when participants became exhausted to finish a full sit-up. Only the total number of appropriately executed performances was documented.

Standing Broad Jump Assessment. To assess the youth's explosive power, the SBJ test was used. The subjects stood at the back of the starting line, after which they were informed to get up and go energetically and hurdled horizontally as far as they could. The children were advised to land simultaneously with both feet and stand through the jump. The assessments were performed twice, with better scores documented. On the other hand, the score with the highest distance out of the two assessments was documented closed to 0.1 centimeters. The "distance between toes at take-off and heels at landing, or whichever body part landed nearest to the take-off point" was recorded.

20 Metres Shuttle Run. A sufficiently large flat and non-slippery space was determined with a measuring tape from which the 20m shuttle run track, which was subsequently demarcated with cones to set up a boundary, was marked. The participants were told to take a position at the rear of one of the lines opposite the subsequent mark, 20m apart, and start to sprint once instructed by the soundtrack, i.e., participants ran continuously between dual marks 20 meters away from each other in tune with soundtrack hoots. At each minute, the echo reflects an intensification in the speediness and the time interval between the beeps progressively shortens. If the subject gets to the mark before the beep soundtrack, he or she needs to slow down for the beep sounds to complete before continuing. Contrarily, if the subject does not get to the line prior to the sound of the hoot, the participant is served with caution and must sprint faster to meet up. The subject was served with caution the first time they could not get to the line and disqualified after the next caution. The weather was favorable, and the testers led the participants to 15 minutes of warming up exercise before the test.

Pilot Test. A pilot study involving 20 school children (male: n=10; female: n=10) of similar age categories was undertaken prior to data gathering to fine-tune the logistics of test administration and data

collection. Specifically, we conducted a training workshop that professionals led in Kinanthropometry. The workshop aimed to sharpen field workers' proficiency in assessing youths' kinanthropometric and musculoskeletal fitness parameters.

Statistical Analysis. Means and standard deviations of kinanthropometric and musculoskeletal aptness variables were analyzed. Deviations in the body weight, height, TMI, BP, SR, SUP, SBJ, and 20 meters shuttle run were estimated by gender using independent samples *t*-test. We performed a bivariate correlation analysis to examine the relationships between the children's TMI and musculoskeletal fitness variables and regression analysis to determine the association between TMI categories, BP, and musculoskeletal fitness among the study participants. We used the Statistical Package for the Social Sciences (SPSS) version 27.0 (SPSS Inc., 2020), while the alpha level was set at $p < 0.05$.

RESULTS

This study examined the association between TMI, blood pressure, and components of musculoskeletal health in youths aged nine to thirteen from Ado-Ekiti, Southwestern Nigeria. Presented in [Table 1](#) are the anthropometric, physiological, and musculoskeletal performance features of the children. There were significant gender differences in almost all the children's measurements, except for waist-to-height ratio and TMI. For example, the mean value for age in girls (11.8 years) was significantly higher than that of boys (11.6 years) ($P = 0.007$). Boys were significantly heavier (38.0 kg) ($P = 0.001$) and taller (144.8 cm) than girls ($P = 0.038$). Mean values of systolic (105.7 mmHg) and diastolic

(62.5 mmHg) BP were higher in boys compared to girls ($P < 0.001$). Furthermore, the mean flexibility score in boys (23.9 cm) was significantly higher when compared to that of girls (22.5 cm) ($P = 0.004$). In contrast, girls performed significantly better in SUP (21.8 no/min) ($P < 0.001$), SBJ (136.0 cm) ($P < 0.001$), and 20 meters shuttle run (4.51) ($P < 0.001$) than their male counterparts.

The results of the correlation analyses are provided in [Table 2](#). There was a significant positive correlation between the TMI and SBP ($r = 0.182$, $P < 0.01$) and DBP ($r = 0.090$, $P < 0.01$). Also, a significant inverse relationship was found between the children's TMI and SR ($r = -0.067$, $P < 0.05$) performances. Similarly, TMI was significantly correlated with SBJ ($r = -0.246$, $P < 0.01$). Furthermore, there was a relationship between the SBP and SR ($r = 0.079$, $P < 0.01$), SUP ($r = -0.077$, $P < 0.01$), and 20m shuttle run ($r = 0.154$, $P < 0.01$). Similar trends in the relationships were observed for DBP and SR ($r = 0.107$, $P < 0.01$) and ($r = 0.090$, $P < 0.01$) [Table 2](#).

Provided in [Table 3](#) are the regression coefficients on the association between TMI and Nigerian school children's musculoskeletal fitness. The children with severe underweight were significantly less likely to perform well in SBJ (OR= 0.96, CI= 0.94; 0.99) and SRCs (OR= 1.39, CI= 1.22; 1.61). Being overweight was significantly associated with poor SBJ (OR= 0.98, CI= 0.97; 0.99) performances compared with participants with normal body weight. Obese children were also significantly associated with poor flexibility (OR= 0.92, CI= 0.87; 0.98) and SBJ (OR= 0.94, CI= 0.93; 0.96) compared with the performances of children with normal weight ([Table 3](#)).

Table 1. Anthropometric, Physiological, and Musculoskeletal Fitness Characteristics of Nigerian Children

Variables	Total (n= 1229)	Male (n=483)	Female (n=746)	P- Value
	M±SD	M±SD	M±SD	
Age (y)	11.7±1.05	11.6±1.05	11.8±1.05	0.007
Body mass (kg)	37.1±7.60	38.0±8.14	36.5±7.17	0.001
Height (cm)	144.2±9.49	144.8±9.54	143.7±9.43	0.038
WHR	0.42±0.06	0.42±0.07	0.42±0.05	0.427
BMI (kg/m ²)	17.8±3.20	18.0±3.49	17.6±2.98	0.019
TMI (Kg/m ³)	12.4±2.89	12.5±3.13	12.3±2.73	0.184
SBP (mmHg)	102.5±15.6	105.7±15.1	100.5±15.7	0.000
DBP (mmHg)	60.4±11.5	62.5±11.4	58.9±11.4	0.000
SR (cm)	23.1±7.34	23.9±8.93	22.5±6.04	0.004
SUP (min)	20.1±7.34	17.5±6.86	21.8±7.13	0.000
SBJ (cm)	130.2±23.3	121.2±22.6	136.0±21.8	0.000
Shuttle run	4.32±2.16	4.02±2.16	4.51±2.14	0.000

Y, year; WHR, waist-to-height ratio; BMI, body mass index; BP, blood pressure; SR, Sit and Reach; SUP, sit-up; SBJ, standing broad jump; Kg, kilogram; Kg/m², kilogram per meter square; TMI, tri-ponderal index; Kg/m³, kilogram per meter cubic; mm, millimeter; M, mean; SD, standard deviation; SBP, Systolic blood pressure; DBP, diastolic blood pressure; mmHg, millimeter mercury; cm, centimeters; m, meters.

Table 2. Pearson's Correlation Matrix Showing Interrelationships between Triponderal Mass Index, Blood Pressure, and Musculoskeletal Fitness (n=1229)

Variables	Age (y)	Body mass (kg)	Height (cm)	WHtR	BMI (kg/m ²)	TMI (kg/m ³)	SBP (mmHg)	DBP (mmHg)	SR (cm)	SUP (sec)	SBJ (cm)	Shuttle run
Age (y)	1	0.297**	0.358**	-0.062*	0.084**	-0.035	0.138**	0.065*	-0.025	0.020	0.213*	0.064*
Body mass (kg)	0.297**	1	0.637**	0.193*	0.684**	0.361**	0.492**	0.318**	0.037	-0.095*	0.002	0.084**
Stature (cm)	0.358**	0.637**	1	-0.155**	-0.094**	-0.430**	0.319**	0.206**	0.070*	-0.117*	0.217*	0.102**
WHtR	-0.062*	0.193**	-0.155**	1	0.357**	0.350**	0.080**	0.164**	0.028	0.014	-0.082*	-0.298**
BMI (kg/m ²)	0.084**	0.684**	-0.094**	0.357**	1	0.924**	0.338**	0.198**	-0.034	0.002	-0.196*	0.034
TMI (kg/m ³)	-0.035	0.361**	-0.430**	0.350**	0.924**	1	0.182**	0.090**	-0.067*	0.054	-0.246*	0.011
SBP (mmHg)	0.138**	0.492**	0.319**	0.080*	0.338**	0.182**	1	0.576**	0.079*	-0.077*	-0.043	0.154**
DBP (mmHg)	0.065*	0.318**	0.206**	0.164*	0.198**	0.090**	0.576**	1	0.107*	-0.023	-0.004	0.090**
SR (cm)	-0.025	0.037	0.070*	0.028	-0.034	-0.067*	0.079**	0.107**	1	-0.025	0.119*	0.020
SUPs (sec)	0.020	-0.095**	-0.117**	0.014	0.002	0.054	-0.077**	-0.023	-0.025	1	0.275*	0.072*
SBJ (cm)	0.213**	0.002	0.217**	-0.082*	-0.196**	-0.246**	-0.043	-0.004	0.119*	0.275*	1	0.084**
20m shuttle run	0.064*	0.084**	0.102**	-0.298**	0.034	0.011	0.154**	0.090**	0.020	0.072*	0.084*	1

Y, year; WHtR, waist-to-height ratio; BMI, body mass index; BP, blood pressure; SR, Sit and Reach; SUP, sit-up; SBJ, standing broad jump; Kg, kilogram; Kg/m², kilogram per meter square; TMI, tri-ponderal index; Kg/m³, kilogram per meter cubic; mm, millimeter; M, mean; SD, standard deviation; SBP, Systolic blood pressure; DBP, diastolic blood pressure; mmHg, millimeter mercury; cm, centimeters; m, meters. **. Correlation is significant at the 0.01 level (2-tailed). *. Correlation is significant at the 0.05 level (2-tailed).

Table 3. Association between TMI and Musculoskeletal Fitness in Children

Musculoskeletal Variables	Sit & Reach (cm)	P-Value	Sit-Ups (sec)	P-Value	SBJ (cm)	P-Value	Shuttle Run	P-Value
	Adjusted OR (95% CI)		Adjusted OR (95% CI)		Adjusted OR (95% CI)		Adjusted OR (95% CI)	
TMI Categories								
Severe underweight	1.00 (0.92; 1.08)	0.93	0.99 (0.90; 1.08)	0.87	0.96 (0.94; 0.99)	0.008	1.39 (1.21; 1.60)	0.000
Underweight	0.98 (0.96; 1.00)	0.20	0.98 (0.96; 1.00)	0.07	1.00 (0.99; 1.01)	0.11	1.02 (0.95; 1.10)	0.52
Overweight	1.01 (0.98; 1.03)	0.38	0.98 (0.95; 1.01)	0.24	0.98 (0.97; 0.99)	0.007	1.04 (0.94; 1.14)	0.37
Obese	0.92 (0.87; 0.98)	0.009	1.03 (0.99; 1.08)	0.09	0.94 (0.93; 0.96)	0.000	1.12 (0.98; 1.28)	0.07
Normal weight	1		1		1		1	

Consequently, underweight children were significantly less likely to be hypertensive than those with normal weight (OR= 0.98, CI= 0.97; 0.99). Results also showed a significant association between overweight and SBP, which suggests that overweight school children have a

higher propensity of being hypertensive (OR= 1.03, CI= 1.01; 1.05) compared to those with normal weight. Similarly, obese children are more likely to be hypertensive (OR= 1.04, CI= 1.02; 1.07) compared to their peers with normal body weight (Table 4).

Table 4. Association between TMI Categories and Blood Pressure Categories in Children

TMI Categories	Severe Underweight	P-Value	Underweight	P-Value	Normal Weight	Overweight	P-Value	Obese	P-Value
	Adjusted OR (95% CI)		Adjusted OR (95% CI)			Adjusted OR (95% CI)		Adjusted OR (95% CI)	
BP Variables									
Systolic BP	1.01 (0.96; 1.06)	0.56	0.98 (0.97; 0.99)	0.008	ref	1.03 (1.01; 1.05)	0.000	1.04 (1.02; 1.07)	0.000
Diastolic BP	0.99 (0.93; 1.06)	0.95	0.98 (0.97; 1.00)	0.07	ref	1.00 (0.98; 1.02)	0.64	0.98 (0.95; 1.01)	0.21

Table 5. Association between Blood Pressure Categories and Musculoskeletal Fitness in Children

Blood Pressure categories	Low	P-Value	Normal	Systolic BP Pre-hypertensive	P-Value	Hypertensive	P-Value	Low	P-Value	Normal	Diastolic BP Pre-hypertensive	P-Value	Hypertensive	P-Value
	Adjusted OR (95% CI)			Adjusted OR (95% CI)		Adjusted OR (95% CI)		Adjusted OR (95% CI)			Adjusted OR (95% CI)		Adjusted OR (95% CI)	
Musculoskeletal fitness														
Sit & reach (cm)	0.965 (0.943; 0.988)	0.003	ref	1.014 (0.993; 1.035)	0.182	0.997 (0.889; 1.118)	0.953	0.964 (0.945; 0.983)	0.000	ref	0.996 (0.952; 1.042)	0.856	0.909 (0.813; 1.016)	0.092
Sit-ups (sec)	1.036 (1.016; 1.057)	0.000	ref	0.988 (0.963; 1.014)	0.368	0.991 (0.890; 1.104)	0.872	0.997 (0.981; 1.014)	0.741	ref	1.015 (0.970; 1.062)	0.526	0.947 (0.860; 1.043)	0.270
SBJ (cm)	1.001 (0.995; 1.008)	0.676	ref	1.001 (0.993; 1.009)	0.840	0.998 (0.965; 1.032)	0.917	1.002 (0.996; 1.007)	0.555	ref	0.989 (0.975; 1.003)	0.120	0.999 (0.970; 1.029)	0.940
Shuttle run	0.891 (0.826; 0.962)	0.000	ref	1.099 (1.024; 1.179)	0.000	1.076 (0.802; 1.443)	0.625	0.897 (0.848; 0.950)	0.000	ref	1.007 (0.875; 1.158)	0.927	0.876 (0.615; 1.249)	0.465

The regression analyses also showed a significant association between the children's SBP and musculoskeletal fitness performance categories. Those with low BP were less likely to perform well in SR (OR=.96, CI=.94; .98) and 20m shuttle run (OR=.89, CI=.82; .96). However, a child with low BP had a significantly higher predisposition to demonstrate superior sit-up performance (OR=1.03, CI=1.01; 1.05). It was also found that children with low BP were more susceptible to scoring high in SBJ, but the result was insignificant (OR=1.00, CI=.99; 1.00). Furthermore, compared to those with normal BP, pre-hypertensive children were more likely to perform well in SR (OR=1.09, CI=1.02; 1.17). For DBP, those with low BP were significantly less likely to exhibit good sit and reach (OR=.96, CI=.94; .98) and shuttle run scores (OR=.89, CI=.84; .95) compared to their normotensive counterparts (Table 5).

DISCUSSION

This study revealed significant relationships between TMI and musculoskeletal fitness in

Nigerian children. Since TMI is currently regarded as a better adiposity marker than BMI in children and adolescents (6, 7, 12), it is important to examine how it relates to BP and musculoskeletal fitness in children. To the best of our knowledge, this is one the few studies to examine how TMI relates to blood pressure and musculoskeletal fitness in Nigerian children. Consistent with the findings of a previous study, which reported that TMI is a better predictor of metabolic syndrome (MetS) in children than BMI (6), our findings showed that a positive relationship exists between TMI and BP. In a study that examined the regression of age-standardized weight for age-standardized height, for which the standardization was attained by articulating weight and height as portions of the fiftieth centile for age from an appropriate growth norm, it was reported that during puberty, a superior power than two is necessary (20), therefore making TMI to be a suitable obesity indicator (6) in predicting BP in youths.

The results of this study also displayed that TMI was significantly inversely related to SBJ.

Furthermore, a relationship was found between BP and musculoskeletal performance. For example, SBP was found to be positively correlated with the children's sit-and-reach, 20-meter shuttle run and inversely correlated with sit-ups performances, while DBP was positively correlated with sit-and-reach and 20-meters shuttle run. Several studies have reported inverse relationships between body conformation and physical fitness (2, 10, 13, 21-23). However, the direction of the relationship might not always be as expected. Although physical fitness could also be used interchangeably with musculoskeletal fitness, this depends on the parameters assessed. A study that examined the longitudinal association between muscular fitness and blood pressure to understand if changes in muscular fitness over twenty-four months were related with BP changes among youths found that muscular health during youth was inversely associated with SBP and DBP. Therefore, a small decrease in muscular capacity could increase cardiometabolic disease risk levels, DBP, and SBP inclusive (24).

Studies have revealed the association between several adiposity indicator categories and musculoskeletal performances in youths (8, 13, 25) but not TMI, which makes it difficult to compare our findings with other fat indicators such as BMI and waist circumference. In this study, the regression analysis of the association between the triponderal mass index and musculoskeletal performance showed that severely underweight individuals performed poorly in SBJ and SR. Similarly, excessively weighty persons may perform poorly in SBJ compared to participants with normal body weight. Being obese could also negatively influence children's flexibility and SBJ performances compared to similar performances in children with normal body weights. A Brazilian study that examined the relations between physical fitness and the anthropometrical and demographical determinants among children stated that unhealthy fitness status was associated with numerous features, such as being female, obesity, and extreme abdominal fat (25). Ajisafe et al. (22) also reported that an increased obesity rate among youths could trigger musculoskeletal incapability, which may indicate deficient motor skill aptitude. It is important to understand that obese persons have disparate physical fitness features compared to normal-weight individuals, which could result in poor running- and jumping-

related effort and strength (26). In addition, a negative association of excessive body fat with cardiorespiratory endurance, explosive muscle strength, and speed has been reported amongst Chinese youth (27).

Furthermore, our study findings based on the association between TMI classifications and BP showed that underweight children might have lesser odds of being hypertensive. Consequently, overweight and obese persons may tend to be hypertensive more than those with normal weight classification. Shim (28), who evaluated the allocation of triponderal mass index under age and sex, and the association of excessive fatness groups in accordance to sex- and age-specific triponderal mass index with MetS and its components, found that children in the overweight and obese groups had a higher propensity for increased BP compared with individuals having normal-weight.

Subsequently, studies have acknowledged the association between BP and musculoskeletal fitness among children (24, 29-31). Similarly, our study findings showed a significant association between BP and musculoskeletal fitness among the participants. Children with low systolic and diastolic BPs were less likely to have good sit-and-reach and 20m shuttle run performances. By contrast, children with low BP are more likely to exhibit better sit-ups. Furthermore, pre-hypertensive individuals were more likely to record superior shuttle run ability. However, this was not the same for the hypertensive children as they were less likely to perform well in sit and reach SUPs and SBJ but more likely to perform well in the 20m shuttle run. These findings are in harmony with those of the Agostinis-Sobrinho et al. (24) study, which also indicated that hypertensive individuals had a reduced capability of exhibiting good musculoskeletal performance. It suggests that youths' increased fitness score is associated with reduced SBP (31). Additionally, the risk of experiencing health problems could be associated with low fitness status among many individuals (30).

Since TMI has now been recognized as a more accurate adiposity evaluation tool, this study adds important information to the body of knowledge regarding the association between TMI, blood pressure, and musculoskeletal fitness among school-going children and adolescents in Nigeria. However, the study has some limitations. First, to date, there is currently no perfect or accurate adiposity evaluation

tool for children, except that the TMI is a more accurate index than BMI. These two adiposity prognosticators of health have shortcomings and will remain in use until a new and accurate evaluation tool for children is invented. Second, this study's musculoskeletal fitness tests are field tests and may not be as precise as laboratory-based protocols. Third, there is a dearth of data on the association between TMI, BP, and musculoskeletal fitness in children to which the present findings could be compared. It calls for the need for more studies to clarify the observed findings. Therefore, it is important that the present findings are not generalized and should be interpreted cautiously.

CONCLUSION

Triponderal mass index was inversely associated with BP and musculoskeletal fitness. It could adversely influence the health and well-being of undernourished, overweight, and obese children.

APPLICABLE REMARKS

- It is important that school feeding and physical activity programs be implemented to promote health-related quality of life in school-going children and adolescents.

REFERENCES

1. Reisberg K, Riso EM, Jurimae J. Physical fitness in preschool children in relation to later body composition at first grade in school. *PLoS One*. 2021;**16**(1):e0244603. doi: 10.1371/journal.pone.0244603 pmid: 33439898
2. Orntoft C, Larsen MN, Madsen M, Sandager L, Lundager I, Moller A, et al. Physical Fitness and Body Composition in 10-12-Year-Old Danish Children in Relation to Leisure-Time Club-Based Sporting Activities. *Biomed Res Int*. 2018;**2018**:9807569. doi: 10.1155/2018/9807569 pmid: 30687761
3. Raistenskis J, Sidlauskienė A, Strukcinskiene B, Ugur Baysal S, Buckus R. Physical activity and physical fitness in obese, overweight, and normal-weight children. *Turk J Med Sci*. 2016;**46**(2):443-450. doi: 10.3906/sag-1411-119 pmid: 27511509
4. Larsen MN, Nielsen CM, Orntoft CO, Randers MB, Manniche V, Hansen L, et al. Physical Fitness and Body Composition in 8-10-Year-Old Danish Children Are Associated With Sports Club Participation. *J Strength Cond Res*. 2017;**31**(12):3425-3434. doi: 10.1519/JSC.0000000000001952 pmid: 28445225
5. Esmaeilzadeh S, Ebadollahzadeh K. Physical fitness, physical activity and sedentary activities of 7 to 11 year old boys with different body mass indexes. *Asian J Sports Med*. 2012;**3**(2):105-112. doi: 10.5812/asjrm.34709 pmid: 22942996
6. Khoshhali M, Heidari-Beni M, Qorbani M, Motlagh ME, Ziaodini H, Heshmat R, et al. Tri-ponderal mass index and body mass index in prediction of pediatric metabolic syndrome: the CASPIAN-V study. *Arch Endocrinol Metab*. 2020;**64**(2):171-178. doi: 10.20945/2359-3997000000206 pmid: 32236304
7. De Lorenzo A, Romano L, Di Renzo L, Gualtieri P, Salimei C, Carrano E, et al. Triponderal mass index rather than body mass index: An indicator of high adiposity in Italian children and adolescents. *Nutrition*. 2019;**60**:41-47. doi: 10.1016/j.nut.2018.09.007 pmid: 30529185
8. Toriola A, Monyeki M, Ajayi-Vincent O, Oyeniyi P, Akindutire I. Relationship between body composition and musculoskeletal fitness in Nigerian children. *Asian J Sci Res*. 2018;**11**:169-176. doi: 10.3923/ajsr.2018.169.176

ACKNOWLEDGMENTS

The author would like to thank all those who contributed in various ways and the schoolteachers for their assistance in the coordination of the participants of this study. Appreciation also goes to the guardians/parents and all the children who partook in this study.

AUTHORS' CONTRIBUTION

Study concept and design: Sunday O. Onagbiye. Acquisition of data: Sunday O. Onagbiye & Olumatoyin O. Toriola. Analysis and interpretation of data: Sunday O. Onagbiye. Drafting of the manuscript: Sunday O. Onagbiye & Olumatoyin O. Toriola. Critical revision of the manuscript for important intellectual content: Sunday O. Onagbiye & Olumatoyin O. Toriola. Statistical analysis: Sunday O. Onagbiye. Administrative, technical, and material support: Sunday O. Onagbiye & Olumatoyin O. Toriola. Study supervision: Sunday O. Onagbiye & Olumatoyin O. Toriola.

CONFLICT OF INTEREST

The authors mention that there is no "Conflict of Interest" in this study.

9. Casonatto J, Fernandes RA, Batista MB, Cyrino ES, Coelho ESMJ, de Arruda M, et al. association between health-related physical fitness and body mass index status in children. *J Child Health Care*. 2016;**20**(3):294-303. doi: 10.1177/1367493515598645 pmid: 26396021
10. Huang YC, Malina RM. Body mass index and individual physical fitness tests in Taiwanese youth aged 9-18 years. *Int J Pediatr Obes*. 2010;**5**(5):404-411. doi: 10.3109/17477160903497902 pmid: 20429737
11. Esco MR, Olson MS, Williford H. Relationship of push-ups and sit-ups tests to selected anthropometric variables and performance results: a multiple regression study. *J Strength Cond Res*. 2008;**22**(6):1862-1868. doi: 10.1519/JSC.0b013e318181fd03 pmid: 18824933
12. Peterson CM, Su H, Thomas DM, Heo M, Golnabi AH, Pietrobelli A, et al. Tri-Ponderal Mass Index vs Body Mass Index in Estimating Body Fat During Adolescence. *JAMA Pediatr*. 2017;**171**(7):629-636. doi: 10.1001/jamapediatrics.2017.0460 pmid: 28505241
13. Monyeki MA, Koppes LL, Kemper HC, Monyeki KD, Toriola AL, Pienaar AE, et al. Body composition and physical fitness of undernourished South African rural primary school children. *Eur J Clin Nutr*. 2005;**59**(7):877-883. doi: 10.1038/sj.ejcn.1602153 pmid: 15915157
14. World Health Organisation (WHO). Mapping the health system response to childhood obesity in the WHO European Region - An overview and country perspectives. Copenhagen, Denmark: 2019 WHO.
15. Kell RT, Bell G, Quinney A. Musculoskeletal fitness, health outcomes and quality of life. *Sports Med*. 2001;**31**(12):863-873. doi: 10.2165/00007256-200131120-00003 pmid: 11665913
16. World Medical A. World Medical Association Declaration of Helsinki: ethical principles for medical research involving human subjects. *JAMA*. 2013;**310**(20):2191-2194. doi: 10.1001/jama.2013.281053 pmid: 24141714
17. The International Society for the Advancement of Kinanthropometry (ISAK). International Standard for Anthropometric Assessment. Potchefstroom, South Africa: ISAK2006.
18. Blood Pressure Association. What do the number mean? [updated 2020 June 16 cited 2021 July 2]. Available from: <http://www.bloodpressureuk.org/your-blood-pressure/understanding-your-blood-pressure/what-do-the-numbers-mean/>.
19. Council of Europe. EUROFIT: Handbook for the EUROFIT Tests of Physical Fitness. Strasbourg: Sports Division Strasbourg, Council of Europe Publishing and Documentation Service1993.
20. Cole TJ. A method for assessing age-standardized weight-for-height in children seen cross-sectionally. *Ann Hum Biol*. 1979;**6**(3):249-268. doi: 10.1080/03014467900007252 pmid: 496386
21. Mak KK, Ho SY, Lo WS, Thomas GN, McManus AM, Day JR, et al. Health-related physical fitness and weight status in Hong Kong adolescents. *BMC Public Health*. 2010;**10**:88. doi: 10.1186/1471-2458-10-88 pmid: 20178615
22. Ajisafe T, Garcia T, Fanchiang HC. Musculoskeletal Fitness Measures Are Not Created Equal: An Assessment of School Children in Corpus Christi, Texas. *Front Public Health*. 2018;**6**:142. doi: 10.3389/fpubh.2018.00142 pmid: 29872650
23. Monyeki MA, Neetens R, Moss SJ, Twisk J. The relationship between body composition and physical fitness in 14 year old adolescents residing within the Tlokwe local municipality, South Africa: the PAHL study. *BMC Public Health*. 2012;**12**:374. doi: 10.1186/1471-2458-12-374 pmid: 22626033
24. Agostinis-Sobrinho C, Ruiz JR, Moreira C, Lopes L, Ramirez-Velez R, Garcia-Hermoso A, et al. Changes in muscular fitness and its association with blood pressure in adolescents. *Eur J Pediatr*. 2018;**177**(7):1101-1109. doi: 10.1007/s00431-018-3164-4 pmid: 29740692
25. Andreasi V, Michelin E, Rinaldi AE, Burini RC. Physical fitness and associations with anthropometric measurements in 7 to 15-year-old school children. *J Pediatr (Rio J)*. 2010;**86**(6):497-502. doi: 10.2223/JPED.2041 pmid: 21076797
26. Kitabayashi T, Demura SI, Noguchi T. Physical Fitness Characterization by Obesity Level in Young Males with Poor Physical Fitness. *Advanc Physic Educat*. 2013;**3**(2):76-79. doi: 10.4236/ape.2013.32012
27. Shang X, Liu A, Li Y, Hu X, Du L, Ma J, et al. The Association of Weight Status with Physical Fitness among Chinese Children. *Int J Pediatr*. 2010;**2010**:515414. doi: 10.1155/2010/515414 pmid: 20652083
28. Shim YS. The Relationship Between Tri-ponderal Mass Index and Metabolic Syndrome and Its Components in Youth Aged 10-20 Years. *Sci Rep*. 2019;**9**(1):14462. doi: 10.1038/s41598-019-50987-3 pmid: 31594996

29. Pate RR, Daniels S. Institute of Medicine report on fitness measures and health outcomes in youth. *JAMA Pediatr.* 2013;**167**(3):221-222. doi: [10.1001/jamapediatrics.2013.1464](https://doi.org/10.1001/jamapediatrics.2013.1464) pmid: 23358923
30. Ramirez-Zea M, Melgar P, Flores R, Hoddinott J, Ramakrishnan U, Stein AD. Physical fitness, body composition, blood pressure, and blood metabolic profile among young Guatemalan adults. *Food Nutr Bull.* 2005;**26**(2 Suppl 1):S88-97. doi: [10.1177/15648265050262S109](https://doi.org/10.1177/15648265050262S109) pmid: 16060215
31. Fraser GE, Phillips RL, Harris R. Physical fitness and blood pressure in school children. *Circulation.* 1983;**67**(2):405-412. doi: [10.1161/01.cir.67.2.405](https://doi.org/10.1161/01.cir.67.2.405) pmid: 6848231