










REVIEW ARTICLE



Impact of Various Types of Exercise on Lipid Metabolism in Patients with Type 2 Diabetes and Concurrent Overweight/Obesity: A Narrative Review

¹Sameer Badri Al-mhanna *, ¹Wan Syaheedah Wan Ghazali *, ²Alexios Batrakoulis , ³Nouf H. Alkhamees , ⁴Clemens Drenowatz , ¹Mahaneem Mohamed , ⁵Mehmet Güllü , ⁶Hafeez Abiola Afolabi , ⁷Georgian Badicu 

¹Department of Physiology, School of Medical Sciences, Universiti Sains Malaysia, Kubang Kerian, Kelantan, Malaysia. ²Department of Physical Education and Sport Science, School of Physical Education, Sport Science and Dietetics, University of Thessaly, Trikala, Greece. ³Department of Rehabilitation College of Health and Rehabilitation Sciences, Princess Nourah bint Abdulrahman University, Riyadh, Kingdom of Saudi Arabia. ⁴Division of Sport, Physical Activity and Health, University of Teacher Education Upper Austria, Linz, Austria. ⁵Department of Sports Management, Faculty of Sport Sciences, Kirikkale University, Kirikkale, Turkey. ⁶Department of General Surgery, School of Medical Sciences, Universiti Sains Malaysia, Kubang Kerian, Kelantan, Malaysia. ⁷Department of Physical Education and Special Motricity, Transilvania University of Brasov, Brasov, Romania.

Submitted December 08, 2023; Accepted in final form February 04, 2024.

ABSTRACT

Background. Type 2 diabetes mellitus (T2DM) and obesity present substantial challenges to global public health, marked by their widespread prevalence and associated morbidities. The ensuing complications, including cardiovascular disease (CVD), metabolic syndrome, cancer, liver disease, and neurodegeneration, underscore the urgent need for effective preventive measures. Despite this, primary prevention of CVD in individuals with T2DM and obesity remains inadequate. Regular exercise emerges as a pivotal factor in ameliorating various cardiometabolic parameters, yet conflicting findings persist regarding the impact of exercise parameters (frequency, intensity, time, and type) on lipid homeostasis. **Objectives.** This review endeavors to scrutinize the effects of diverse exercise types and parameters on individuals with T2DM and concurrent obesity. A focus is placed on investigating the influence of exercise on conventional lipids, such as LDL-C, HDL-C, TG, total TC, and VLDL-C. Additionally, the review briefly delves into the mechanisms underlying exercise-induced effects on lipids and lipoproteins. **Methods.** A literature search was conducted using the PubMed, Scopus, and Google Scholar databases. **Results.** Regular exercise proves instrumental in elevating HDL-C levels while concurrently reducing TG, TC, VLDL, and LDL-C. Notably, exercise mitigates CVD risk, lowers BMI, and enhances insulin resistance, contingent on exercise types, volume, intensity, frequency, and duration. **Conclusion.** Future research must delve into the dose-response effects of real-world exercise programs to guide tailored interventions. This comprehensive understanding should inform clinicians and practitioners, empowering them to prescribe personalized exercise regimens for individuals grappling with compromised metabolic health.

KEYWORDS: *Aerobic Exercise, Resistance Training, Physical Activity, Metabolic Syndrome.*

*. Corresponding Author:

Sameer Badri Al-mhanna, Ph.D.

E-mail: Sameer@student.usm.my

Wan Syaheedah Wan Ghazali, Ph.D.

E-mail: syaheeda@usm.my

INTRODUCTION

Diabetes mellitus (DM) is a 21st-century global public health concern, as shown by the high incidence of associated morbidities and complications (1, 2). The global prevalence of diabetes among individuals aged 20 to 79 was approximately 10.5% (536.6 million people) in 2021, and it is anticipated to rise to 12.2% (783.2 million) by 2045 (3). The prevalence of diabetes was comparable between men and women, with the highest rates observed in the 75–79 age group. In 2021, urban areas had a higher estimated prevalence of 12.1% compared to 8.3% in rural areas, and high-income countries had a prevalence of 11.1% compared to 5.5% in low-income countries. The most significant relative increase in diabetes prevalence between 2021 and 2045 is expected in middle-income countries at 21.1%, surpassing projections for high-income (12.2%) and low-income (11.9%) countries. In terms of global diabetes-related health expenditures, estimates reached 966 billion USD in 2021 and are projected to reach 1,054 billion USD by 2045 (3).

The Type 2 diabetes mellitus (T2DM) physiopathway is linked to reduced high-density lipoprotein cholesterol (HDL-C), increased density of low-density lipoprotein (LDL) particles, elevated triglycerides and a cluster of interrelated plasma lipid and lipoprotein disorders (4). These are correlated with a significant part of the insulin resistance syndrome, which is linked to T2DM. In general, pre-diabetic people also have an atherogenic trend of risk factors, such as higher total cholesterol (TC), low-density lipoprotein cholesterol (LDL-C), and TG, as well as lower HDL-C, than people who do not experience diabetes (5, 6). Insulin resistance has a serious influence on very low-density lipoprotein (VLDL), LDL-C, and high-density lipoprotein (HDL) particle size and concentrations (7). In addition, obesity is known as a global epidemic that has a significant impact on population health (8), because of its association with T2DM and metabolic syndrome. Obesity and overweight affect more than 1.9 billion individuals worldwide, contributing to an increase in various health risks that include insulin resistance, type 2 diabetes, cardiovascular disease, cancer, liver disease, and neurodegeneration (9). Obesity, specifically central obesity, might be associated with insulin resistance, elevated triglycerides, and low HDLC; which is known as metabolic syndrome (10).

It is well-known that obese diabetic patients have an increased risk of CVD, but unfortunately, in diabetic patients, primary CVD prevention remains insufficient (11). Therefore, engagement in regular exercise is associated with reducing the risk of CVD, improving the lipid profile, and providing many benefits on metabolic syndrome and its risk factors such as abdominal obesity, hypertension, and insulin resistance (12). Bharath, Choi (13), for example, showed a reduced risk of metabolic syndrome, such as central obesity, high blood pressure (14), dyslipidemia, and insulin resistance following combined aerobic and resistance training (15).

In fact, exercise for improving overall health among people affected by obesity and T2DM has been recently reported as one of the most popular reasons for exercise worldwide (16). Ideally, a combination of aerobic exercise (AE) and diet has been documented as the optimal strategy (17, 18). Despite the well-characterized health benefits of exercise on lipid and lipoprotein profile, there remains some controversy surrounding the optimal exercise modality, duration, frequency, and/or intensity amongst DM patients with obesity. Since, the improvement of the lipid profile as a result of exercise depends on the intensity, frequency, and duration of each session as well as the type of exercise and the time spent on such a program there have been conflicting results on the efficacy of exercise DM patients with obesity (19). Hence, the purpose of this study was to examine the effects of different types of exercise, duration, frequency, and/or intensity for DM patients with obesity. Furthermore, this review aimed to provide information on the effects of various exercise training modalities on both conventional lipids, such as LD, high-density lipoprotein (HDL), triglycerides TG, cholesterol, very-low-density lipoprotein (VLDL), and lipoproteins.

Exercise training can be broadly categorized into three main types: AE, resistance exercise (RE), and concurrent training.

AEROBIC EXERCISE

AE is defined as any activity that utilizes large muscle groups, can be sustained over a prolonged period of time, is rhythmic, and stimulates muscles using aerobic metabolism to derive energy in the form of adenosine triphosphate (ATP) from amino acids, fatty acids, and

carbohydrates. Cycling, dancing, jogging, diving, and walking are forms of aerobic activities (20).

The effectiveness of AE in improving lipid profiles among type overweight/ obese T2DM patients can be attributed to several mechanisms. These exercises have been found to positively affect lipid metabolism, improving lipid profile parameters such as TC, LDL-c, HDL-C, and TG (17).

AE has been shown to enhance insulin sensitivity in individuals with T2DM. Improved insulin sensitivity helps to regulate blood glucose levels and reduces the production of TG and VLDL in the liver, leading to favorable changes in the lipid profile. AE can potentially reduce visceral fat stored around the organs in the abdominal area. Visceral fat is metabolically active and contributes to increased TG and LDL-C levels, often called “bad cholesterol”. Exercise can improve the lipid profile by reducing visceral fat by decreasing TG and LDL-C levels (21, 22). Regular exercise, including AE, has increased HDL-C levels, commonly known as “good cholesterol”. HDL-C helps transport excess cholesterol from peripheral tissues to the liver for excretion, thus reducing the risk of cardiovascular disease. Furthermore, exercise can modify the size and density of LDL-C particles, making them less atherogenic (less likely to cause plaque buildup in the arteries) (23).

The mechanism responsible for the acute rise in HDL-C following exercise is likely associated with the catabolism of lipoproteins rich in TG through lipoprotein lipase (LPL) (24). Increased activity of LPL following exercise results in hydrolysis of chylomicron and VLDL, thereby decreasing plasma TG levels (25), and subsequently, surface remnants are converted into nascent HDL-C; nevertheless, following the exercise, the levels of HDL-C can increase. In comparison, drops in plasma TG levels following acute activity tend to correlate with the amount of intensity and duration of the EX (25). Increased LPL activity leads to a reduction in circulating TG and an improvement in the lipid profile. Studies showed that exercise could induce a decrease in TG levels within 24 hours (26, 27), while shorter workout durations will lead to only moderate reductions in plasma TG in obese women (28).

On the other hand, obese T2DM patients are associated with chronic low-grade inflammation, which contributes to dyslipidemia and cardiovascular risk. AE exercise has been shown

to reduce inflammation markers such as C-reactive protein (CRP) and interleukin-6 (IL-6). By decreasing inflammation, exercise can positively influence the lipid profile and reduce the risk of cardiovascular complications (29, 30). Moreover, AE primarily utilize fat as fuel for exercise. By increasing the intensity and duration of exercise, patients can enhance fat oxidation, reducing overall body fat and improving their lipid profile (31-33).

Lipoproteins. CVD is the main cause of death around the world, with hypertension, smoking, physical inactivity, obesity, elevated glucose levels/ DM mellitus, and dyslipidemia being major CVD risk factors (30). Among these, dyslipidemia which is described by a low level of HDL-C is closely and contrarily linked to the risk of cardiovascular disease (34). Also, low levels of HDL-C are part of the atherogenic dyslipidemia complex that is linked to T2DM (35). Increased atherogenesis has been described in DM individuals due to many abnormalities in lipoprotein metabolism, particularly in VLDL, LDL-C, and HDL-C (36).

Some studies indicated that HDL-C efflux capacity was a stronger predictor for the prevalence and incidence of CHD, and HDL-C efflux capacity was also revealed to be diminished in obese patients with T2DM (37, 38). Furthermore, it was recently found that in overweight adults-only those with an elevated TG and reduced HDL-C levels had hypertension, increased C-reactive protein (CRP) levels, and insulin resistance (39). These findings triggered a surge of interest in raising HDL-C levels in obese T2DM patients by encouraging participation in AE as a CVD therapy and to reduce insulin resistance.

It was reported that AE is a preventive strategy against atherosclerosis and has been proposed as medical assistance in the treatment of DM patients, which can help bring high blood glucose levels down into the normal range by improving insulin sensitivity and minimizing the risk of CVD (40, 41). An AE program for 3 months, with each session consisting of a 10-min warm-up, 30-40 min of walking, cycling, or jogging, or a combination at 60% to 65% of maximum oxygen consumption (VO₂max), and a 10-min cool-down, once or twice a week showed a significant enhancement in lipid and lipoprotein profiles and a rise in HDL-C levels, while LDL-C and TG decreased in type 2 diabetic patients (42).

Low-density lipoprotein. Individuals who commenced the training program with lower HDL-C/LDL-C ratios exhibited the most significant improvements after engaging in exercise, emphasizing positive enhancements in those who are at the highest risk, which holds clinical relevance for their overall health and well-being (43).

Azizi et al., (2019) (44) conducted a study in twenty-four middle-aged adults (40-50 years, 44.0 ± 2.3 mean \pm SD) with T2DM, subjected to 35-50 min of AE at 40-55% of their heart rate reserve (HRR) for 3 days a week over 8 weeks and reported a decline in TC (33.2%), TG (55.9%) and LDL-C (30.0%) in the training group while HDL-C (38.5%) was higher after 8 weeks of AE compared to the control group. In contrast to the control group, the training group's fasting blood sugar, fasting insulin, and insulin resistance index all decreased.

While another study found that 5 months of daily cycling exercise for 20 min did not affect LDL-C and other main lipid profile indices in elderly women with diabetes, these participants showed a minor increase in VO₂ max (45). According to an observational study comparing an endurance training group with an inactive control, the endurance training group displayed significantly lower LDL-C density with no variations in the mass of the larger LDL-C and IDL particles (46). Since endurance training lowers LDL-C particle mass, the chance of coronary artery disease (CAD) should be lower in people with T2DM. However, endurance training in men did not result in any changes in these parameters over a one year period (47, 48). Hence, the effects of 4 weeks of AE on lipid profile parameters, HDL-C, LDL-C, and chylomicrons in Type II diabetic patients were not significantly different after the intervention (49).

Very low-density lipoprotein. Diabetic patients are more likely to develop dyslipidemia, which is one of the leading causes of CVD (50). One mechanism explaining this association is elevated free fatty acid (FFA) production in insulin-resistant fat cells. High levels of FFA boost TG production, which in turn enhances apolipoprotein B (ApoB) and VLDL cholesterol secretion. High levels of ApoB and VLDL are linked to an elevated risk of CVD in obese and DM patients (50, 51). In addition, hyperglycemia may also have a detrimental effect on lipoproteins

(especially LDL and VLDL) by raising glycosylation and oxidation, reducing vascular compliance, and promoting the progression of atherosclerosis (52). It has been reported that engagement in an AE program is one of the most important strategies to improve VLDL when compared to inactive people. Physically active adults also had higher HDL-C levels and lower TG, LDL-C, and VLDL levels, as well as a decreased CVD risk (53).

According to Turcotte and Fisher (2008) (54), AE can decrease unfavorable lipid and lipoprotein profiles, and these positive changes are not affected by gender, body weight, or diet; however, they can be affected by the degree of glucose tolerance (25). Alam et al., (2004) (55) stated that 6 months of AE in moderately obese individuals with T2DM resulted in a substantial reduction in VLDL level. Twenty sedentary males and females with T2DM completed 30 minutes of AE 4 times a week. Specifically, they performed 10 min of stepping on a 9-inch stepper in addition to 20 min of spinning at a resistance of 5, which resulted in a substantial decline in VLDL and TG parameters (49). This reduction can be attributed to enhanced TG catabolism, reduced hepatic synthesis and secretion, or a combination of the two because of increased catabolism of VLDL-triglycerides. There is also evidence that hepatic synthesis is decreased. Another study, on the other hand, did not show a substantial reduction in VLDL-C levels following AE (42). The inconsistent findings may be attributed to variations in study populations, exercise protocols, and individual responses to AE. Factors such as genetic differences, baseline fitness levels, and dietary habits could influence the impact of AE on VLDL-C levels.

Several studies also demonstrated that AE might have no effect on fasting blood LDL-C levels unless there was a reduction in body weight (56, 57). While, a recent meta-analysis conducted among overweight and obese adults by Kelley et al., (2005) (57), which included 13 studies with a total of 613 overweight and obese adults performed AE, showed a nonsignificant reduction in blood LDL-C concentration of less than 1%, independent of body weight. Changes in blood LDL-C however, were shown to have a significant association with body weight after conducting sensitivity analyses. Furthermore, AE was associated with a significant decrease in LDL-C in conjunction with significant weight

loss (58). Another study also showed that each kilogram of weight loss resulted in a drop of roughly 0.8-mg/dL-1 in LDL-C (59).

High-density lipoprotein. Obesity/overweight is a leading cause of HDL-C dysfunction because it lowers HDL-C levels (4). The impact of various forms of exercise on HDL-C function is complicated and relies on several factors, including exercise and individual features, but the effects of regular AE on these measures are unknown (4). There is, however, strong evidence that exercise, especially aerobic exercise performed in clinical practice, is more effective in the management of obese patients with T2DM than resistance exercise (RE) (60).

A recent randomized controlled trial conducted among 102 patients of both genders aged 40 to 70 years with T2DM completed an AE program for 25 weeks with exercise sessions on 3 days a week resulting in a total of 150 M per week, reported a substantial raise in HDL-C level compared to the control group (61). On the other hand, Nassef et al., (2019) (62) reported in a large study with almost 8,000 participants between 30 and 70 years of age that AE and daily badminton were linked to higher levels of HDL-C following participation in both AE and regular badminton. Another study, however, found that moderate AE for 2 days a week over a period of 12 months did not substantially improve cardiometabolic markers including TC, LDL-c, and CRP levels in obese adults following the exercise intervention. Nevertheless, AE was associated with reduced HDL-C in overweight and obese persons after 6 and 12 months of exercise engagement (19).

In summary, AE of different intensities and durations has been associated with an increase (63, 64) decrease (65), or no change (66) in HDL-C levels in overweight and obese adults. Such inconsistencies in the recorded efficacy of AE on HDL-C were due to discrepancies in experimental design, e.g. not all experiments were controlled for collateral changes in diet, alcohol consumption, and smoking, all of which could significantly modify lipoprotein profiles. Moreover, the time of plasma processing may have a profound influence on levels of HDL-C (67).

Triglycerides. In obese patients with T2DM, daily AE has been linked to lower plasma TG concentrations, lower BMI, and improved insulin resistance. A study performed in India between September 2011 and June 2012 demonstrated a

substantial reduction in total TG in 22 obese adults without T2DM who exercised moderately for 40 min/day for 5 days/week over a period of 15 weeks (68). Hayashino et al., (2012) (69) indicated that HDL-C, LDL-C, and TC levels were significantly improved in repose to AE. In addition, a total of 9 randomized controlled trials with 619 obese or overweight adults showed significantly decreased TG levels following AE even though there was no effect on other lipid and glucose indices (70).

While many studies showed that, AE did not alter TG levels following a single exercise session (71). Kelley and Kelley (2007) (72) reported that LDL-C was decreased in response to endurance training, even though these exercises had no significant effects on HDL-C, TC, and TG. It was documented that endurance training typically decreases plasma TG concentrations (73, 74) when baseline concentrations were elevated, but there were also exceptions (75).

With aerobic activity, TG was not improved substantially in inactive people. However, the findings of studies in obese with T2DM have been very promising. Several studies noted that when participants' TG levels were lower at the onset, there was only a small reduction in TG during exercise (43). Whereas, when TG baseline concentration was elevated, there was a significant reduction (43). Hence, TG levels at baseline appear to be a major determinant on the impact of exercise on TG response.

Several studies have shown that lower TG levels were often followed by no change in HDL-C, or that increasing HDL-C was accompanied by no noticeable change in TG levels (43). Similarly, an increase in HDL-C was not associated with TG levels in previously inactive people. However, after AE training, sedentary participants with hyperlipidemia displayed an increase in HDL-C levels while their TG levels decreased (76). Grandjean et al., (2000) (77) documented that after two days of AE, TG levels declined, with a 14% increase in HDL-C. This rise in HDL-C concentrations might be attributed to an increase of almost 11% in HDL3-C levels. Researchers are still looking for explanations for the diverse responses of TG and HDL-C to AE. Bodyweight, body fat, cardiovascular health, exercise status, geographic lipid concentrations, lifestyle changes, and genetic factors all tend to play a part. Additionally, exercise duration, blood collecting time, and subject sample size are all factors to be considered.

Total cholesterol. A randomized controlled trial including 28 young men with DM aged 20–40 who participated in a 12 to 16-week AE program for 3 to 5 days/week with 30 to 60 min of moderate-intensity running, found a significant reduction in Serum TC following AE compared to the control group (78). A 16-week program of 60 min mixed and AE three times a week, showed a decrease in TC without effects on HDL-C, TG, body weight, and glycemic control in diabetic patients (79).

On the other hand, it was reported that diabetic obese patients with nonalcoholic fatty liver disease following an AE program at 60% to 70% VO₂max for 3 sessions/week over 8 week period exhibited a significant decrease in TC, TG, LDL-C, and increase HDL-C level (80). Gondim et al., (2015) (19), however, reported no significant improvement in TC in overweight or obese participants following an exercise intervention twice a week over a period of 6 to 12 months. A meta-analysis of randomized controlled trials implementing AE over periods of 8 weeks or more in T2DM patients, included seven studies with a total of 220 male and female participants. Results showed a substantial reduction of nearly 5% in LDL-C, while TC, HDL-C, TC/HDL-C, and TG did not change significantly (72). Furthermore, AE training consisting of 75-minute sessions for 2 days/week over 12 weeks did not affect TC in 26 patients with T2DM (81). TC, LDL-C, and triglyceride levels, however, did change after 21 weeks of high-intensity exercise (82). These inconsistent results may be attributed to differences in the duration of the training session, the amount of exercise accomplished, changes in body composition, nutritional intake, weight loss, and pre-training TG concentration across studies.

RESISTANCE EXERCISE

Resistance training, alternatively referred to as strength training or weight training, involves engaging in exercises that test the muscles against an external force. The primary objective of resistance training is to enhance muscle strength, endurance, and power (83). RT, which involves powerlifting, dumbbell, and barbell training, allows muscles to contract against external resistance to improve strength and mass (84).

Lipoproteins. As a result of the growing scientific evidence of RT's benefits additional

evidence on the advantages of RT has been added since it was addressed in the guidelines of the American College of Sports Medicine (ACSM) (85). According to a systematic review and meta-analysis, RT can result in enhanced levels of coordination and muscle strength/power in obese older adults (86). Tayebi et al., (2023) (83) found that engaging in eight weeks of circuit resistance training, with low to high intensities, can lead to elevated resting levels of METRNL in men diagnosed with T2DM. This increase in METRNL is notably correlated with enhancements in fasting blood glucose levels and a reduction in insulin resistance.

Pollock et al., (2000) (87) also indicated that RT, implemented at least twice a week is a healthy and effective way to improve muscle strength and endurance by 25% to 100% or more in individuals with and without cardiovascular disease. Studies further demonstrated that when adequately administered and controlled, complementary RT can increase muscle strength and endurance, cardiovascular function, appetite, and psychosocial well-being in obese and DM patients (17). Concerning additional positive effects of RT, there remains, however, a lot of discussion. Progressive RT, for example, has been shown to lower blood LDL-C and non-HDL-C levels, as well as postprandial TG levels in some studies (88).

Gordon et al., (2016) (88) reported that maintaining muscle mass, along with increasing resting metabolic rate, better insulin balance, and improved fat metabolism are all likely underlying mechanisms for lowering blood lipid and lipoprotein levels and improving metabolic syndrome following RT. Increased skeletal muscle LPL activity and decreased hepatic TG lipase activity are two mechanisms by which physical exercise can yield beneficial changes in lipoprotein and lipid metabolism even in the absence of weight loss (89). Even with weight loss, these mechanisms affecting body structure may have played a role when BMI dramatically decreased while lean body mass significantly increased in obese individuals, leading to a noticeable change in the lipid profile (90). Furthermore, regular physical activity can have a positive impact on reverse cholesterol transport by lowering cholesterol ester transferase protein concentration or activity (91, 92) and increasing lecithin: acyl-cholesterol transferase activity

(93), which increases HDL2 levels preferentially.

When investigating the effects of RT on lipid and lipoprotein Jangio-Borazjani et al., (2021) (94) demonstrated that RT for 3 days/week consisting of a 10-min of warm-up, 45-min RT, conducted with 3 sets of 10 to 12 repetitions and loads ranging from 60% to 80% of 1-repetition maximum (1-RM) along with a 10-min cool-down, over a period of 8 weeks in 10 T2DM patients, resulted in a significant reduction in TG, TC, LDL-C, glucose, an increase in HDL-C levels, improved glycemic control, and muscle substrate metabolism. Particularly training programs shorter than 12 weeks, however, are unlikely to affect serum lipid and lipoprotein levels (59) as there may not be enough stimulus or enough time to trigger a change in blood lipids and lipoproteins (56). However, according to a meta-analysis by Kelley and Kelley (2009) (95), which included 29 studies representing 1329 men and women (676 exercise, 653 control), RT was related to a significant decrease in LDL-C. Another meta-analysis by Tambalis et al., (2009) (96) showed a significant reduction in LDL-C. Although each of these meta-analyses discovered lower LDL-C levels, neither research evaluated the association between lower LDL-C and improvements in body weight or body fat. Nonetheless, higher muscle mass and resulting changes in LPL function may have resulted in lower LDL-C levels (97). Hence, these findings contradict previous research suggesting that RT had no effect on LDL-C levels (98).

Low-density lipoprotein. In a systematic review and meta-analysis including 555 obese people, concurrent exercise with RT led to an increased body mass due to a higher lean mass, while fat mass and LDL-C decreased compared to AE alone (99). Moraleda et al., (2013) (100) further indicated that strength training in 24 obese individuals for 24 weeks was effective in improving the blood lipid profile by decreasing LDL-C, and TC values, while there was no change in HDL-C. According to a meta-analysis of 42 randomized controlled trials, structured exercise in three groups AE, RT, and a mixture of AE+RT was associated with improvement in blood pressure control, lower LDL-C, and higher HDL-C levels in DM individuals. The author stated that longer duration and higher volume were associated with significant improvements in SBP, HDL-C, and LDL-C in T2DM patients (69).

Baldi and Snowling (2003) (101), on the other hand, showed that 10 weeks of RT in T2DM patients with obesity, resulted in a significant reduction in fasting glucose and HbA1c levels while there was no substantial change in BMI, body fat, waist-hip ratio, and lipid profile TC, TG, LDL-C, and HDL-C. Accordingly, the study concluded that 12 weeks of RT did not induce adequate energy expenditure and lipolysis which is essential to decrease obesity and alter the lipid profile. In addition, some studies reported no alteration in LDL-C concentrations in men (102-104) and obese women (104) following RT.

Very low-density lipoprotein. Dâmaso et al., (2014) (105) conducted a study involving obese teenagers. A total of 139 participants were randomly divided into two groups: one undergoing AE alone (n=55) and the other undergoing a combination of AE and RT (n=61). The results indicated that the AE plus RT group showed superior outcomes in terms of body fat, body mass, and LDL-C ratio compared to the group undergoing AE alone. Additionally, both groups experienced a significant reduction in TG and VLDL.

Another study examined 30 obese women, aged 35 to 45 years, who were divided into three groups: control (n=10) AE (n=10), and RT (n=10) with the intervention lasting 6 weeks. AE consisted of 3 exercise sessions per week at 60-70% HR and RT was performed on alternating days for 6 weeks. AT resulted in a reduction in VLDL and a substantial rise in HDL-C, while both exercise groups displayed significant improvements in BMI and body fat ratio (106). In addition, a study by Gómez-Tomás et al., (2018) (107) showed a significant reduction in waist circumference, TC, and LDL-C following 1 year of elastic band RT. While Roh et al., (2020) (108) revealed that, twenty-six obese elderly women who underwent RT using elastic bands three days a week for 12 weeks, with each session consisting of 40 min of the main exercise, and a respective 10 min warm-up and cool-down, no substantial changes in serum lipid profiles following exercise.

High-density lipoprotein. For the control of obesity and T2DM, RT appears to be a crucial addition to diet and fitness regimens, since it increases muscle mass, functional capability, and improves lipoprotein profile and insulin resistance (4). A one-year resistance band training has been shown to dramatically enhance HDL-C

levels and decrease TC, LDL-C, and the TC/HDL-C ratio (90, 107). Hamasaki et al., (2015) (109) reported that low-intensity RT in 26 obese patients with T2DM subjected to 12-week exercise, displayed a significant increase in lower extremity muscle mass relative to body weight, while body fat mass and body fat percentage decreased. There was also a significant increase in HDL-C.

On the other hand, there were no significant changes in HDL-C levels in obese individuals engaging in strength training, endurance training, or a combination over a period of 24 weeks (100). Previous studies reported elevated (110), reduced (111), or stable HDL-C levels (112) with weight loss following exercise intervention. The disparity in findings appears to be due to the divergent consequences of weight loss. Some studies, however, have shown that lowering fat consumption in the diet lowers HDL-C levels, even when weight loss is achieved, in both short (113) and long-term (114) studies.

Total cholesterol. Metabolism is dysregulated in obese DM patients. From a medical perspective, elevated blood sugar levels and an abnormal lipid profile may damage different organs, including the heart, lungs, liver, and kidneys. It may also cause disorders like atherosclerosis, liver failure, and kidney disease (94). RT is considered a beneficial strategy to reduce these harmful disorders in obese DM patients. A total of 35 obese postmenopausal women with metabolic syndrome, were assigned to two groups: the control group (n=17) or the resistance band exercise training group (n=18) with resistance band exercise being conducted three days per week over a period of 12 weeks, consisting of 10 min warm-up, and 40 min of resistance band exercise. The intensity of the exercise was steadily increased from 40–50% 1-RM and a rating of perceived exertion (RPE) of 11–12 in weeks 1 to 4, to 50–60% 1RM and RPE 13–14 in weeks 5–8, and finally to 60–70% 1RM and RPE 15–16 in weeks 9–12. The results showed a significant decrease in blood glucose, insulin, TG, LDL-C, systolic BP, waist circumference, and body fat level, along with a significant increase in HDL-C with RT and the study concluded that RT can be an effective therapeutic intervention in obese adults with metabolic syndrome that, possibly decreases the risk for the development of CVD (92). Twenty-

one obese women who participated 3 days per week in whole-body RT (consisting of 10 exercises with 3 sets at 8-RM) for 12 weeks showed an improvement in TC and LDL-C levels (115). Another study also showed significant improvements in the levels of TC, TG, LDL-C, and HDL-C in overweight postmenopausal women following a conventional RT (116). A study with 30 teenagers between the ages of 15 and 19, who were randomly assigned either AT alone or AT+RT also showed slightly higher increases in body composition and lower, TC, waist circumference, glucose, and adiponectin in the combined training group than the AT group alone (117).

While others have reported no differences in TC concentrations after resistance training in obese women (104), Kokkinos and Hurley (1990) (118) suggested that this may be due to the differences in day-to-day lipid and lipoprotein patterns, nutritional structure, body composition, and plasma density. Further, limited sample sizes, no distinction between a single workout session and daily performed exercise, anabolic steroid use, the different types of RT, and the absence of or incorrectly chosen control groups may have contributed to inconsistent findings.

Triglycerides. In people with T2DM, accumulating experimental evidence suggests that serum TG is a leading indicator of atherosclerotic cardiovascular disease, comparable to LDL-C (36). Engagement in regular exercise may be beneficial to reduce TG in obese and DM patients. Gavin et al., (2010) (119) conducted a study in 64 adults with type 2 diabetes, who performed RT 3 times per week with a progressive increase in intensity and found a significant reduction in TG level following the exercise intervention.

A study of adults between 35 and 65 years of age participating in an RT program that lasted 60 to 75 min twice a week (150 minutes) for 12 weeks with a progressive increase in exercise intensity, on the other hand, showed no improvement in TC, HDL-C, LDL-C, and TG ratio following the exercise, and there was no change in TC ratio (81). In addition, cross-sectional trials evaluating regular involvement in RT training did not show any differences in plasma TG concentrations for males (120), and females (121) compared to the control group.

HIGH-INTENSITY EXERCISE

Obesity and type 2 diabetes, when combined with a lack of physical exercise might create serious health concerns in terms of death and morbidity (122). The intensity and duration of an exercise session are important considerations that must be considered when evaluating health benefits. Low-intensity training that is performed over prolonged periods uses fat as the main energy substrate, while high-intensity interval exercise uses glucose instead of fat as the dominant energy substrate. The differences in metabolism has lead to the conclusion that conventional low to moderate-intensity exercise, as compared to high-intensity exercise are beneficial for lipid parameter improvements (123, 124). Nevertheless, various exercise types can improve blood lipid profiles in people with metabolic dysregulation (125, 126). There are, however, mixed results, with some suggesting high-intensity exercise and some suggesting low-to moderate-intensity exercise is superior for improving lipid, lipoprotein, and blood pressure levels in obese diabetic patients.

In a recent study conducted by Abdelbasset et al., (2020) (80) 47 diabetic obese patients with nonalcoholic fatty liver disease, were randomly split into 16 participants performing high-intensity interval training (HIIT) exercise between 50% and 80% to 85% of HRmax, 3 times per week for 8 weeks and 15 patients performing moderate-intensity continuous exercise at 60% to 70% of HRmax 3 times per week showed a significant reduction in TC, TG, LDL-C and increased HDL-C levels in both HIIT and moderate-intensity exercise groups compared to inactive controls (80). In addition, a total of 51 obese adults with a sedentary lifestyle were studied for a total of 15 weeks. Participants completed moderate intensity (n=22) and HIIT exercise (n=29) on a bicycle ergometer for 40 min/day, five days/week, and 20 min/day, 3 days/week, respectively. Comparing LDL-C levels following moderate and HIIT exercise, the HIIT exercise group demonstrated a significant decrease. This may be due to the fact that with HIIT energy expenditure is higher during the recovery period (68). Excess post-exercise oxygen consumption may take several minutes after moderate-intensity exercise but several hours after HIT exercise, even though fat contains more Kcal of energy per gram than carbohydrate. According to the findings by, Kannan et al.,

(2014) (68) HIIT is comparatively healthy and can be progressively implemented in obese patients, while RT necessitates monitoring, instruction, and the need to increase repetitions. Nevertheless, significant changes in TC, LDL-C, HDL-C, TG, and VLDL were observed in the moderate-intensity group and changes in TC, HDL-C, LDL-C, and BMI were observed in the HIIT exercise group (127).

COMBINED AEROBIC AND RESISTANCE TRAINING

A population-based, epidemiological investigation has shown that individuals with T2DM have dyslipidemia (128). In another study, 24 T2DM patients were randomly placed in two groups: combined AE and RT (n=12) with 3 exercise sessions per week for 12 weeks or control (n=12). Results showed a substantial decline in TC, TG, LDL-c, and a rise in HDL-c level after the exercise intervention compared to the control group (129).

Exercise training has been shown in previous findings to enhance the lipid profile in T2DM; these improvements include a rise in HDL-C and a reduction in TC, TG, and LDL-C (130-136). In these studies, the study participants' average age was in the middle of fifty. Only two of these studies included older participants (>65 years old); however, the authors did not disclose the results from these elderly participants separately (131, 132). The improvement in the lipid profile was also found in the previous study following 12 weeks of combined aerobic and resistance training among overweight T2DM patients (137). Zarei et al., (2021) (138) reported that the combination of AE and RT performed three times a week for 12 weeks significantly reduced TC and TG in the exercise group compared to the control group. Magalhães et al., (2020) (139) conducted a randomized controlled trial among obese T2DM patients three times per week and found significant improvement in LDL-C and TC following the combination of AE and RT compared to the control group. Tan et al., (2012) (140) reported that three times a week of exercise, each session was divided into the following modules: a warm-up, 30 minutes of moderate AE, 10 minutes of RT with five leg exercises (two sets of 10–12 repetitions at 50–70% of 1RM for each activity), and a cool-down demonstrated a significant improvement in the lipid profile. In another study, twenty-four individuals diagnosed

with T2DM and obesity were assigned randomly to two groups: the first group, consisting of twelve participants, underwent a program involving AE and RT, while the second group, also comprising twelve participants, as the control group. After a 12-week intervention, the findings revealed a significant reduction in TC, TG, and LDL-c levels and an increase in HDL-c levels in the exercise group compared to the control group (129). Tokmakidis et al., (2004) (141) examined the effects of an AE and RT program on lipid profile in women with T2DM. The results showed that the exercise program improved the lipid profile, significantly reducing TC, LDL-C, and TG levels. Kwon et al., (2011) (142) compared the effects of AE and RT on lipid profile in women with T2DM. The findings demonstrated that aerobic exercise and resistance training reduced LDL-C and TG levels increased HDL-C levels, and improved the lipid profile, including reduced TC and TG levels. A study by Balducci et al., (2010) (143) demonstrated that a 12-week combined training programme significantly reduced TC and LDL-c in obese T2DM patients.

A study by Sigal et al., (2007) (144) found that a 22-week combined exercise program significantly increased HDL-C levels in obese T2DM patients. A study by Dunstan et al., (2002) (145) reported a reduction in TG levels following a 10-week combined training program in obese T2DM patients. A study by Tokmakidis et al., (2004) (141) showed that a 12-week combined AE and RT program significantly improved lipid profile ratios in T2DM patients. Azarbayjani et al., (2014) (146) examined the effect of combined AE and RT for 16 weeks among obese women. The results showed significant reductions in TC, LDL-C, and TG, while increasing HDL-C levels. Other studies have shown a significant reduction in TG (147-149), and TC (150-154).

Previous studies have shown a significant decrease in weight loss and a significant increase in HDL-C (148, 150). The literature describes an increase in HDL-C associated with decreased cardiovascular risk by a combination of aerobic and resistance training (155-157). The fact that combination training involved a larger muscle mass might be one of the reasons for these improvements. This could have caused a greater reduction in intramyocytic fat content and/or an increase in fatty acid oxidation capacity compared to the control groups. These beneficial effects may have caused an increase in the clearance of

lipids from circulation (76). Moreover, the training might explain these positive results, which have been demonstrated to increase fatty acid oxidation capacity and decrease intramyocytic lipid content (158). In the combination group, more muscles in the upper and lower limbs were worked out. Assuming that the combination program included higher muscle mass, it is likely that the total intramyocytic fat level decreases and the fatty acid oxidation ability enhances more.

A recent meta-analysis examined 12 weeks of AE and RT-affected lipid profiles in people with T2DM (159). The studies' findings were varied, with two claiming that exercise had no impact on LDL-C, HDL-C, TC, and TG (160, 161), and one study linked exercise to lower TC and alterations in HDL-C and LDL-C (162). Meanwhile, no difference in LDL-c and HDL-c was found following a combination of AE and RT (151, 152). The differences between these results may be explained by the participants' baseline values, which did not change significantly after exercise (139).

OTHER TYPES OF EXERCISE

Other types of exercise are Yoga and Pilates. Both are mind-body fitness modalities that induce favorable changes in lipid homeostasis in people with impaired lipid profiles (163-166). Yoga is an exercise that is a cost-effective and simple therapeutic modality for patients with obesity and T2DM (163). Yoga assists in the redistribution of body fat and the elimination of central obesity, all of which contribute to insulin resistance. Yoga has been associated with physical, neurological, psychological, and endocrinological improvements. Gordon et al., (2008) (167) reported that after a 6-month yoga intervention in 77 T2DM patients, there was a reduction in TC, TG, LDL-C, VLDL, and increased HDL-C level (167). Patients with CAD further showed substantial decreases in serum TC, TG, and LDL-C after one year of yoga therapy (168). Since yoga does not require excessive muscle movement or energy generation, its beneficial role in the treatment of hyperlipidemia and obesity cannot be due exclusively to increased energy expenditure (165).

Repeated stress is believed to induce a long-term increase in cortisol, which leads to central obesity and insulin resistance (169, 170). It enhances gluconeogenesis, thus lowering glucose

absorption in the peripheral tissues (171). Stress-induced growth hormone and endorphin release may indeed cause reduced glucose absorption and insulin secretion (172). Cortisol overproduction is associated with dyslipidemia, elevated blood pressure, and elevated levels of fatty molecules (lipids) in the blood, such as cholesterol and triglycerides both of which are associated with an increased risk of cardiovascular complications, for instance, heart failure (173). Yoga has been shown to lower sympathetic hormone and cortisol levels. Pranayama helps to relieve tension by lowering sympathetic tone and increasing parasympathetic activity (174). Meditation can also decrease stress-induced sympathetic overactivity and may induce a hypometabolic condition (175). A greater ability to deal with stress, resulting in lower cortisol levels, has been proposed as a potential explanation for improved lipid profiles in yoga practitioners. In the case of insulin resistance, dyslipidemia is typically associated with disturbances in lipolysis, TG metabolism, and free fatty acid turnover. Insulin resistance in DM is believed to induce reduced LPL and elevated HL function. Chronic susceptibility to high levels of free fatty acids has been linked to decreased insulin secretion (176). Increased HL and LPL, which would increase TG absorption by adipose tissue and influence lipoprotein metabolism, may explain the change in lipid profile with yoga practice.

On the other hand, some researchers have made adjustments for plasma volume change when measuring the results of a single session of physical exercise. In nine men, it was found that TG and TC concentrations were not altered directly after or in the days following a short-duration, low-intensity exercise session (177). TG concentrations have been reported to be reduced when the volume of work completed is high. Wallace et al., (1991) (178) reported that triglycerides were significantly decreased 24 hours post-exercise following a high-volume session in comparison to immediate post-exercise values. The findings of this research indicate that the volume of exercise has the greatest impact on acute changes in the lipoprotein profile. In general, triglyceride concentrations will typically be lower directly following and/or during the days after a single workout session if the exercise session is lengthy and requires a lot of energy. There was no alteration in HDL-C after 1.5 hours of exercise requiring about 77% of maximal heart

rate (179). On the other hand, HDL-C concentrations are normally elevated shortly following extended activity (more than 1.5 hours) (180). Thus, future research should focus on the dose-response effects of various of real-world exercise programs as previously reported (181). Clinicians and practitioners should be aware of this valuable information in order to promote exercise prescription to individuals with impaired metabolic health (182).

CONCLUSION

Physical activity has a positive effect on the lipid and lipoprotein profile in patients with obesity and T2DM. Regular exercise increases HDL-C and lowers plasma triglyceride concentrations, total cholesterol, VLDL, and LDL-C. Engaging in adequate and regular exercise also decreases the risk of CVD, high BMI and improves insulin resistance. However, these benefits are dependent on the types of exercise, volume, intensity, frequency, and duration of each session.

APPLICABLE REMARKS

- Physical activity induces beneficial changes in blood lipid levels among patients with obesity and T2DM.
- Exercise increases HDL-C and lowers plasma TG concentrations, TC, VLDL, and LDL-C.
- Engaging in adequate and regular physical exercise also reduces the risk of insulin resistance and CVD.
- Exercise-induced positive alterations are dependent on the exercise training parameters.

ACKNOWLEDGMENTS

School of Medical Sciences of the Universiti Sains Malaysia, grant number 304.PPSP.6315639.

AUTHORS' CONTRIBUTIONS

Study concept and design: Sameer Badri Al-mhanna, Wan Syaheedah Wan Ghazali, Alexios Batrakoulis, Nouf H. Alkhamees. Acquisition of data: Clemens Drenowatz, Mahaneem Mohamed. Analysis and interpretation of data: Mehmet Güllü, Hafeez Abiola Afolabi, Georgian Badicu. Drafting the manuscript: Alexios Batrakoulis, Nouf H. Alkhamees, Clemens Drenowatz. Critical revision of the manuscript for important intellectual content: Nouf H. Alkhamees, Clemens Drenowatz, Mehmet Güllü, Hafeez Abiola Afolabi, Georgian Badicu. Statistical analysis: Sameer Badri Al-

mhanna, Wan Syaheedah Wan Ghazali , Alexios Batrakoulis, Nouf H. Alkhamees. Administrative, technical, and material support: Clemens Drenowatz, Mahaneem Mohamed, Mehmet Güli, Hafeez Abiola Afolabi, Georgian Badicu. Study supervision: Sameer Badri Al-mhanna, Wan

Syaheedah Wan Ghazali, Nouf H. Alkhamees, Mahaneem Mohamed.

CONFLICT OF INTEREST

No potential conflict of interest was reported by the authors.

REFERENCES

1. Kimura T, Itoh T, Fusazaki T, Matsui H, Sugawara S, Ogino Y, Endo H, Kobayashi K, Nakamura M. Low-Density Lipoprotein-Cholesterol/High-Density Lipoprotein-Cholesterol Ratio Predicts Lipid-Rich Coronary Plaque in Patients With Coronary Artery Disease—Integrated-Backscatter Intravascular Ultrasound Study. *Circulation Journal*. 2010;74(7):1392-8. [doi:10.1253/circj.CJ-09-0849] [PMid:20484825]
2. Saeidi A, Hackney AC, Tayebi SM, Ahmadian M, Zouhal H. Diabetes, insulin resistance, fetuin-B and exercise training. *Annals of applied sport science*. 2019;7(2):1-2. [doi:10.29252/aassjournal.7.2.1] [PMid:31321377]
3. Sun H, Saeedi P, Karuranga S, Pinkepank M, Ogurtsova K, Duncan BB, et al. IDF Diabetes Atlas: Global, regional and country-level diabetes prevalence estimates for 2021 and projections for 2045. *Diabetes research and clinical practice*. 2022;183:109119. [doi:10.1016/j.diabres.2021.109119] [PMid:34879977]
4. Badri Al-mhanna S, Leão C, Wan Ghazali WS, Mohamed M, Batrakoulis A, Abiola Afolabi H, et al. Impact of Exercise on High-Density Lipoprotein Cholesterol in Adults with Overweight and Obesity: A Narrative Review. *Annals of Applied Sport Science*.0-0.
5. Haffner SM, Mykkänen L, Festa A, Burke JP, Stern MP. Insulin-resistant prediabetic subjects have more atherogenic risk factors than insulin-sensitive prediabetic subjects: implications for preventing coronary heart disease during the prediabetic state. *Circulation*. 2000 Mar 7;101(9):975-80. [doi:10.1161/01.CIR.101.9.975] [PMid:10704163]
6. Haffner SM, Stern MP, Hazuda HP, Mitchell BD, Patterson JK. Cardiovascular risk factors in confirmed prediabetic individuals: does the clock for coronary heart disease start ticking before the onset of clinical diabetes?. *Jama*. 1990 Jun 6;263(21):2893-8. [doi:10.1001/jama.263.21.2893] [PMid:2338751]
7. Garvey WT, Kwon S, Zheng D, Shaughnessy S, Wallace P, Hutto A, Pugh K, Jenkins AJ, Klein RL, Liao Y. Effects of insulin resistance and type 2 diabetes on lipoprotein subclass particle size and concentration determined by nuclear magnetic resonance. *Diabetes*. 2003 Feb 1;52(2):453-62. [doi:10.2337/diabetes.52.2.453] [PMid:12540621]
8. Afolabi HA, Zakaria Z, Salleh SM, Ch'ng ES, Nafi SNM, Aziz AABA, et al. Obesity: A Prerequisite for Major Chronic Illnesses. *Obesity-Recent Insights and Therapeutic Options: IntechOpen*; 2023.
9. Afolabi HA, Salleh SM, Zakaria Z, Ch'ng ES, Mohd Nafi SN, Abdul Aziz AAB, et al. A GNAS Gene Mutation's Independent Expression in the Growth of Colorectal Cancer: A Systematic Review and Meta-Analysis. *Cancers*. 2022;14(22):5480. [doi:10.3390/cancers14225480] [PMid:36428574]
10. Grundy SM, Cleeman JI, Daniels SR, Donato KA, Eckel RH, Franklin BA, Gordon DJ, Krauss RM, Savage PJ, Smith Jr SC, Spertus JA. Diagnosis and management of the metabolic syndrome: an American Heart Association/National Heart, Lung, and Blood Institute scientific statement. *Circulation*. 2005 Oct 25;112(17):2735-52. [doi:10.1161/CIRCULATIONAHA.105.169404] [PMid:16157765]
11. Afolabi HA, bin Zakariya Z, Shokri ABA, Hasim MNBM, Vinayak R, Afolabi-Owolabi OT, et al. The relationship between obesity and other medical comorbidities. *Obesity Medicine*. 2020;17:100164. [doi:10.1016/j.obmed.2019.100164]
12. Paley CA, Johnson MI. Abdominal obesity and metabolic syndrome: exercise as medicine?. *BMC Sports Science, Medicine and Rehabilitation*. 2018 Dec;10(1):1-8. [doi:10.1186/s13102-018-0097-1] [PMid:29755739]
13. Bharath LP, Choi WW, Cho JM, Skobodzinski AA, Wong A, Sweeney TE, Park SY. Combined resistance and aerobic exercise training reduces insulin resistance and central adiposity in adolescent girls who are obese: randomized clinical trial. *European journal of applied physiology*. 2018 Aug;118:1653-60. [doi:10.1007/s00421-018-3898-8] [PMid:29846794]

14. Pekas EJ, Shin J, Son WM, Headid III RJ, Park SY. Habitual combined exercise protects against age-associated decline in vascular function and lipid profiles in elderly postmenopausal women. *International journal of environmental research and public health*. 2020 Jun;17(11):3893. [doi:10.3390/ijerph17113893] [PMid:32486335]
15. Shaw IN, Shaw BS, Krasilshchikov OL. Comparison of aerobic and combined aerobic and resistance training on low-density lipoprotein cholesterol concentrations in men: cardiovascular topic. *Cardiovascular journal of Africa*. 2009 Sep 1;20(5):290-5.
16. Kercher VM, Kercher K, Levy P, Bennion T, Alexander C, Amaral PC, et al. 2023 Fitness Trends from Around the Globe. *ACSMs Health Fit J*. 2023;27(1):19-30. [doi:10.1249/FIT.0000000000000836]
17. Al-Mhanna SB, Rocha-Rodriguesc S, Mohamed M, Batrakoulis A, Aldhahi MI, Afolabi HA, et al. Effects of combined aerobic exercise and diet on cardiometabolic health in patients with obesity and type 2 diabetes: a systematic review and meta-analysis. *BMC Sports Science, Medicine and Rehabilitation*. 2023;15(1):165. [doi:10.1186/s13102-023-00766-5] [PMid:38049873]
18. Yapici H, Gülü M, Yagin FH, Ugurlu D, Comertpay E, Eroglu O, et al. The effect of 8-weeks of combined resistance training and chocolate milk consumption on maximal strength, muscle thickness, peak power and lean mass, untrained, university-aged males. *Frontiers in physiology*. 2023;14:1148494. [doi:10.3389/fphys.2023.1148494] [PMid:37007992]
19. Gondim OS, Camargo VT, Gutierrez FA, Martins PF, Passos ME, Momesso CM, Santos VC, Gorjão R, Pithon-Curi TC, Cury-Boaventura MF. Benefits of regular exercise on inflammatory and cardiovascular risk markers in normal weight, overweight and obese adults. *PloS one*. 2015 Oct 16;10(10):e0140596. [doi:10.1371/journal.pone.0140596] [PMid:26474157]
20. Wahid A, Manek N, Nichols M, Kelly P, Foster C, Webster P, Kaur A, Friedemann Smith C, Wilkins E, Rayner M, Roberts N. Quantifying the association between physical activity and cardiovascular disease and diabetes: a systematic review and meta-analysis. *Journal of the American Heart Association*. 2016 Sep 14;5(9):e002495. [doi:10.1161/JAHA.115.002495] [PMid:27628572]
21. Marandi SM, Abadi NGB, Esfarjani F, Mojtahedi H, Ghasemi G. Effects of intensity of aerobics on body composition and blood lipid profile in obese/overweight females. *International journal of preventive medicine*. 2013;4(Suppl 1):S118.
22. Janssen I, Fortier A, Hudson R, Ross R. Effects of an energy-restrictive diet with or without exercise on abdominal fat, intermuscular fat, and metabolic risk factors in obese women. *Diabetes care*. 2002;25(3):431-8. [doi:10.2337/diacare.25.3.431] [PMid:11874926]
23. Halverstadt A, Phares DA, Wilund KR, Goldberg AP, Hagberg JM. Endurance exercise training raises high-density lipoprotein cholesterol and lowers small low-density lipoprotein and very low-density lipoprotein independent of body fat phenotypes in older men and women. *Metabolism*. 2007;56(4):444-50. [doi:10.1016/j.metabol.2006.10.019] [PMid:17378998]
24. Visich P, Goss F, Warty V, Denys B, Robertson R, Metz K, Gordon P. Effect of caloric expenditure on acute changes in high density lipoprotein-cholesterol and related subfractions: 1061. *Medicine & Science in Sports & Exercise*. 1992 May 1;24(5):S177. [doi:10.1249/00005768-199205001-01062]
25. Larrydurstine J, Haskell WL. Effects of exercise training on plasma lipids and lipoproteins. *Exercise and sport sciences reviews*. 1994 Jan 1;22(1):477-522. [doi:10.1249/00003677-199401000-00017]
26. Farber H. Acute metabolic effects of an endurance triathlon. *Ann. Sports Med.*. 1987;3:131-8.
27. Lamon-Fava S, McNamara JR, Farber HW, Hill NS, Schaefer EJ. Acute changes in lipid, lipoprotein, apolipoprotein, and low-density lipoprotein particle size after an endurance triathlon. *Metabolism*. 1989 Sep 1;38(9):921-5. [doi:10.1016/0026-0495(89)90243-6] [PMid:2505019]
28. Lee R, Nieman D, Raval R, Blankenship J, Lee J. The effects of acute moderate exercise on serum lipids and lipoproteins in mildly obese women. *International journal of sports medicine*. 1991 Dec;12(06):537-42. [doi:10.1055/s-2007-1024730] [PMid:1797694]
29. Fiuza-Luces C, Santos-Lozano A, Joyner M, Carrera-Bastos P, Picazo O, Zugaza JL, et al. Exercise benefits in cardiovascular disease: beyond attenuation of traditional risk factors. *Nature Reviews Cardiology*. 2018;15(12):731-43. [doi:10.1038/s41569-018-0065-1] [PMid:30115967]
30. Al-Mhanna SB, Ghazali WSW, Mohamed M, Rabaan AA, Santali EY, Alestad JH, et al. Effectiveness of physical activity on immunity markers and quality of life in cancer patient: a systematic review. *Peer J*. 2022;10:e13664. [doi:10.7717/peerj.13664] [PMid:35935260]

31. Hansen D, Dendale P, Jonkers R, Beelen M, Manders R, Corluy L, et al. Continuous low-to moderate-intensity exercise training is as effective as moderate-to high-intensity exercise training at lowering blood HbA 1c in obese type 2 diabetes patients. *Diabetologia*. 2009;52:1789-97. [doi:10.1007/s00125-009-1354-3] [PMid:19370339]
32. Talanian JL, Galloway SD, Heigenhauser GJ, Bonen A, Spriet LL. Two weeks of high-intensity aerobic interval training increases the capacity for fat oxidation during exercise in women. *Journal of applied physiology*. 2007 Apr 1. [doi:10.1152/jappphysiol.01098.2006] [PMid:17170203]
33. Racil G, Ben Ounis O, Hammouda O, Kallel A, Zouhal H, Chamari K, et al. Effects of high vs. moderate exercise intensity during interval training on lipids and adiponectin levels in obese young females. *European journal of applied physiology*. 2013;113:2531-40. [doi:10.1007/s00421-013-2689-5] [PMid:23824463]
34. Hayashi T, Kawashima S, Itoh H, Yamada N, Sone H, Watanabe H, Hattori Y, Ohru T, Yokote K, Nomura H, Umegaki H. Low HDL cholesterol is associated with the risk of stroke in elderly diabetic individuals: changes in the risk for atherosclerotic diseases at various ages. *Diabetes care*. 2009 Jul 1;32(7):1221-3. [doi:10.2337/dc08-1677] [PMid:19509009]
35. Xiao C, Dash S, Morgantini C, Hegele RA, Lewis GF. Pharmacological targeting of the atherogenic dyslipidemia complex: the next frontier in CVD prevention beyond lowering LDL cholesterol. *Diabetes*. 2016 Jul 1;65(7):1767-78. [doi:10.2337/db16-0046] [PMid:27329952]
36. Hirano T. Pathophysiology of diabetic dyslipidemia. *Journal of atherosclerosis and thrombosis*. 2018 Sep 1;25(9):771-82. [doi:10.5551/jat.RV17023] [PMid:29998913]
37. Saleheen D, Scott R, Javad S, Zhao W, Rodrigues A, Picataggi A, Lukmanova D, Mucksavage ML, Luben R, Billheimer J, Kastelein JJ. Association of HDL cholesterol efflux capacity with incident coronary heart disease events: a prospective case-control study. *The lancet Diabetes & endocrinology*. 2015 Jul 1;3(7):507-13. [doi:10.1016/S2213-8587(15)00126-6] [PMid:26025389]
38. Khera AV, Cuchel M, de la Llera-Moya M, Rodrigues A, Burke MF, Jafri K, French BC, Phillips JA, Mucksavage ML, Wilensky RL, Mohler ER. Cholesterol efflux capacity, high-density lipoprotein function, and atherosclerosis. *New England Journal of Medicine*. 2011 Jan 13;364(2):127-35. [doi:10.1056/NEJMoa1001689] [PMid:21226578]
39. Barter P, McPherson YR, Song K, Kesäniemi YA, Mahley R, Waeber G, et al. Serum insulin and inflammatory markers in overweight individuals with and without dyslipidemia. 2007;92(6):2041-5. [doi:10.1210/jc.2006-2219] [PMid:17389699]
40. *care ADAJD*. Diabetes mellitus and exercise. 2002;25(suppl 1):s64-s. [doi:10.2337/diacare.25.2007.S64]
41. Al-Mhanna SB, Mohamed M, Mohd Noor N, Aldhahi MI, Afolabi HA, Mutalub YB, et al., editors. Effects of Circuit Training on Patients with Knee Osteoarthritis: A Systematic Review and Meta-Analysis. *Healthcare*; 2022: MDPI. [doi:10.3390/healthcare10102041] [PMid:36292488]
42. Rigla M, Sánchez-Quesada JL, Ordóñez-Llanos J, Prat T, Caixàs A, Jorba O, et al. Effect of physical exercise on lipoprotein (a) and low-density lipoprotein modifications in type 1 and type 2 diabetic patients. 2000;49(5):640-7. [doi:10.1016/S0026-0495(00)80041-4] [PMid:10831176]
43. Wang Y, Xu D. Effects of aerobic exercise on lipids and lipoproteins. *Lipids in health and disease*. 2017;16(1):1-8. [doi:10.1186/s12944-015-0172-5] [PMid:30611256]
44. Azizi R, Mohammadi Domieh A. Effects of aerobic exercise on lipids profile and insulin resistance in patients with type 2 diabetes. *Journal of Physical Activity and Hormones*. 2019 Sep 1;3(3):23-34.
45. Wallberg-Henriksson H, Gunnarsson R, Rössner S, Wahren J. Long-term physical training in female type 1 (insulin-dependent) diabetic patients: absence of significant effect on glycaemic control and lipoprotein levels. *Diabetologia*. 1986 Jan;29:53-7. [doi:10.1007/BF02427281] [PMid:3956895]
46. Williams PT, Krauss RM, Wood PD, Lindgren FT, Giotas C, Vranizan KM. Lipoprotein subfractions of runners and sedentary men. *Metabolism*. 1986 Jan 1;35(1):45-52. [doi:10.1016/0026-0495(86)90094-6] [PMid:3941608]
47. Wood PD, Haskell W, Klein H, Lewis S, Stern MP, Farquhar JW. The distribution of plasma lipoproteins in middle-aged male runners. *Metabolism*. 1976 Nov 1;25(11):1249-57. [doi:10.1016/S0026-0495(76)80008-X] [PMid:185487]

48. Williams PT, Krauss RM, Vranizan KM, Albers JJ, Wood PD. Effects of weight-loss by exercise and by diet on apolipoproteins AI and A-II and the particle-size distribution of high-density lipoproteins in men. *Metabolism*. 1992 Apr 1;41(4):441-9. [doi:10.1016/0026-0495(92)90082-L] [PMid:1556953]
49. Aggarwala J, Sharma S, Saroochi JA, Sarkar A. Effects of aerobic exercise on blood glucose levels and lipid profile in Diabetes Mellitus type 2 subjects. *Al Ameen J Med Sci*. 2016 Jan 1;9(1):65-9.
50. Mooradian AD. Dyslipidemia in type 2 diabetes mellitus. *Nature Reviews Endocrinology*. 2009 Mar;5(3):150-9. [doi:10.1038/ncpendmet1066] [PMid:19229235]
51. Solano MP, Goldberg RB. Management of dyslipidemia in diabetes. *Cardiology in review*. 2006 May 1;14(3):125-35. [doi:10.1097/01.crd.0000188034.76283.5e] [PMid:16628021]
52. Hamilton SJ, Watts GF. Endothelial dysfunction in diabetes: pathogenesis, significance, and treatment. *The review of diabetic studies: RDS*. 2013;10(2-3):133. [doi:10.1900/RDS.2013.10.133] [PMid:24380089]
53. Lampman RM, Schteingart DE. Effects of exercise training on glucose control, lipid metabolism, and insulin sensitivity in hypertriglyceridemia and non-insulin dependent diabetes mellitus. *Medicine and science in sports and exercise*. 1991 Jun 1;23(6):703-12. [doi:10.1249/00005768-199106000-00009]
54. Turcotte LP, Fisher JS. Skeletal muscle insulin resistance: roles of fatty acid metabolism and exercise. *Physical therapy*. 2008 Nov 1;88(11):1279-96. [doi:10.2522/ptj.20080018] [PMid:18801860]
55. Alam S, Stolinski M, Pentecost C, Boroujerdi MA, Jones RH, Sonksen PH, Umpleby AM. The effect of a six-month exercise program on very low-density lipoprotein apolipoprotein B secretion in type 2 diabetes. *The Journal of Clinical Endocrinology & Metabolism*. 2004 Feb 1;89(2):688-94. [doi:10.1210/jc.2003-031036] [PMid:14764782]
56. Durstine JL, Grandjean PW, Cox CA, Thompson PD. Lipids, lipoproteins, and exercise. *J Cardiopulm Rehabil*. 2002;22:385-98. [doi:10.1097/00008483-200211000-00002] [PMid:12464825]
57. Kelley GA, Kelley KS, Vu Tran Z. Aerobic exercise, lipids and lipoproteins in overweight and obese adults: a meta-analysis of randomized controlled trials. *International journal of obesity*. 2005 Aug;29(8):881-93. [doi:10.1038/sj.ijo.0802959] [PMid:15824746]
58. Keller C, Treviño RP. Effects of two frequencies of walking on cardiovascular risk factor reduction in Mexican American women. *Research in nursing & health*. 2001 Oct;24(5):390-401. [doi:10.1002/nur.1039] [PMid:11746068]
59. Goldberg AC, Hopkins PN, Toth PP, Ballantyne CM, Rader DJ, Robinson JG, Daniels SR, Gidding SS, De Ferranti SD, Ito MK, McGowan MP. Familial hypercholesterolemia: screening, diagnosis and management of pediatric and adult patients: clinical guidance from the National Lipid Association Expert Panel on Familial Hypercholesterolemia. *Journal of clinical lipidology*. 2011 May 1;5(3):133-40. [doi:10.1016/j.jacl.2011.03.001] [PMid:21600517]
60. Thent ZC, Das S, Henry LJ. Role of exercise in the management of diabetes mellitus: the global scenario. *PloS one*. 2013 Nov 13;8(11):e80436. [doi:10.1371/journal.pone.0080436] [PMid:24236181]
61. Shakil-ur-Rehman S, Karimi H, Gillani SA. Effects of supervised structured aerobic exercise training program on high and low density lipoprotein in patients with type II diabetes mellitus. *Pakistan Journal of Medical Sciences*. 2017 Jan;33(1):96. [doi:10.12669/pjms.331.11758]
62. Nassef Y, Lee KJ, Nfor ON, Tantoh DM, Chou MC, Liaw YP. The impact of aerobic exercise and badminton on HDL cholesterol levels in adult Taiwanese. *Nutrients*. 2019 Feb 28;11(3):515. [doi:10.3390/nu11030515] [PMid:30823398]
63. Boyer M, Mitchell PL, Poirier P, Alméras N, Tremblay A, Bergeron J, Després JP, Arsenault BJ. Impact of a one-year lifestyle modification program on cholesterol efflux capacities in men with abdominal obesity and dyslipidemia. *American Journal of Physiology-Endocrinology and Metabolism*. 2018 Oct 1;315(4):E460-8. [doi:10.1152/ajpendo.00127.2018] [PMid:29870675]
64. Furuyama F, Koba S, Yokota Y, Tsunoda F, Shoji M, Kobayashi Y. Effects of cardiac rehabilitation on high-density lipoprotein-mediated cholesterol efflux capacity and paraoxonase-1 activity in patients with acute coronary syndrome. *Journal of atherosclerosis and thrombosis*. 2018 Feb 1;25(2):153-69. [doi:10.5551/jat.41095] [PMid:28855433]
65. Leon AS, Conrad J, Hunninghake DB, Serfass R. Effects of a vigorous walking program on body composition, and carbohydrate and lipid metabolism of obese young men. *The American Journal of Clinical Nutrition*. 1979 Sep 1;32(9):1776-87. [doi:10.1093/ajcn/32.9.1776] [PMid:474467]

66. Lewis S, Haskell WL, Wood PD, Manoogian N, Bailey JE, Pereira MB. Effects of physical activity on weight reduction in obese middle-aged women. *The American Journal of Clinical Nutrition*. 1976 Feb 1;29(2):151-6. [doi:10.1093/ajcn/29.2.151] [PMid:1251807]
67. Rashid S, Genest J. Effect of obesity on high-density lipoprotein metabolism. *Obesity*. 2007 Dec;15(12):2875-88. [doi:10.1038/oby.2007.342] [PMid:18198293]
68. Kannan U, Vasudevan K, Balasubramaniam K, Yerrabelli D, Shanmugavel K, John NA. Effect of exercise intensity on lipid profile in sedentary obese adults. *Journal of clinical and diagnostic research: JCDR*. 2014 Jul;8(7):BC08. [doi:10.7860/JCDR/2014/8519.4611] [PMid:25177556]
69. Hayashino Y, Jackson JL, Fukumori N, Nakamura F, Fukuhara S. Effects of supervised exercise on lipid profiles and blood pressure control in people with type 2 diabetes mellitus: a meta-analysis of randomized controlled trials. *Diabetes research and clinical practice*. 2012 Dec 1;98(3):349-60. [doi:10.1016/j.diabres.2012.10.004] [PMid:23116535]
70. Cai M, Zou Z. Effect of aerobic exercise on blood lipid and glucose in obese or overweight adults: A meta-analysis of randomised controlled trials. *Obesity research & clinical practice*. 2016;10(5):589-602. [doi:10.1016/j.orep.2015.10.010] [PMid:26777793]
71. Kantor MA, Cullinane EM, Sady SP, Herbert PN, Thompson PD. Exercise acutely increases high density lipoprotein-cholesterol and lipoprotein lipase activity in trained and untrained men. *Metabolism*. 1987;36(2):188-92. [doi:10.1016/0026-0495(87)90016-3] [PMid:3807790]
72. Kelley GA, Kelley KS. Effects of aerobic exercise on lipids and lipoproteins in adults with type 2 diabetes: a meta-analysis of randomized-controlled trials. *Public health*. 2007 Sep 1;121(9):643-55. [doi:10.1016/j.puhe.2007.02.014] [PMid:17544042]
73. Thompson PD, Cullinane EM, Sady SP, Flynn MM, Bernier DN, Kantor MA, et al. Modest changes in high-density lipoprotein concentration and metabolism with prolonged exercise training. *Circulation*. 1988;78(1):25-34. [doi:10.1161/01.CIR.78.1.25] [PMid:3383408]
74. Wilund KR, Feeney LA, Tomayko EJ, Weiss EP, Hagberg JM. Effects of endurance exercise training on markers of cholesterol absorption and synthesis. *Physiological research*. 2009;58(4):545-52. [doi:10.33549/physiolres.931515] [PMid:18656998]
75. Tarnopolsky MA, Rennie CD, Robertshaw HA, Fedak-Tarnopolsky SN, Devries MC, Hamadeh MJ. Influence of endurance exercise training and sex on intramyocellular lipid and mitochondrial ultrastructure, substrate use, and mitochondrial enzyme activity. *American journal of physiology Regulatory, integrative and comparative physiology*. 2007;292(3):R1271-8. [doi:10.1152/ajpregu.00472.2006] [PMid:17095651]
76. Wang Y, Xu D. Effects of aerobic exercise on lipids and lipoproteins. *Lipids in health and disease*. 2017;16(1):132. [doi:10.1186/s12944-017-0515-5] [PMid:28679436]
77. Grandjean PW, Crouse SF, Rohack JJ. Influence of cholesterol status on blood lipid and lipoprotein enzyme responses to aerobic exercise. *Journal of applied physiology (Bethesda, Md : 1985)*. 2000;89(2):472-80. [doi:10.1152/jappl.2000.89.2.472] [PMid:10926628]
78. Laaksonen DE, Atalay MU, Niskanen LK, Mustonen JU, Lakka TA, Uusitupa MI. Aerobic exercise and the lipid profile in type 1 diabetic men: a randomized controlled trial. *Medicine & Science in Sports & Exercise*. 2000 Sep 1;32(9):1541-8. [doi:10.1097/00005768-200009000-00003] [PMid:10994902]
79. Wallberg-Henriksson H, Gunnarsson R, Henriksson J, DeFronzo R, Felig P, Östman J, Wahren J. Increased peripheral insulin sensitivity and muscle mitochondrial enzymes but unchanged blood glucose control in type I diabetics after physical training. *Diabetes*. 1982 Dec 1;31(12):1044-50. [doi:10.2337/diacare.31.12.1044] [PMid:6757018]
80. Abdelbasset WK, Tantawy SA, Kamel DM, Alqahtani BA, Elnegamy TE, Soliman GS, Ibrahim AA. Effects of high-intensity interval and moderate-intensity continuous aerobic exercise on diabetic obese patients with nonalcoholic fatty liver disease: a comparative randomized controlled trial. *Medicine*. 2020 Mar;99(10). [doi:10.1097/MD.00000000000019471] [PMid:32150108]
81. Ranasinghe C, Devage S, Constantine GR, Katulanda P, Hills AP, King NA. Glycemic and cardiometabolic effects of exercise in South Asian Sri Lankans with type 2 diabetes mellitus: A randomized controlled trial Sri Lanka diabetes aerobic and resistance training study (SL-DARTS). *Diabetes & Metabolic Syndrome: Clinical Research & Reviews*. 2021 Jan 1;15(1):77-85. [doi:10.1016/j.dsx.2020.12.011] [PMid:33310265]

82. Akcakoyun F. Changes in serum lipid profile following moderate exercise. *Afr J Pharm Pharmacol*. 2010 Nov 1;4(11):829-33.
83. Tayebi SM, Golmohammadi M, Eslami R, Shakiba N, Costa PB. The Effects of Eight Weeks of Circuit Resistance Training on Serum METRNL Levels and Insulin Resistance in Individuals with Type 2 Diabetes. *Journal of Diabetes & Metabolic Disorders*. 2023:1-8. [doi:10.1007/s40200-023-01225-1] [PMid:37975113]
84. Tan J, Krasilshchikov O, Kuan G, Hashim HA, Aldhahi MI, Al-Mhanna SB, et al., editors. The Effects of Combining Aerobic and Heavy Resistance Training on Body Composition, Muscle Hypertrophy, and Exercise Satisfaction in Physically Active Adults. *Healthcare*; 2023: MDPI. [doi:10.3390/healthcare11172443] [PMid:37685476]
85. Haskell WL, Lee IM, Pate RR, Powell KE, Blair SN, Franklin BA, Macera CA, Heath GW, Thompson PD, Bauman A. Physical activity and public health: updated recommendation for adults from the American College of Sports Medicine and the American Heart Association. *Circulation*. 2007;116(9):1081. [doi:10.1161/CIRCULATIONAHA.107.185649] [PMid:17671237]
86. Lacroix A, Hortobagyi T, Beurskens R, Granacher U. Effects of supervised vs. unsupervised training programs on balance and muscle strength in older adults: a systematic review and meta-analysis. *Sports medicine*. 2017 Nov;47:2341-61. [doi:10.1007/s40279-017-0747-6] [PMid:28573401]
87. Pollock ML, Franklin BA, Balady GJ, Chaitman BL, Fleg JL, Fletcher B, Limacher M, Piña IL, Stein RA, Williams M, Bazzarre T. Resistance exercise in individuals with and without cardiovascular disease: benefits, rationale, safety, and prescription an advisory from the committee on exercise, rehabilitation, and prevention, council on clinical cardiology, American Heart Association. *Circulation*. 2000 Feb 22;101(7):828-33. [doi:10.1161/01.CIR.101.7.828] [PMid:10683360]
88. Gordon B, Chen S, Durstine JL. The effects of exercise training on the traditional lipid profile and beyond. *Current sports medicine reports*. 2014 Jul 1;13(4):253-9. [doi:10.1249/JSR.0000000000000073] [PMid:25014391]
89. Thompson PD, Yurgalevitch SM, Flynn MM, Zmuda JM, Spannaus-Martin D, Saritelli A, Bausserman L, Herbert PN. Effect of prolonged exercise training without weight loss on high-density lipoprotein metabolism in overweight men. *Metabolism*. 1997 Feb 1;46(2):217-23. [doi:10.1016/S0026-0495(97)90305-X] [PMid:9030832]
90. Bembien DA, Bembien MG. Effects of resistance exercise and body mass index on lipoprotein-lipid patterns of postmenopausal women. *The Journal of Strength & Conditioning Research*. 2000 Feb 1;14(1):80-5. [doi:10.1519/00124278-200002000-00014]
91. Seip RL, Angelopoulos TJ, Semenkovich CF. Exercise induces human lipoprotein lipase gene expression in skeletal muscle but not adipose tissue. *American Journal of Physiology-Endocrinology and Metabolism*. 1995 Feb 1;268(2):E229-36. [doi:10.1152/ajpendo.1995.268.2.E229] [PMid:7864098]
92. Son WM, Park JJ. Resistance band exercise training prevents the progression of metabolic syndrome in obese postmenopausal women. *Journal of sports science & medicine*. 2021 Jun;20(2):291. [doi:10.52082/jssm.2021.291] [PMid:34211322]
93. Marniemi J, Dahlström S, Kvist M, Seppänen A, Hietanen E. Dependence of serum lipid and lecithin: cholesterol acyltransferase levels on physical training in young men. *European Journal of Applied Physiology and Occupational Physiology*. 1982 Jun;49(1):25-35. [doi:10.1007/BF00428960] [PMid:7201926]
94. Jangjo-Borazjani S, Dastgheib M, Kiyamarsi E, Jamshidi R, Rahmati-Ahmadabad S, Helalizadeh M, Irajji R, Cornish SM, Mohammadi-Darestani S, Khojasteh Z, Azarbayjani MA. Effects of resistance training and nigella sativa on type 2 diabetes: implications for metabolic markers, low-grade inflammation and liver enzyme production. *Archives of Physiology and Biochemistry*. 2023 Jul 4;129(4):913-21. [doi:10.1080/13813455.2021.1886117] [PMid:33612031]
95. Kelley GA, Kelley KS. Impact of progressive resistance training on lipids and lipoproteins in adults: a meta-analysis of randomized controlled trials. *Preventive medicine*. 2009 Jan 1;48(1):9-19. [doi:10.1016/j.ypmed.2008.10.010] [PMid:19013187]
96. Tambalis K, Panagiotakos DB, Kavouras SA, Sidossis LS. Responses of blood lipids to aerobic, resistance, and combined aerobic with resistance exercise training: a systematic review of current evidence. *Angiology*. 2009 Nov;60(5):614-32. [doi:10.1177/0003319708324927] [PMid:18974201]

97. Brinkley TE, Halverstadt A, Phares DA, Ferrell RE, Prigeon RL, Hagberg JM, Goldberg AP. Hepatic lipase gene-514C> T variant is associated with exercise training-induced changes in VLDL and HDL by lipoprotein lipase. *Journal of Applied Physiology*. 2011 Dec;111(6):1871-6. [doi:10.1152/jappphysiol.00567.2011] [PMid:21960661]
98. Williams MA, Haskell WL, Ades PA, Amsterdam EA, Bittner V, Franklin BA, Gulanick M, Laing ST, Stewart KJ. Resistance exercise in individuals with and without cardiovascular disease: 2007 update: a scientific statement from the American Heart Association Council on Clinical Cardiology and Council on Nutrition, Physical Activity, and Metabolism. *Circulation*. 2007 Jul 31;116(5):572-84. [doi:10.1161/CIRCULATIONAHA.107.185214] [PMid:17638929]
99. García-Hermoso A, Ramírez-Vélez R, Ramírez-Campillo R, Peterson MD, Martínez-Vizcaíno V. Concurrent aerobic plus resistance exercise versus aerobic exercise alone to improve health outcomes in paediatric obesity: a systematic review and meta-analysis. *British journal of sports medicine*. 2016 Dec 16. [doi:10.1136/bjsports-2016-096605] [PMid:27986760]
100. Moraleda BR, Morencos E, Peinado AB, Bermejo L, Candela CG, Benito PJ, PRONAF Study Group. Can the exercise mode determine lipid profile improvements in obese patients?. *Nutrición hospitalaria*. 2013;28(3):607-17.
101. Baldi J, Snowling NJ. *Josm*. Resistance training improves glycaemic control in obese type 2 diabetic men. 2003;24(06):419-23. [doi:10.1055/s-2003-41173] [PMid:12905089]
102. Stein RA, Michielli DW, Glantz MD, Sardy H, Cohen A, Goldberg N, Brown CD. Effects of different exercise training intensities on lipoprotein cholesterol fractions in healthy middle-aged men. *American heart journal*. 1990 Feb 1;119(2):277-83. [doi:10.1016/S0002-8703(05)80017-1] [PMid:2301216]
103. Kokkinos PF, Hurley BF, Vaccaro P, Patterson JC, Gardner LB, Ostrove SM, Goldberg AP. Effects of low-and high-repetition resistive training on lipoprotein-lipid profiles. *Medicine and science in sports and exercise*. 1988 Feb 1;20(1):50-4. [doi:10.1249/00005768-198802000-00007] [PMid:3343916]
104. Kokkinos PF, Hurley BF, Smutok MA, Farmer C, Reece C, Shulman R, Charabogous C, Patterson J, Will S, Devane-Bell J. Strength training does not improve lipoprotein-lipid profiles in men at risk for CHD. *Medicine and Science in Sports and Exercise*. 1991 Oct 1;23(10):1134-9. [doi:10.1249/00005768-199110000-00004]
105. Dâmaso AR, da Silveira Campos RM, Caranti DA, de Piano A, Fisberg M, Foschini D, de Lima Sanches P, Tock L, Lederman HM, Tufik S, de Mello MT. Aerobic plus resistance training was more effective in improving the visceral adiposity, metabolic profile and inflammatory markers than aerobic training in obese adolescents. *Journal of sports sciences*. 2014 Sep 14;32(15):1435-45. [doi:10.1080/02640414.2014.900692] [PMid:24730354]
106. Chaudhary S, Kang MK, Sandhu JS. The effects of aerobic versus resistance training on cardiovascular fitness in obese sedentary females. *Asian journal of sports medicine*. 2010 Dec;1(4):177. [doi:10.5812/asjasm.34835] [PMid:22375205]
107. Gomez-Tomas C, Chulvi-Medrano I, Carrasco JJ, Alakhdar Y. Effect of a 1-year elastic band resistance exercise program on cardiovascular risk profile in postmenopausal women. *Menopause*. 2018 Sep 1;25(9):1004-10. [doi:10.1097/GME.0000000000001113] [PMid:29787478]
108. Roh HT, Cho SY, So WY. A cross-sectional study evaluating the effects of resistance exercise on inflammation and neurotrophic factors in elderly women with obesity. *Journal of clinical medicine*. 2020 Mar 20;9(3):842. [doi:10.3390/jcm9030842] [PMid:32244926]
109. Hamasaki H, Kawashima Y, Tamada Y, Furuta M, Katsuyama H, Sako A, Yanai H. Associations of low-intensity resistance training with body composition and lipid profile in obese patients with type 2 diabetes. *PLoS One*. 2015 Jul 15;10(7):e0132959. [doi:10.1371/journal.pone.0132959] [PMid:26176700]
110. Hughes TA, Gwynne JT, Switzer BR, Herbst C, White G. Effects of caloric restriction and weight loss on glycemic control, insulin release and resistance, and atherosclerotic risk in obese patients with type II diabetes mellitus. *The American journal of medicine*. 1984 Jul 1;77(1):7-17. [doi:10.1016/0002-9343(84)90429-7] [PMid:6377892]
111. Wing RR, Jeffery RW, Burton LR, Thorson C, Kuller LH, Folsom AR. Change in waist-hip ratio with weight loss and its association with change in cardiovascular risk factors. *The American journal of clinical nutrition*. 1992 Jun 1;55(6):1086-92. [doi:10.1093/ajcn/55.6.1086] [PMid:1595579]

112. So AC, Richter WO, Schwandt P. Benefit from hypocaloric diet in obese men depends on the extent of weight-loss regarding cholesterol, and on a simultaneous change in body fat distribution regarding insulin sensitivity and glucose tolerance. *Metabolism*. 1992 Sep 1;41(9):1035-9. [doi:10.1016/0026-0495(92)90134-V] [PMid:1518420]
113. Noakes M, Clifton PM. Changes in plasma lipids and other cardiovascular risk factors during 3 energy-restricted diets differing in total fat and fatty acid composition. *The American journal of clinical nutrition*. 2000 Mar 1;71(3):706-12. [doi:10.1093/ajcn/71.3.706] [PMid:10702163]
114. Kasim-Karakas SE, Almario RU, Mueller WM, Peerson J. Changes in plasma lipoproteins during low-fat, high-carbohydrate diets: effects of energy intake. *The American journal of clinical nutrition*. 2000 Jun 1;71(6):1439-47. [doi:10.1093/ajcn/71.6.1439] [PMid:10837283]
115. Wooten JS, Phillips MD, Mitchell JB, Patrizi R, Pleasant RN, Hein RM, Menzies RD, Barbee JJ. Resistance exercise and lipoproteins in postmenopausal women. *International journal of sports medicine*. 2010 Nov 17;7-13. [doi:10.1055/s-0030-1268008] [PMid:21086242]
116. Fahlman MM, Boardley D, Lambert CP, Flynn MG. Effects of endurance training and resistance training on plasma lipoprotein profiles in elderly women. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*. 2002 Feb 1;57(2):B54-60. [doi:10.1093/gerona/57.2.B54] [PMid:11818424]
117. de Mello MT, de Piano A, Carnier J, Sanches PD, Corrêa FA, Tock L, Ernandes RM, Tufik S, Dâmaso AR. Long-term effects of aerobic plus resistance training on the metabolic syndrome and adiponectinemia in obese adolescents. *The journal of clinical hypertension*. 2011 May;13(5):343-50. [doi:10.1111/j.1751-7176.2010.00388.x] [PMid:21545395]
118. Kokkinos PF, Hurley BF. Strength training and lipoprotein-lipid profiles: a critical analysis and recommendations for further study. *Sports medicine*. 1990 May;9:266-72. [doi:10.2165/00007256-199009050-00002] [PMid:2188329]
119. Gavin C, Sigal RJ, Cousins M, Menard ML, Atkinson M, Khandwala F, Kenny GP, Proctor S, Ooi TC. Resistance exercise but not aerobic exercise lowers remnant-like lipoprotein particle cholesterol in type 2 diabetes: a randomized controlled trial. *Atherosclerosis*. 2010 Dec 1;213(2):552-7. [doi:10.1016/j.atherosclerosis.2010.08.071] [PMid:20870234]
120. Farrell PA, Maksud MG, Pollock ML, Foster C, Anholm J, Hare J, Leon AS. A comparison of plasma cholesterol, triglycerides, and high density lipoprotein-cholesterol in speed skaters, weightlifters and non-athletes. *European Journal of Applied Physiology and Occupational Physiology*. 1982 Feb;48:77-82. [doi:10.1007/BF00421167] [PMid:7199456]
121. Morgan DW, Cruise RJ, Girardin BW, Lutz-Schneider V, Morgan DH, Qi WM. HDL-C concentrations in weight-trained, endurance-trained, and sedentary females. *The Physician and Sportsmedicine*. 1986 Mar 1;14(3):166-81. [doi:10.1080/00913847.1986.11709017] [PMid:27467348]
122. Awad SF, O'Flaherty M, El-Nahas KG, Al-Hamaq AO, Critchley JA, Abu-Raddad LJ. Preventing type 2 diabetes mellitus in Qatar by reducing obesity, smoking, and physical inactivity: mathematical modeling analyses. *Population health metrics*. 2019 Dec;17(1):1-3. [doi:10.1186/s12963-019-0200-1] [PMid:31888689]
123. Mameletzi D, Kouidi E, Koutlianos N, Deligiannis A. Effects of long-term exercise training on cardiac baroreflex sensitivity in patients with coronary artery disease: a randomized controlled trial. *Clinical rehabilitation*. 2011 Mar;25(3):217-27. [doi:10.1177/0269215510380825] [PMid:20943717]
124. Ardoy DN, Artero EG, Ruiz Ruiz J, Labayen I, Sjöström M, Castillo Garzón MJ, et al. Effects on adolescents' lipid profile of a fitness-enhancing intervention in the school setting: the EDUFIT study. 2013.
125. Batrakoulis A, Jamurtas AZ, Draganidis D, Georgakouli K, Tsimeas P, Poullos A, et al. Hybrid Neuromuscular Training Improves Cardiometabolic Health and Alters Redox Status in Inactive Overweight and Obese Women: A Randomized Controlled Trial. *Antioxidants (Basel)*. 2021;10(10). [doi:10.3390/antiox10101601] [PMid:34679738]
126. Batrakoulis A, Jamurtas AZ, Metsios GS, Perivoliotis K, Liguori G, Feito Y, et al. Comparative Efficacy of 5 Exercise Types on Cardiometabolic Health in Overweight and Obese Adults: A Systematic Review and Network Meta-Analysis of 81 Randomized Controlled Trials. *Circ Cardiovasc Qual Outcomes*. 2022;15(6):5(6):e008243. [doi:10.1161/CIRCOUTCOMES.121.008243] [PMid:35477256]

127. Batrakoulis A, Jamurtas AZ, Fatouros IG. High-Intensity Interval Training in Metabolic Diseases: Physiological Adaptations. *ACSM's Health & Fitness Journal*. 2021;25(5):54-9. [doi:10.1249/FIT.0000000000000703]
128. Lorenzo C, Okoloise M, Williams K, Stern MP, Haffner SM. The metabolic syndrome as predictor of type 2 diabetes: the San Antonio heart study. *Diabetes care*. 2003;26(11):3153-9. [doi:10.2337/diacare.26.11.3153] [PMid:14578254]
129. Zarei M, Nakhzari Khodakheyr J, Rashidlamir A, Montazeri A. The effect of combined resistance aerobic exercise training on concentrations of asprosin and complement C1q tumor necrosis factor-related protein-1 in men with type 2 diabetes. *Sport Sciences for Health*. 2021:1-9. [doi:10.1007/s11332-021-00738-7]
130. Christos ZE, Tokmakidis SP, Volaklis KA, Kotsa K, Touvra A-M, Douda E, et al. Lipoprotein profile, glycemic control and physical fitness after strength and aerobic training in post-menopausal women with type 2 diabetes. *European journal of applied physiology*. 2009;106(6):901-7. [doi:10.1007/s00421-009-1078-6] [PMid:19458961]
131. Lehmann R, Vokac A, Niedermann K, Agosti K, Spinass G. Loss of abdominal fat and improvement of the cardiovascular risk profile by regular moderate exercise training in patients with NIDDM. *Diabetologia*. 1995;38(11):1313-9. [doi:10.1007/BF00401764] [PMid:8582541]
132. McAuley KA, Williams SM, Mann JI, Goulding A, Chisholm A, Wilson N, et al. Intensive lifestyle changes are necessary to improve insulin sensitivity: a randomized controlled trial. *Diabetes care*. 2002;25(3):445-52. [doi:10.2337/diacare.25.3.445] [PMid:11874928]
133. Rönnemaa T, Marniemi J, Puukka P, Kuusi T. Effects of long-term physical exercise on serum lipids, lipoproteins and lipid metabolizing enzymes in type 2 (non-insulin-dependent) diabetic patients. *Diabetes Research (Edinburgh, Scotland)*. 1988;7(2):79-84.
134. Ruderman NB, Ganda OP, Johansen K. The effect of physical training on glucose tolerance and plasma lipids in maturity-onset diabetes. *Diabetes*. 1979;28(Supplement_1):89-92. [doi:10.2337/diab.28.1.S89] [PMid:761721]
135. Dunstan DW, Puddey IB, Beilin LJ, Burke V, Morton AR, Stanton K. Effects of a short-term circuit weight training program on glycaemic control in NIDDM. *Diabetes research and clinical practice*. 1998;40(1):53-61. [doi:10.1016/S0168-8227(98)00027-8] [PMid:9699091]
136. Jorge MLMP, de Oliveira VN, Resende NM, Paraiso LF, Calixto A, Diniz ALD, et al. The effects of aerobic, resistance, and combined exercise on metabolic control, inflammatory markers, adipocytokines, and muscle insulin signaling in patients with type 2 diabetes mellitus. *Metabolism*. 2011;60(9):1244-52. [doi:10.1016/j.metabol.2011.01.006] [PMid:21377179]
137. Sabouri M, Hatami E, Pournemati P, Shabkhiz F. Inflammatory, antioxidant and glycemic status to different mode of high-intensity training in type 2 diabetes mellitus. *Molecular Biology Reports*. 2021;48(6):5291-304. [doi:10.1007/s11033-021-06539-y] [PMid:34228273]
138. Zarei M, Nakhzari Khodakheyr J, Rashidlamir A, Montazeri A. The effect of combined resistance aerobic exercise training on concentrations of asprosin and complement C1q tumor necrosis factor-related protein-1 in men with type 2 diabetes. *Sport Sciences for Health*. 2021;17(4):863-71. [doi:10.1007/s11332-021-00738-7]
139. Magalhães JP, Santos DA, Correia IR, Hetherington-Rauth M, Ribeiro R, Raposo JF, et al. Impact of combined training with different exercise intensities on inflammatory and lipid markers in type 2 diabetes: A secondary analysis from a 1-year randomized controlled trial. *Cardiovascular diabetology*. 2020;19(1):1-11. [doi:10.1186/s12933-020-01136-y] [PMid:33028418]
140. Tan S, Li W, Wang J. Effects of six months of combined aerobic and resistance training for elderly patients with a long history of type 2 diabetes. *Journal of sports science & medicine*. 2012;11(3):495.
141. Tokmakidis SP, Zois CE, Volaklis KA, Kotsa K, Touvra A-M. The effects of a combined strength and aerobic exercise program on glucose control and insulin action in women with type 2 diabetes. *European journal of applied physiology*. 2004;92:437-42. [doi:10.1007/s00421-004-1174-6] [PMid:15232701]
142. Kwon HR, Min KW, Ahn HJ, Seok HG, Lee JH, Park GS, et al. Effects of aerobic exercise vs. resistance training on endothelial function in women with type 2 diabetes mellitus. *Diabetes & metabolism journal*. 2011;35(4):364-73. [doi:10.4093/dmj.2011.35.4.364] [PMid:21977456]

143. Balducci S, Zanuso S, Nicolucci A, De Feo P, Cavallo S, Cardelli P, et al. Effect of an intensive exercise intervention strategy on modifiable cardiovascular risk factors in subjects with type 2 diabetes mellitus: a randomized controlled trial: the Italian Diabetes and Exercise Study (IDES). *Archives of internal medicine*. 2010;170(20):1794-803. [doi:10.1001/archinternmed.2010.380] [PMid:21059972]
144. Sigal RJ, Kenny GP, Boulé NG, Wells GA, Prud'homme D, Fortier M, et al. Effects of aerobic training, resistance training, or both on glycemic control in type 2 diabetes: a randomized trial. *Annals of internal medicine*. 2007;147(6):357-69. [doi:10.7326/0003-4819-147-6-200709180-00005] [PMid:17876019]
145. Dunstan DW, Daly RM, Owen N, Jolley D, De Courten M, Shaw J, et al. High-intensity resistance training improves glycemic control in older patients with type 2 diabetes. *Diabetes care*. 2002;25(10):1729-36. [doi:10.2337/diacare.25.10.1729] [PMid:12351469]
146. Azarbayjani MA, Abedi B, Peeri M, Rasae MJ, Stannard SR. Effects of combined aerobic and resistant training on lipid profile and glycemic control in sedentary men. *Int Med J*. 2014;21(2):132-6.
147. Yavari A, Najafipoor F, Aliasgarzadeh A, Niafar M, Mobasser M. Effect of aerobic exercise, resistance training or combined training on glycaemic control and cardio-vascular risk factors in patients with type 2 diabetes. *Biology of Sport*. 2012;29(2):135-43. [doi:10.5604/20831862.990466]
148. Gibbs BB, Dobrosielski DA, Bonekamp S, Stewart KJ, Clark JM. A randomized trial of exercise for blood pressure reduction in type 2 diabetes: effect on flow-mediated dilation and circulating biomarkers of endothelial function. *Atherosclerosis*. 2012;224(2):446-53. [doi:10.1016/j.atherosclerosis.2012.07.035] [PMid:22889573]
149. Duncan GE, Perri MG, Theriaque DW, Hutson AD, Eckel RH, Stacpoole PW. Exercise training, without weight loss, increases insulin sensitivity and postheparin plasma lipase activity in previously sedentary adults. *Diabetes care*. 2003;26(3):557-62. [doi:10.2337/diacare.26.3.557] [PMid:12610001]
150. Lambers S, Van Laethem C, Van Acker K, Calders P. Influence of combined exercise training on indices of obesity, diabetes and cardiovascular risk in type 2 diabetes patients. *Clinical Rehabilitation*. 2008;22(6):483-92. [doi:10.1177/0269215508084582] [PMid:18511528]
151. Annibalini G, Lucertini F, Agostini D, Vallorani L, Gioacchini A, Barbieri E, et al. Concurrent aerobic and resistance training has anti-inflammatory effects and increases both plasma and leukocyte levels of IGF-1 in late middle-aged type 2 diabetic patients. *Oxidative medicine and cellular longevity*. 2017;2017. [doi:10.1155/2017/3937842] [PMid:28713486]
152. Mann S, Beedie C, Jimenez A. Differential effects of aerobic exercise, resistance training and combined exercise modalities on cholesterol and the lipid profile: review, synthesis and recommendations. *Sports medicine*. 2014;44(2):211-21. [doi:10.1007/s40279-013-0110-5] [PMid:24174305]
153. Hansen D, Dendale P, Jonkers R, Beelen M, Manders R, Corluy L, et al. Continuous low-to moderate-intensity exercise training is as effective as moderate-to high-intensity exercise training at lowering blood HbA1c in obese type 2 diabetes patients. *Diabetologia*. 2009;52(9):1789-97. [doi:10.1007/s00125-009-1354-3] [PMid:19370339]
154. Gordon LA, Morrison EY, McGrowder DA, Young R, Fraser YTP, Zamora EM, et al. Effect of exercise therapy on lipid profile and oxidative stress indicators in patients with type 2 diabetes. *BMC complementary and alternative medicine*. 2008;8(1):1-10. [doi:10.1186/1472-6882-8-21] [PMid:18477407]
155. Wagner H, Degerblad M, Thorell A, Nygren J, Stahle A, Kuhl J, et al. Combined treatment with exercise training and acarbose improves metabolic control and cardiovascular risk factor profile in subjects with mild type 2 diabetes. *Diabetes Care*. 2006;29(7):1471-7. [doi:10.2337/dc05-2513] [PMid:16801564]
156. Maiorana A, O'Driscoll G, Cheetham C, Dembo L, Stanton K, Goodman C, et al. The effect of combined aerobic and resistance exercise training on vascular function in type 2 diabetes. *Journal of the American College of Cardiology*. 2001;38(3):860-6. [doi:10.1016/S0735-1097(01)01439-5] [PMid:11527646]
157. Maiorana A, O'Driscoll G, Goodman C, Taylor R, Green D. Combined aerobic and resistance exercise improves glycemic control and fitness in type 2 diabetes. *Diabetes research and clinical practice*. 2002;56(2):115-23. [doi:10.1016/S0168-8227(01)00368-0] [PMid:11891019]

158. Kim HJ, Lee JS, Kim CK. Effect of exercise training on muscle glucose transporter 4 protein and intramuscular lipid content in elderly men with impaired glucose tolerance. *European journal of applied physiology*. 2004;93(3):353-8. [doi:10.1007/s00421-004-1214-2] [PMid:15480742]
159. De Nardi AT, Tolves T, Lenzi TL, Signori LU, da Silva AMV. High-intensity interval training versus continuous training on physiological and metabolic variables in prediabetes and type 2 diabetes: a meta-analysis. *Diabetes research and clinical practice*. 2018;137:149-59. [doi:10.1016/j.diabres.2017.12.017] [PMid:29329778]
160. Terada T, Friesen A, Chahal BS, Bell GJ, McCargar LJ, Boulé NG. Feasibility and preliminary efficacy of high intensity interval training in type 2 diabetes. *Diabetes research and Clinical practice*. 2013;99(2):120-9. [doi:10.1016/j.diabres.2012.10.019] [PMid:23183390]
161. Maillard F, Rousset S, Pereira B, Traore A, Del Amaze PdP, Boirie Y, et al. High-intensity interval training reduces abdominal fat mass in postmenopausal women with type 2 diabetes. *Diabetes & metabolism*. 2016;42(6):433-41. [doi:10.1016/j.diabet.2016.07.031] [PMid:27567125]
162. Mitranun W, Deerochanawong C, Tanaka H, Suksom D. Continuous vs interval training on glycemic control and macro-and microvascular reactivity in type 2 diabetic patients. *Scandinavian journal of medicine & science in sports*. 2014;24(2):e69-e76. [doi:10.1111/sms.12112] [PMid:24102912]
163. Batrakoulis A. Psychophysiological Adaptations to Yoga Practice in Overweight and Obese Individuals: A Topical Review. *Diseases*. 2022;10(4). [doi:10.3390/diseases10040107] [PMid:36412601]
164. Batrakoulis A. Psychophysiological Adaptations to Pilates Training in Overweight and Obese Individuals: A Topical Review. *Diseases*. 2022;10(4). [doi:10.3390/diseases10040071] [PMid:36278570]
165. Batrakoulis A. Role of Mind-Body Fitness in Obesity. *Diseases*. 2022;11(1). [doi:10.3390/diseases11010001] [PMid:36648866]
166. Adıgüzel S, Aras D, Gülü M, Aldhahi MI, Alqahtani AS, Al-Mhanna SB. Comparative effectiveness of 10-week equipment-based pilates and diaphragmatic breathing exercise on heart rate variability and pulmonary function in young adult healthy women with normal BMI—a quasi-experimental study. *BMC Sports Science, Medicine and Rehabilitation*. 2023;15(1):82. [doi:10.1186/s13102-023-00693-5] [PMid:37434197]
167. Gordon LA, Morrison EY, McGrowder DA, Young R, Fraser YT, Zamora EM, Alexander-Lindo RL, Irving RR. Effect of exercise therapy on lipid profile and oxidative stress indicators in patients with type 2 diabetes. *BMC complementary and alternative medicine*. 2008 Dec;8(1):1-0. [doi:10.1186/1472-6882-8-21] [PMid:18477407]
168. Singh S, Malhotra V, Singh KP, Madhu SV, Tandon OP. Role of yoga in modifying certain cardiovascular functions in type 2 diabetic patients. *JAPI*. 2004 Mar;52:203-6.
169. Geer EB, Islam J, Buettner C. Mechanisms of glucocorticoid-induced insulin resistance: focus on adipose tissue function and lipid metabolism. *Endocrinology and Metabolism Clinics*. 2014 Mar 1;43(1):75-102. [doi:10.1016/j.ecl.2013.10.005] [PMid:24582093]
170. Morais JB, Severo JS, Beserra JB, de Oiveira AR, Cruz KJ, de Sousa Melo SR, do Nascimento GV, de Macedo GF, do Nascimento Marreiro D. Association between cortisol, insulin resistance and zinc in obesity: a mini-review. *Biological trace element research*. 2019 Oct 15;191:323-30. [doi:10.1007/s12011-018-1629-y] [PMid:30617901]
171. Manchanda SC, Narang R, Reddy KS, Sachdeva U, Prabhakaran D, Dharmanand S, Rajani M, Bijlani R. Retardation of coronary atherosclerosis with yoga lifestyle intervention. *The Journal of the Association of Physicians of India*. 2000 Jul 1;48(7):687-94.
172. Chrousos GP, Gold PW. A healthy body in a healthy mind—and vice versa—the damaging power of “uncontrollable” stress. *The Journal of Clinical Endocrinology & Metabolism*. 1998 Jun 1;83(6):1842-5. [doi:10.1210/jcem.83.6.4908] [PMid:9626106]
173. Mendonça FM, Silva MM, Borges-Canha M, Neves JS, Costa C, Cabral PM, Guerreiro V, Lourenço R, Meira P, Ferreira MJ, Salazar D. Statin Therapy Among Bariatric Patients: The Impact on Metabolic Outcomes and Diabetes Status. *Experimental and Clinical Endocrinology & Diabetes*. 2022 Aug;130(08):539-45. [doi:10.1055/a-1743-2335] [PMid:35320845]

174. Park CL, Finkelstein-Fox L, Sacco SJ, Braun TD, Lazar S. How does yoga reduce stress? A clinical trial testing psychological mechanisms. *Stress and Health*. 2021 Feb;37(1):116-26. [[doi:10.1002/smi.2977](https://doi.org/10.1002/smi.2977)] [[PMid:32797683](https://pubmed.ncbi.nlm.nih.gov/32797683/)]
175. Sandeep Bhattacharya UP. Improvement In Oxidative Status With Yogic Breathing In Young Healthy Males Sandeep Bhattacharya', Us Pandey And Ns Verma. *Indian J Physio! Pharmacol*. 2002;46(3):349-54.
176. Perez-De-Albeniz A, Holmes J. Meditation: Concepts, effects and uses in therapy. *International Journal of Psychotherapy*. 2000 Mar 1;5(1):49-58. [[doi:10.1080/13569080050020263](https://doi.org/10.1080/13569080050020263)]
177. Angelopoulos TJ, Robertson RJ, Goss FL, Metz KF, LaPorte RE. Effect of repeated exercise bouts on high density lipoprotein-cholesterol and its subfractions HDL2-C and HDL3-C. *Int J Sports Med*. 1993;14(4):196-201. [[doi:10.1055/s-2007-1021163](https://doi.org/10.1055/s-2007-1021163)] [[PMid:8325718](https://pubmed.ncbi.nlm.nih.gov/8325718/)]
178. Rebolledo-Cobos R, Florez AP, Sarmiento-Rubiano L, Ardila-Pereira L, Enriquez JB, Rodríguez-Delgado A. Acute effects of resistance and concurrent exercise on the lipid profile of postmenopausal women. *Campa d*. 2019 Mar:79.
179. Anuzzi G, Jansson E, Kaijser L, Holmquist L, Carlson LA. Increased removal rate of exogenous triglycerides after prolonged exercise in man: time course and effect of exercise duration. *Metabolism*. 1987 May 1;36(5):438-43. [[doi:10.1016/0026-0495\(87\)90040-0](https://doi.org/10.1016/0026-0495(87)90040-0)] [[PMid:3574133](https://pubmed.ncbi.nlm.nih.gov/3574133/)]
180. Griffin BA, Skinner ER, Maughan RJ. The acute effect of prolonged walking and dietary changes on plasma lipoprotein concentrations and high-density lipoprotein subfractions. *Metabolism*. 1988 Jun 1;37(6):535-41. [[doi:10.1016/0026-0495\(88\)90168-0](https://doi.org/10.1016/0026-0495(88)90168-0)] [[PMid:3163762](https://pubmed.ncbi.nlm.nih.gov/3163762/)]
181. Batrakoulis A, Fatouros IG, Chatzinikolaou A, Draganidis D, Georgakouli K, Papanikolaou K, et al. Dose-response effects of high-intensity interval neuromuscular exercise training on weight loss, performance, health and quality of life in inactive obese adults: Study rationale, design and methods of the DoIT trial. *Contemp Clin Trials Commun*. 2019;15:100386. [[doi:10.1016/j.conctc.2019.100386](https://doi.org/10.1016/j.conctc.2019.100386)] [[PMid:31193901](https://pubmed.ncbi.nlm.nih.gov/31193901/)]
182. Batrakoulis A, Jamurtas AZ, Fatouros IG. Exercise and type II diabetes mellitus: A brief guide for exercise professionals. *Strength Cond J*. 2022;44(6):64-72. [[doi:10.1519/SSC.0000000000000731](https://doi.org/10.1519/SSC.0000000000000731)]