

## REVIEW ARTICLE



# Treatment Methods According to Each Pain Region of the Body in Dancers

<sup>1</sup>Hea Kyung Choi \*

<sup>1</sup>Department of Dance and Arts, College of Humanities and Convergence Arts, Sungshin Women's University, Seoul, Korea.

Submitted January 12, 2024; Accepted in final form March 19, 2024.

## ABSTRACT

**Background.** Dancers experience frequent pain and injuries as they perform the same daily training and movements. Therefore, if pain caused by repeated movements is treated early, training effectiveness will improve, leading to improved performance. **Objectives.** This study aims to suggest a method of treating pain based on published papers that have been studied so far. **Methods.** This study has searched the scientific literature published between 2000 and 2023. Indexed terms and text words include dance, dancer, sports injury, musculoskeletal pain, treatment, initial treatment, recovery, and dance effect. **Results.** Pain was prevalent in the head, lower back, foot, and legs, including the hip joint, and injuries were also concentrated in the lower extremity and lower back region. **Conclusion.** The treatment method proposed in this study is expected to help treat pain in dancers through analysis of the structural causes, considering that the cutaneous nerve, which quickly feels pain, is a treatment target that should relax the structures in the path to the superficial fascia.

**KEYWORDS:** *Dancers, Treatment, Musculoskeletal Pain, Headache, Injury.*

## INTRODUCTION

Dancers often experience physical injuries or pain due to excessive joint movements and practice for durability, which causes them to be unable to demonstrate their skills in actual performances. The dancers' most common musculoskeletal pain or injuries are in the lower extremities. The incidence of thigh injury in dancers is approximately 20%, with musculoskeletal region and pain having even higher incidences (1). In particular, it has been reported that excessive stretching can cause hamstring injuries in dancers, with 12-31% cases of reinjury (2). Other studies reported that the injury region of the lower extremity includes the quadriceps, adductors, iliotibial band, and bone (3).

Although there are several experimental studies to show that dance has a positive effect on

the central nervous system, such as Parkinson's, and to prevent injuries, it should be considered necessary in sports science that frequent injuries have a negative impact in terms of performance and treatment purposes (4-9). For the dancer who practices dance every day, the best way to prevent musculoskeletal injury or pain is to do well recovery after practice every day (10).

Pain ultimately comes from sensory nerves, and the cutaneous nerves are sensitive to pain. However, since the human body is not composed of muscles and nerves independently, it is necessary to analyze the whole body and think about the cause of the pain. Therefore, analyzing the area where pain is felt and the imbalance in the body is essential to understanding body balance and pain (11, 12).

---

\*. Corresponding Author:

Hea Kyung Choi, Ph.D.


E-mail: [queenndance@sungshin.ac.kr](mailto:queenndance@sungshin.ac.kr)

This study aims to suggest a method of treating pain based on published papers that have been studied.

## MATERIALS AND METHODS

This study has been searched in the scientific literature published between 2000 and 2023.

Indexed terms and text words include dance, dancer, sports injury, musculoskeletal pain, treatment, initial treatment, recovery, and dance effect. Then, this study examined the reference lists of all relevant studies. Then, for each study found, injuries, pain, and recovery methods were organized, made into one table, and analyzed (Figure 1).



**Figure 1.** The processing of this study.

## RESULTS

This study investigated where the dancers experienced pain or injury region. According to research in Sweden in 1994, the lower back, foot, ankle, and neck regions were painful. In a study reported in America between 2010 and 2017, the hamstring, adductor muscles, and iliotibial tract were reported as the injury region, and the lumbosacral, hip, and patellofemoral region were reported as the pain region. In 2019, the knee, lumbar, foot, and ankle were reported as injured areas in the UK. According to two studies reported in Germany, the pain regions were the head, trunk, and lower limb, which were the same in both studies. A Croatian study reported back, knee, and toe pain, and a study published in Irish reported pain in the knee, leg, and plantar region and was diagnosed with a problem with the foot bones. According to a recently published study in China, pain occurred in the Achilles, forefoot, and midfoot regions (Table 1).

## DISCUSSION

Back pain is a common symptom among dancers and a propounding health problem that increases with age and cannot be resolved by decreasing training intensity (13). According to another study (14), patients did not think much of the importance of diagnosis, drug treatment, or

surgery, and medical staff did not do enough to explain the cause of pain to patients. In many cases, finding the cause of back pain is difficult, so the primary approach is to find the tight region and apply pressure as a noninvasive treatment method (15, 16). Depending on the direction of attachment of the muscles around the lower back that attach to the lumbar vertebra and iliac crest, there is a method of palpating and gently pressing the area that feels a little hard and waiting until it loosens. Also, it is an excellent way to relieve it in the same way as above by palpating the hamstring muscles based on the origin of the hamstring muscles in the ischial tuberosity in the lower part of the pelvic bone. This is because back pain often occurs due to the shortening of the hamstrings (17, 18).

The pain in the head with the neck was included in previous studies (Table 1). Anatomically, structures sensitive to pain in the head region include the periosteum, meninges, and scalp, and the trigeminal nerve innervates as the sensory function (27). A few researchers reported that the reason for head pain was a trigger point of the near muscle of the head and neck (28-30). The greater and lesser occipital nerves that cause headaches are affected by the structures of the neck and shoulders, so in a headache, it would be good to find the trigger point in the above region and recover. The myodural bridge also might be

related to headaches (31). In the dura mater in the vertebral foramen, it is reflected that the relaxation of these muscles affects the circulation of cerebrospinal fluid and positively affects the purpose of recovery of the human body (31). The muscles forming the suboccipital triangle are attached to the superior and inferior nuchal line

based on the external occipital protuberance. Therefore, it is believed that relaxing muscles by providing physical stimulation centered on the above bony landmark will help treat headaches and improve the overall physical condition by promoting cerebrospinal fluid circulation (Figure 2; 32).

**Table 1. The pain and injury region**


population	Year	Pain or diagnosis	injury
American (3)	2010		Hamstring Iliotibial tract Adductor
Croatia (13)	2015	Back pain (53.5%) Knee (43%) Toe (40.5%)	
Irish (19)	2023	Ankle sprain Posterior tibialis tendonitis Patellofemoral pain syndrome Plantar fasciitis	Sesamoid, metatarsal, navicular, 1 <sup>st</sup> proximal phalanx, tibia
Sweden (20)	1994	Low Back Foot, ankle neck	
UK (21)	2019		Knee (36%) Lumbar (19%) Foot and ankle (15%)
American (22)	2017		Hip and groin
American (23)	2015	Lumbosacral pain (62%) Snapping hip (58%) Patellofemoral pain (29%)	Lower extremity (66%-91%) Foot and ankle (14%-57%)
Chinese (24)	2022	Archilles region Forefoot and midfoot	
Germany (25)	2019	Lower limb in a Ballet dancer Head and trunk in JMC	
Germany (26)	2021	Lower extremity Head Trunk	

JMC: jazz/modern/contemporary dancers.


The posterior femoral cutaneous nerve handles pain in the back of the leg. Since this nerve passes through the back of the thigh, relaxation of the hamstring muscles can effectively reduce pain in the region (33). According to Table 1, this study recommends relaxing the hamstring as a treatment method for feeling pain in the posterior leg. Structurally, depending on the condition of the hamstrings or fascia, abnormal sensations may be felt in the affected cutaneous nerve (Figure 3; 34).

According to another study (35), various causes of foot pain in dancers have been reported as interdigital neuromas, tarsal tunnel syndrome, medial hallucal nerve compression, anterior tarsal tunnel syndrome, superficial and deep peroneal nerve entrapment, and sural nerve entrapment (Figure 4). Although there are cases where the


exact cause of various pains cannot be diagnosed, tarsal tunnel syndrome is usually caused by the tibial nerve passing through the tarsal tunnel and becoming the plantar nerve, which is the nerve responsible for motor and sensory function on the plantar region (36). Therefore, pain in the plantar region or ankle is caused by the tarsal tunnel. This study thought there are many cases where the pressure increases and causes pain. The structures that pass through the tarsal tunnel are the tibial nerve, tibialis posterior, flexor digitorum longus, and flexor hallucis longus. For dancers with pain in the plantar or ankles, if the back of the tibia is tight when palpated, it is a good idea to relax this region (37). Although it is good to analyze the body type of a dancer in pain, it is also recommended to identify and treat the cutaneous nerve, which is responsible for the pain region.



**Figure 2.** The suboccipital structures are related to headaches. GAN, greater occipital nerve; SsC, semispinalis capitis; OCS, oblique capitis superior; RCPm, rectus capitis posterior minor; RCPM, rectus capitis posterior major; VA, vertebral artery; SON, suboccipital nerve; LC, longissimus capitis; SC, splenius capitis.



**Figure 3.** Photographs of the distribution of the sural cutaneous nerve. Arrow, branch out point of sural cutaneous nerve.



**Figure 4.** The structures that pass through the tarsal tunnel. TP, tibialis posterior; FDL, flexor digitorum longus; TN, tibial nerve; PTA, posterior tibial artery; FHL, flexor hallucis longus; CT, calcaneal tendon.

## CONCLUSION

As a result of collecting and analyzing papers on musculoskeletal pain and injury as conditioning methods for recovery, each method will be effective in the pain region. Early treatment for pain relief is expected to strengthen training and have a positive effect on athletic performance. If further research is conducted on the effectiveness of the recovery method presented this time, it can be expected that pain prevention will not lead to injury. The limitation of this study is that it did not treat patients with these symptoms in clinical practice and prove the results.

## APPLICABLE REMARKS

- When a dancer has a headache, it is recommended to check the tension of the suboccipital triangle; when there is back pain, check the tension of surrounding muscles such as the hamstring.
- When there is pain in the ankles and plantar region, it is a good idea to check the structure passing through the tarsal tunnel as the relaxation target.

## AUTHORS' CONTRIBUTIONS

Study concept and design: Hea Kyung Choi. Acquisition of data: Hea Kyung Choi. Analysis and interpretation of data: Hea Kyung Choi. Drafting the manuscript: Hea Kyung Choi. Critical revision of the manuscript for important intellectual content: Hea Kyung Choi. Statistical analysis: Hea Kyung Choi. Administrative, technical, and material support: Hea Kyung Choi. Study supervision: Hea Kyung Choi.

## CONFLICT OF INTEREST

The authors declare that they have no conflicts of interest.

## ETHICAL CONSIDERATION

This study protocol conforms to the ethical guidelines of the 1975 Declaration of Helsinki.

## FUNDING/SUPPORT

This study has no funding or support for the paper.

## ROLE OF THE SPONSOR

The funding organizations are public institutions and had no role in the design and conduct of the study.

**FINANCIAL DISCLOSURE**

This study has no financial interests related to the material in the manuscript.

**ARTIFICIAL INTELLIGENCE (AI) USE**

This study agrees with the journal's policy in this section.

**REFERENCES**

1. Jacobs CL, Hincapié CA, Cassidy JD. Musculoskeletal injuries and pain in dancers: a systematic review update. *Journal of dance medicine & science*. 2012;16(2):74-84. [doi:10.1177/1089313X1201600204]
2. Hincapié CA, Morton EJ, Cassidy JD. Musculoskeletal injuries and pain in dancers: a systematic review. *Archives of physical medicine and rehabilitation*. 2008;89(9):1819-1829. [doi:10.1016/j.apmr.2008.02.020] [PMid:18760170]
3. Deleget A. Overview of thigh injuries in dance. *Journal of Dance Medicine & Science*. 2010;14(3):97-102. [doi:10.1177/1089313X1001400304]
4. Ko MG, Lee MM, Song CH. A comparison of the effects of different stretching methods on flexibility, muscle activity, and pain threshold in ballet dancers; a preliminary randomized controlled trial. *Journal of Bodywork and Movement Therapies*. 2020;24(4):354-360. [doi:10.1016/j.jbmt.2020.06.019] [PMid:33218533]
5. Li H, Qiu X, Yang Z, Zhang Z, Wang G, Kim Y, Kim S. Effects of Cha-Cha Dance Training on the Balance Ability of the Healthy Elderly. *International Journal of Environmental Research and Public Health*. 2022;19(20):13535. [doi:10.3390/ijerph192013535] [PMid:36294115]
6. Hong JJ, Chang KT, Villinger F. The dynamics of T and B cells in lymph node during chronic HIV infection: TFH and HIV, unhappy dance partners?. *Frontiers in immunology*. 2016;7:522. [doi:10.3389/fimmu.2016.00522] [PMid:27920778]
7. Morrin N, Redding E. Acute effects of warm-up stretch protocols on balance, vertical jump height, and range of motion in dancers. *Journal of dance medicine & science*. 2013;17(1):34-40. [doi:10.12678/1089-313X.17.1.34] [PMid:23498355]
8. Lima CD, Brown LE, Ruas CV, Behm DG. Effects of static versus ballistic stretching on hamstring: quadriceps strength ratio and jump performance in ballet dancers and resistance trained women. *Journal of Dance Medicine & Science*. 2018;22(3):160-167. [doi:10.12678/1089-313X.22.3.160] [PMid:30139422]
9. Koh K, Park YS, Park DW, Shim JK. Dance training improves the CNS's ability to utilize the redundant degrees of freedom of the whole body. *Scientific Reports*. 2020;10(1):22197. [doi:10.1038/s41598-020-79081-9] [PMid:33335153]
10. Chamberlain R. Hip pain in adults: evaluation and differential diagnosis. *American family physician*. 2021;103(2):81-89.
11. Abdo H, Calvo EL, Lopez JM et al. Specialized cutaneous Schwann cells initiate pain sensation. *Science*. 2019;365(6454):695-699. [doi:10.1126/science.aax6452] [PMid:31416963]
12. Stucky C, Mikesell AR. Cutaneous pain in disorders affecting peripheral nerves. *Neuroscience Letter*. 2021;20:765. [doi:10.1016/j.neulet.2021.136233] [PMid:34506882]
13. Miletic D, Miletic A, Milavic B. Age-related progressive increase of lower back pain among male dance sport competitors. *Journal of Back and Musculoskeletal Rehabilitation*. 2015;28(3):551-560. [doi:10.3233/BMR-140555] [PMid:25391325]
14. Smuck M, Barrette K, Martinez-Ith A, Sultana G, Zheng P. What does the patient with back pain want? A comparison of patient preferences and physician assumptions. *The Spine Journal*. 2022;22(2):207-213. [doi:10.1016/j.spinee.2021.09.007] [PMid:34551322]
15. Maher C, Underwood M, Buchbinder R. Non-specific low back pain. *Lancet*. 2017;18:736-747. [doi:10.1016/S0140-6736(16)30970-9] [PMid:27745712]
16. Vlaeyen JW, Maher CG, Wiech K et al. Low back pain. *Nature Reviews Disease Primers*. 2018;13;4(1):52. [doi:10.1038/s41572-018-0052-1] [PMid:30546064]
17. Sadler SG, Spink MJ, Ho A et al. Restriction in lateral bending range of motion, lumbar lordosis, and hamstring flexibility predicts the development of low back pain: a systematic review of prospective cohort studies. *BMC Musculoskeletal Disorder*. 2017;5;18(1):179. [doi:10.1186/s12891-017-1534-0] [PMid:28476110]

18. Shamsi M, Mirzaei M, Shahsavari S, Safari A, Saeb M. Modeling the effect of static stretching and strengthening exercise in lengthened position on balance in low back pain subject with shortened hamstring: a randomized controlled clinical trial. *BMC musculoskeletal disorders*. 2020;21(1):809. [doi:10.1186/s12891-020-03823-z] [PMid:33276764]
19. Póvoa AR, Costa CM, Simões S, Azevedo AM, Oliveira R. Irish Dancing Injuries and Associated Risk Factors: A Systematic Review. *International Journal of Environmental Research and Public Health*. 2023;20(12):6190. [doi:10.3390/ijerph20126190] [PMid:37372775]
20. Ramel E, Moritz U. Self-reported musculoskeletal pain and discomfort in professional ballet dancers in Sweden. *Journal of Rehabilitation Medicine*. 1994;26(1):11-16. [doi:10.2340/1650197719942611116]
21. Jubb C, Bell L, Cimelli S, Wolman R. Injury patterns in hip hop dancers. *Journal of Dance Medicine & Science*. 2019;23(4):145-149. [doi:10.12678/1089-313X.23.4.145] [PMid:31775952]
22. Trentacosta N, Sugimoto D, Micheli LJ. Hip and groin injuries in dancers: a systematic review. *Sports Health*. 2017;9(5):422-427. [doi:10.1177/1941738117724159] [PMid:28783444]
23. Smith PJ, Gerrie BJ, Varner KE, McCulloch PC, Lintner DM, Harris JD. Incidence and prevalence of musculoskeletal injury in ballet: a systematic review. *Orthopaedic journal of sports medicine*. 2015;3(7):2325967115592621. [doi:10.1177/2325967115592621] [PMid:26673541]
24. Lai JH, Fung NP, Yeung ST, Siu RW, Pak NK, Surgenor B, Yung PS, Ling SK. Comparison of Dance-Related foot and ankle injuries among Pre-Professional ballet, contemporary, and Chinese dancers. *Journal of Dance Medicine & Science*. 2022;26(2):134-142. [doi:10.12678/1089-313X.061522f] [PMid:35287790]
25. Lampe J, Groneberg DA, Borgetto B, Ohlendorf D, Wanke EM. Assessment of musculoskeletal pain in dance focusing on dance-style related differences. *The Physician and sportsmedicine*. 2019;47(4):433-440. [doi:10.1080/00913847.2019.1613120] [PMid:31099296]
26. Wanke EM, Haenel J, Schoettker-Koeniger T, Groneberg DA. Determinants of pain intensity in physical education teachers focusing on dance teachers: a cross-sectional study. *International Journal of Environmental Research and Public Health*. 2021;18(4): 2193. [doi:10.3390/ijerph18042193] [PMid:33672286]
27. Robertson CE, Benarroch EE. The anatomy of head pain. *Handbook of Clinical Neurology*. 2023;198:41-60. [doi:10.1016/B978-0-12-823356-6.00001-9] [PMid:38043970]
28. Fernández-de-Las-Peñas C. Myofascial head pain. *Current pain and headache reports*. 2015;19(7):28. [doi:10.1007/s11916-015-0503-2] [PMid:26049772]
29. Do TP, Heldarskard GF, Kolding LT, Hvedstrup J, Schytz HW. Myofascial trigger points in migraine and tension-type headache. *The journal of headache and pain*. 2018;19(1):84. [doi:10.1186/s10194-018-0913-8] [PMid:30203398]
30. Fernández-de-las-Peñas C, Alonso-Blanco C, Cuadrado ML, Gerwin RD, Pareja JA. Myofascial trigger points and their relationship to headache clinical parameters in chronic tension-type headache. *Headache: The Journal of Head and Face Pain*. 2006;46(8):1264-1272. [doi:10.1111/j.1526-4610.2006.00440.x] [PMid:16942471]
31. Kitamura K, Cho KH, Yamamoto M, Ishii M, Murakami G, Rodríguez-Vázquez JF, Abe SI. Suboccipital myodural bridges revisited: Application to cervicogenic headaches. *Clinical Anatomy*. 2019;32(7):914-928. [doi:10.1002/ca.23411] [PMid:31116454]
32. Enix DE, Scali F, Pontell ME. The cervical myodural bridge, a review of literature and clinical implications. *The Journal of the Canadian Chiropractic Association*. 2014;58(2):184-192.
33. Feigl GC, Schmid M, Zahn PK, González CA, Litz RJ. The posterior femoral cutaneous nerve contributes significantly to sensory innervation of the lower leg: an anatomical investigation. *British Journal of Anaesthesia*. 2020;124(3):308-313. [doi:10.1016/j.bja.2019.10.026] [PMid:31973825]
34. Wilke J, Macchi V, De Caro R, Stecco C. Fascia thickness, aging and flexibility: is there an association?. *Journal of anatomy*. 2019; 234(1):43-49. [doi:10.1111/joa.12902] [PMid:30417344]
35. Kennedy JG, Baxter DE. Nerve disorders in dancers. *Clinics in sports medicine*. 2008;27(2):329-334. [doi:10.1016/j.csm.2008.01.001] [PMid:18346547]
36. Nelson SC. Tarsal tunnel syndrome. *Clinics in podiatric medicine and surgery*. 2021;38(2):131-141. [doi:10.1016/j.cpm.2020.12.001] [PMid:33745647]

37. Fortier LM, Leethy KN, Smith M et al. An update on posterior tarsal tunnel syndrome. Orthopedic reviews. 2022;31;14(4):35444. [[doi:10.52965/001c.35444](https://doi.org/10.52965/001c.35444)]