

Ann Appl Sport Sci 13(1): e1447, 2025. e-ISSN: 2322-4479; p-ISSN: 2476-4981



REVIEW ARTICLE

Impact of Aerobic, Resistance, and Combined Training on Blood Pressure in Patients with Chronic Diseases: A Narrative Review

¹Nouf H. Alkhamees¹⁰, ^{2,3}Sameer Badri Al-Mhanna¹⁰, ⁴Gerasimos V. Grivas¹⁰, ⁵Yuni Astuti¹⁰, ³Norsuhana Omar¹⁰*, ⁶Hafeez Abiola Afolabi¹⁰, ⁷Bishir Daku Abubakar¹⁰, ⁸Khalid Mohamed, ^{9,10}Alexios Batrakoulis¹⁰

¹Department of Rehabilitation Sciences, College of Health and Rehabilitation Sciences, Princess Nourah bint Abdulrahman University, P.O. Box 84428, Riyadh 11671, Saudi Arabia.

²Center for Global Health Research, Saveetha Medical College and Hospitals, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai 602105, India.

³Department of Physiology, School of Medical Sciences, Universiti Sains Malaysia, Kubang Kerian 16150, Kelantan, Malaysia.

⁸Staff Physician, Adult Stem Cell Transplant and Cellular Therapy, Department of Oncology, King Abdul-Aziz Medical City, Riyadh, Kingdom of Saudi Arabia.

⁹Department of Physical Education and Sport Science, School of Physical Education, Sport Science and Dietetics, University of Thessaly, Karies 42100, Trikala, Greece.

¹⁰School of Physical Education and Sport Science, Department of Physical Education and Sport Science, Komotini 69100, Greece.

*. Corresponding Author: Norsuhana Omar; E-mail: suhanakk@usm.my

Submitted September 01, 2024; Accepted November 04, 2024.



KEYWORDS

Physical Activity, Cardiovascular Health, Health Outcomes, Exercise Prescription.

ABSTRACT

Background. Hypertension, a major contributor to heart disease, is linked to severe complications such as heart attacks, strokes, myocardial infarctions, and chronic kidney disease. Various types of exercise present a promising strategy for lowering BP in both hypertensive and normotensive individuals, thereby reducing the risk of cardiovascular diseases. **Objectives.** This study aimed to assess the impact of aerobic, resistance, and combined training on BP regulation in patients with chronic diseases. **Methods.** A comprehensive search was conducted using relevant keywords across SCOPUS, PubMed, the Cochrane Central Database, and Google Scholar. Eighteen articles meeting predefined inclusion and exclusion criteria were selected for review. The search covered papers published from January 1, 1995, through May 31, 2024. **Results.** The analysis revealed that aerobic exercise performed three times per week for 10 weeks led to significant reductions in systolic (SBP) and diastolic (DBP) blood pressure in obese patients with cardiovascular disease. Aerobic exercise showed a meaningful alteration in BP by reducing systemic vascular resistance in patients with metabolic syndrome. Moreover, aerobic and resistance training were more effective in lowering mean arterial pressure in individuals with pre- and stage-1 hypertension and type 2 diabetes. Combined aerobic and resistance training, performed three times per week for 12 weeks, led to a significant reduction in both SBP and DBP. **Conclusion.** These findings highlight the crucial role of exercise therapy as a nonpharmacological approach to managing BP and reducing cardiovascular risk factors.

⁴Physical Education and Sports, Division of Humanities and Political Sciences, Hellenic Naval Academy, Piraeus, Greece.

⁵Faculty of Sport Science, Universitas Negeri Padang, Padang, Indonesia.

⁶Department of General Surgery, School of Medical Sciences, Universiti Sains Malaysia, Kelantan, Malaysia.

⁷Department of Human Physiology, Federal University Dutse, Jigawa, Nigeria.

INTRODUCTION

Hypertension is a significant risk factor for cardiovascular disease (1). Cardiovascular diseases encompass a range of conditions, including coronary artery disease (CAD), arrhythmias, cardiomyopathy, and valvular disease (2). Among these, hypertension is a primary driver, leading to serious complications such as heart attack, myocardial infarction, stroke, and chronic kidney disease (3).

Cardiovascular disease mortality and morbidity increase in a curvilinear fashion with rising BP, without evidence of a definitive risk threshold (4, 5). Hypertension, a critical component of metabolic syndrome, is typically defined by a systolic blood pressure (SBP) of 130 mm Hg or higher or a diastolic blood pressure (DBP) of 85 mm Hg or higher (6). High BP is a significant public health issue in the United States, affecting approximately 50 million Americans (7). Over the past 80 years, the increasing cardiovascular disease mortality rate has led to annual healthcare costs nearing \$320 billion, impacting around 60 million Americans (7). Therefore, Whelton et al. (8) stated that uncontrolled high BP is a major contributor to coronary heart disease, stroke, end-stage renal disease, and congestive heart failure. Clinical research consistently shows that reducing BP can lower cardiovascular mortality (9). Even a modest reduction of 2 mm Hg in the average DBP across the general population could substantially decrease the risk of diseases associated with elevated BP (9).

Exercise can reduce the risk of numerous health disorders and chronic diseases (10). It is broadly defined as any bodily movement produced by skeletal muscles that requires energy expenditure. Various forms of exercise include cycling, jogging, walking, and participating in sports and recreational activities. Physical inactivity is a significant risk factor for cardiovascular disease, with individuals who are less physically active having a 30% to 50% higher risk of developing hypertension (11). Clinical studies have shown that regular physical activity can reduce BP in hypertensive and normotensive individuals, independent of weight loss (12). For instance, a study found that older women with metabolic syndrome experienced reductions in systolic, diastolic, and mean BP, along with improvements in heart rate variability, 24 hours after a low-intensity resistance exercise session with 60% to 80% blood flow restriction (13). However, there is conflicting evidence regarding how exercise contributes to BP reduction in the general population and specific subgroups (14).

Many systematic reviews and meta-analyses have investigated the impact of diverse exercise modalities on blood pressure. The findings indicate that aerobic exercise is associated with significant reductions in SBP, although the effect on DBP is less pronounced. These observations suggest that aerobic exercise may contribute to a reduction in cardiovascular disease risk. Such aerobic exercise-induced adaptations may be primarily attributed to the significant improvements in cardiac output, oxygen uptake, nitric oxide production, and stress levels that have antihypertensive properties (15-17). It has been well-documented that improved endothelial function and autonomic regulation, as well as reduced oxidative stress, are considered to be the potential physiological mechanisms responsible for these positive adaptations in BP levels (18). These advantageous effects of aerobic exercise are critical for middle-aged and older adults struggling with lifestyle-related chronic diseases commonly linked to impaired metabolic health that increases the risk of mortality (19).

Resistance training that involves muscle contraction against external forces has been demonstrated to enhance muscle strength, facilitate fat loss, and regulate blood pressure. A literature review reveals that studies comparing the effects of aerobic exercise and resistance training on BP in individuals with hypertension indicate that both types of exercise are effective in lowering BP, with no significant difference between them (20). However, it is possible that resistance training may be more beneficial in populations with metabolic dysregulation, as it has been shown to induce favorable effects on various glycolipid metabolism indicators (21).

The objective of the present review was to provide a synthesis of the research methods employed and the findings reported in studies where blood pressure was examined. A review article may disseminate the main research findings on blood pressure responses to exercise in various clinical populations to identify research issues, considerations, and gaps in the literature. Given the paucity of research on blood pressure responses to exercise, this narrative review aimed to provide a comprehensive overview of the

existing evidence by capturing all published research on the topic, regardless of the study design. The review addressed the general question: What is known from published research about blood pressure adaptations to aerobic, resistance, and combined training in patients with chronic diseases?

MATERIALS AND METHODS

Protocol and Registration. This manuscript is a narrative review, which does not require protocol registration. Therefore, no registration code is applicable for this type of study.

Literature Search Strategy. The studies included in this review were selected based on freely accessible data. We identified relevant studies by searching multiple databases, including SCOPUS, PubMed, Cochrane Central Database, and Google Scholar.

Eligibility Criteria. Following a comprehensive literature search, all studies examining the effects of exercise on BP in patients with chronic diseases were reviewed. The search spanned papers published between January 1, 1995, and May 31, 2024, utilizing keywords such as "BP", "endurance training", "cardiovascular risk factors", "hypertension", "aerobic exercise", "meta-analysis", and "resistance training". Two authors (S.B. and N.K.) applied the PICOS method to assess the full texts of the identified studies and

establish the inclusion and exclusion criteria. In cases of disagreement, a third author (A.B.) was consulted to reach a final decision.

Study Selection. Two authors (S.B. and N.O.) conducted the study selection process and applied the inclusion and exclusion criteria through a stepwise evaluation of titles, abstracts, and full texts. In cases where discrepancies arose, a third author (W.G.) was consulted to discuss and finalize the decision on whether to include or exclude articles for the current review.

Methodological Quality Assessment. The quality of the included studies was evaluated using the Joanna Briggs Institute (JBI) critical appraisal checklist (JBI, 2017), which examines nine key criteria: (i) appropriate sampling frame, (ii) suitable sampling technique, (iii) sufficient sample size, (iv) clear description of study subjects and setting, (v) thorough data analysis, (vi) valid methods for the conditions studied, (vii) consistent measurement across participants, (viii) appropriate statistical analysis, and (ix) adequate response rate. Each item was rated as yes, no, unclear, or not applicable. A "yes" received a score of 1, while "no" and "unclear" received 0. The average score for each study was calculated, and those with scores above the mean were classified as high-quality, while those below were deemed low-quality. Studies were included or excluded based on this methodological quality assessment (Table 1).

Table 1. Inclusion and exclusion criteria were used for this review. Characteristics Inclusion criteria

Inclusion criteria	Exclusion criteria		
Original studies from a peer-reviewed journal systematic review and meta-analysis.	Non-peer-reviewed or non-original research Editorials, opinions, and discussions		
Full-text articles available Available in English	Studies without full texts In a language other than English		
Patients with Diabetes, hypertension, congestive heart failure, stroke, asthma, human immunodeficiency virus infection, acquired immune deficiency syndrome comorbidities, liver disease, and chronic obstructive pulmonary	Patients without a history of chronic diseases.		
	Original studies from a peer-reviewed journal systematic review and meta-analysis. Full-text articles available Available in English Patients with Diabetes, hypertension, congestive heart failure, stroke, asthma, human immunodeficiency virus infection, acquired immune deficiency syndrome comorbidities, liver		

RESULTS

The data from the studies were imported into the EndNote X9 citation manager. Articles were then reviewed for relevance to the review's objectives, and duplicates were removed. Initially, 2,735 articles were identified. Following the application of predefined inclusion and exclusion criteria, 17 articles met the established criteria and were selected for inclusion in this review (as detailed in Table 2 and Figure 1).




Figure 1. PRISMA flowchart for search strategy.

Our review of relevant studies indicates that aerobic exercise is associated with significant benefits for individuals with obesity and cardiovascular disease. For instance, Banz et al. (22) found that obese patients with cardiovascular disease who participated in 40 minutes of aerobic exercise three times a week for ten weeks, at intensities between 60% and 80% of their maximum heart rate, experienced significant reductions in both SBP and DBP. Similarly, Grant et al. (23) observed that overweight women also experienced a reduction in BP following aerobic exercise interventions.

Furthermore, a meta-analysis examining hypertensive and non-hypertensive adults aged 20 to 72 years demonstrated that engaging in aerobic exercise—specifically, three sessions per week, each lasting 40 minutes at 65% of maximum heart rate—resulted in significant reductions in both resting and daytime ambulatory BP over durations ranging from 4 to 52 weeks. This observed decrease in BP was linked to a reduction in systemic vascular resistance (24).

Additionally, aerobic exercise has been shown to lower both SBP and DBP in individuals with

metabolic syndrome, indicating its potential as an effective intervention for improving BP and alleviating chronic diseases (25).

Our literature review indicates that DBP significantly decreased after 16 and 32 weeks of exercise interventions, leading to reductions in both blood BP and left ventricular hypertrophy in individuals with severe hypertension (25). Maeda et al. (26) reported that exercise training resulted in elevated plasma concentrations of nitric oxide and cyclic guanosine monophosphate, which were associated with significant reductions in BP.

Similarly, Ruangthai and Phoemsapthawee (27) observed that combined aerobic and resistance training led to substantial decreases in SBP, correlated with increased plasma total nitrite/nitrate levels. Conversely, while aerobic and resistance training have been shown to reduce mean arterial pressure in prehypertension and stage-1 hypertension individuals, resistance training alone did not result in significant BP changes (28) (Table 2).

Table 2. Characteristics of the included studies.

Study	Total sample size/ Gender	Population	Age (Mean years ± SD)	Intervention	Duration (week)	Session per week	Main finding	Quality assessment JBI score out of 9 criteria
Whelton et al., 2002 (8)	A meta-analysis of 54 RCTs included 2419 subjects	hypertensive and non-hypertensive patients	-	Aerobic exercise	12	3	A significant decrease in mean systolic and diastolic BP was observed in both hypertensive and normotensive participants.	6
Banz et al., 2003 (22)	26/M	Obesity and at least one other heart disease	48±6	Aerobic and resistance training	10	3	No significant differences in BP were noted across groups; however, aerobic exercise significantly reduced DBP (P<0.05). Resistance training reduced systolic BP.	8
Grant et al., 2004 (23)	26/F	Overweight/obese hypertensive patients	63±4	Aerobic and resistance training	12	2	Both training significantly reduced DBP and SBP (P<0.05) compared to the control group	9
Fagard and Cornelissen, 2007 (24)	A meta-analysis of 105 studies included 72 trials	hypertensive and non-hypertensive	20 to 72	Aerobic exercise	4-52	3	The exercise intervention significantly decreased resting and daytime ambulatory BP (P<0.001) through decreased systemic vascular resistance.	6
Kokkinos et al., 1995 (25)	46/M	Hypertensive patients	35-76	Aerobic exercise	16	3	Intervention decreases DBP, BP, and left ventricular hypertrophy.	7
Maeda et al., 2004 (26)	15/F	Chronic diseases	59-69	Aerobic exercise	12	5	The intervention significantly reduced resting BP (P<0.05) and increased the plasma concentration of NO and cGMP.	7
Ruangthai et al., 2019 (27)	54/M	Hypertension	67±6 years	Aerobic and resistance training	12	3	Endurance training, combined training, and strength training reduced SBP by 7.9%, 8.2%, and 19.1%, respectively, correlating with increased plasma NO concentrations.	7
Collier et al., 2008 (28)	30/M & F	Hypertensive patients	48.2±1.3	Aerobic and resistance training	4	3	Significantly reduced resistance and aerobic training SBP and DBP. Resistance training increases central and peripheral arterial stiffness, while aerobic training decreases arterial stiffness.	8
Katzmarzyk et al., 2003 (29)	621/M & F	Chronic diseases, participants with BP<160/100 mm Hg)	45.8±13.9	Aerobic exercise	20	3	Aerobic exercise significantly decreases BP, SBP, and DBP (P<0.05).	8
Figueroa et al., 2011 (30)	24/F	Postmenopausal women	47-68	Circuit resistance training followed by aerobic exercise	12	-	Significantly reduced SBP and DBP compared with the control. The 12 weeks of moderate-intensity combined circuit resistance training and aerobic exercise improve hemodynamics, arterial stiffness, and muscle strength	8

Study	Total sample size/ Gender	Population	Age (Mean years ± SD)	Intervention	Duration (week)	Session per week	Main finding	Quality assessment JBI score out of 9 criteria
Dimeo et al., 2012 (31)	50/M & F	Hypertension	67.9±6.2	Aerobic exercise	12	3	It significantly reduced SBP and DBP (P=0.03). Exercise decreased BP on exertion and improved physical performance even in subjects with low responsiveness to medication.	8
Ho et al., 2011 (32)	20/F	Hypertensive women	50-67 years	Aerobic and resistance training	12	3	significantly reduced in SBP and DBP. A moderate-intensity 30-minute resistance or aerobic training bout could decrease cardiovascular disease risk factors.	9
Carpio-Rivera et al., 2016 (33)	A meta-analysis included 1408 participants	Chronic disease	36.1±15.1 years	Aerobic and resistance training	12	3	Intervention decreased significantly both DBP and SBP (p<0.05). The decrease in blood pressure was significant irrespective of the subjects' initial blood pressure levels, sex, or physical activity levels.	6
Park et al., 2021 (34)	-	Hypertension and T2DM.	-	Aerobic/resistance And combined training	-	-	Exercise improved SBP (95% CI, -8.39 to -2.12) and DBP (95% CI, -4.91 to -1.40). Aerobic activity showed the greatest impact on reducing SBP and DBP.	6
Lopes et al., 2021 (35)	A meta-analysis included 642 participants	hypertension	-	Aerobic, resistance, and combined aerobic and resistance	-	-	Exercise training, particularly aerobic, combined, and isometric resistance exercises, reduced pulse wave velocity in adults with hypertension; however, dynamic resistance training did not demonstrate this effect.	6
Al-Mhanna et al., 2024c (36)	A meta-analysis / 1,148 F	overweight/ obese T2DM patients	66±6	Aerobic and resistance training	12	3	The training intervention showed significant improvements in SBP and DBP.	6
Al-Mhanna et al., 2024d (37)	A meta-analysis. Included 1,192 participants	Cancer	57±7	Aerobic and resistance training	12	3	Combined aerobic and resistance training demonstrated significant improvements in SBP and DBP	6

BP: Blood Pressure; SBP: Systolic Blood Pressure; DBP: Diastolic Blood Pressure; JBI criteria: (1) appropriate sampling frame, (2) proper sampling technique, (3) adequate sample size, (4) study subject and setting description, (5) sufficient data analysis, (6) use of valid methods for the identified conditions, (7) valid measurement for all the participants, (8) using appropriate statistical analysis, and (9) adequate response rate; T2DM: Type 2 Diabetes Mellitus; F: Female; M: Male.

DISCUSSION

The study aimed to assess the effects of different exercise modalities on blood pressure in patients with chronic diseases. The analysis showed that consistent aerobic and resistance training significantly lowered SBP and DBP in individuals with cardiovascular Furthermore, the current evidence suggests that mind-body fitness modalities, such as Pilates, yoga, and tai chi, benefit BP and enhance physical fitness (38-40). A meta-analysis further indicated that aerobic exercise can significantly lower resting and daytime ambulatory BP, primarily due reduced systemic vascular resistance. Moreover, exercise-induced elevations in nitric oxide and cyclic guanosine monophosphate levels were identified as fundamental mechanisms contributing to BP reduction (41).

Exercise plays a crucial role in promoting the formation of new blood vessels and angiogenesis by increasing the expression of vascular endothelial growth factor (VEGF) stimulating blood flow to tissues. This enhanced blood vessel formation improves oxygen delivery and reduces vascular resistance, contributes to lowering blood pressure (42). Regular aerobic and resistance exercise can help regulate blood pressure by improving endothelial function and increasing nitric oxide production, leading to vasodilation. These effects collectively support cardiovascular health and help manage conditions like hypertension (42). Aerobic and resistance training have been shown to reduce mean arterial pressure in individuals with prehypertension and stage-1 hypertension, consistent with findings from previous studies (8). This finding may suggest that a combined approach involving aerobic and resistance training may be efficient in managing BP in individuals with hypertension or cardiovascular disease, as it potentially targets multiple physiological pathways involved regulation. MacDonald, Johnson (43) conducted a meta-analysis involving 64 trials and 2,344 participants, demonstrating that resistance training is equally as practical as aerobic training in improving SBP, DBP, and overall health outcomes.

Exercise therapy is a well-established nonpharmacological intervention for managing hypertension and other cardiovascular disorders. Among the various forms of exercise, aerobic continuous training is endorsed by international

guidelines for primary and secondary hypertension prevention (44). A notable phenomenon associated with exercise is postexercise hypotension (PEH), where BP decreases to levels below the pre-exercise baseline following a single bout of exercise. This transient BP reduction is considered a predictor of the longterm effects of exercise training on chronic hypertension (44). The occurrence and magnitude of PEH have been observed in both normotensive and hypertensive individuals, with varying observation periods following exercise (45). The precise mechanisms underlying the reduction in BP associated with exercise remain unclear. However, insulin resistance and hyperinsulinemia have been implicated in the development of hypertension trials (46).Clinical demonstrated that aerobic exercise improves insulin resistance and reduces insulin levels in hypertensive patients (47). Moreover, changes in BP during exercise have been closely associated with decreased overall cholesterol levels and improved insulin resistance (48).

proposed mechanism antihypertensive effects of exercise is the enhancement endothelium-dependent of vasodilation. Aerobic exercise has increased nitric oxide bioavailability, improving endothelium-dependent vasorelaxation hypertensive individuals (48). Reductions in SBP due to exercise are linked to increases in plasma total nitrite/nitrate concentrations. Additionally, previous research indicates that aerobic exercise can lower BP in pre-hypertensive hypertensive populations (46). The endothelial adaptation associated with exercise is believed to be driven by exercise-induced increases in shear stress, which result in the upregulation of endothelial nitric oxide synthase and greater nitric oxide bioavailability (49). It has been suggested that this adaptation is primarily due to a significant increase in nitric oxide production and/or a decrease in nitric oxide scavenging by reactive oxygen species (49). Furthermore, strength training has been reported to enhance endothelial-dependent vasodilation, although this effect appears localized to the trained limbs (50).

Circuit resistance training followed by endurance exercise at 60% of maximal heart rate performed three times per week for 12 weeks has been shown to significantly reduce SBP and DBP while improving hemodynamics, arterial stiffness, and muscle strength compared to a

control group (30). Circuit-based resistance training interventions conducted in various settings demonstrated favorable changes in cardiovascular disease risk factors, including BP (51-53). Recent systematic reviews and metaanalyses on the effects of exercise in adults with both hypertension and type 2 diabetes mellitus (T2DM) have further confirmed these benefits. Specifically, exercise interventions appear to improve SBP and DBP, with aerobic exercise being the most effective modality. Moreover, combined aerobic and isometric resistance training has been shown to reduce pulse wave velocity, reducing arterial stiffness (34). These findings suggest that exercise reduces BP by lowering systemic vascular resistance, potentially modulating the sympathetic nervous system and renin-angiotensin the system, collectively positively impacting cardiovascular risk factors. The antihypertensive effects of exercise appear to be more pronounced in hypertensive individuals than those with normal BP, indicating that regular physical activity helps regulate BP in hypertensive patients and may also serve as a preventative measure against hypertension in normotensive individuals (24). However, it is essential to note that not all studies report significant reductions in BP following exercise interventions. For instance, García-Hermoso et al. (54) found that higher-intensity exercise did not produce a greater antihypertensive effect in patients with chronic kidney disease, highlighting the potential influence of underlying health conditions on exercise outcomes. These results underscore the importance of tailored exercise interventions for managing BP and reducing cardiovascular disease risk in hypertensive and individuals, contributing normotensive improved quality of life and long-term health outcomes. It is also noteworthy that interval training programs appear to be a promising exercise strategy for inducing beneficial alterations in SBP and DBP in individuals with metabolic derangements (55, 56). Such positive effects may be attributed to the advantageous role of vigorous-intensity, intermittent-type exercise regimens in several mental health indicators associated with vascular function (57). This highlights the need for further research in this particular area.

Current American and European hypertension guidelines emphasize the importance of regular physical activity as a primary recommendation for

managing high BP (58). Engaging in physical activity has been demonstrated to lower BP, even in individuals who exhibit a limited response to medical treatments, highlighting its critical role in managing resistant hypertension (58). Therefore, exercise intervention is fundamental in treating and preventing chronic diseases, significantly impacting BP control. The current evidence base supports the recommendation of engaging in moderate-intensity physical activity for minimum of 30 minutes daily, with the optimal approach being daily participation. This regimen should include aerobic exercise modalities daily (24). Prior to undertaking vigorous exercise (>60% of maximum intensity), individuals hypertension should undergo medically supervised peak or symptom-limited exercise testing, incorporating an electrocardiogram and BP monitoring (59). Asymptomatic individuals at low risk participating in light-to-moderate physical activity (<60% of maximum intensity) generally do not require additional testing beyond routine care. However, asymptomatic high-risk patients may benefit from exercise testing before engaging in moderate-intensity activities (40-60% of maximum intensity), though not for light or very light activities (<40% of maximum intensity). Patients experiencing symptoms such as chest discomfort, exertional dyspnea, or palpitations should be closely monitored and may need further evaluation with tests like Holter monitoring, echocardiography, or a combination of these. physical activity, Alongside patients encouraged to follow non-pharmacological strategies, including increasing fruit and vegetable reducing salt and saturated intake, consumption, quitting smoking, limiting alcohol intake, and incorporating daily moderate exercise. For those at elevated or high risk of cardiovascular prompt complications, initiation antihypertensive drug therapy is recommended (24). For BP prevention and management, moderate-intensity whole-body aerobic exercise (50–65% of maximum oxygen intake) focusing on major muscle groups is recommended. The optimal duration for each session is between 30 and 60 minutes, with a frequency of three to four times per week (60). Additionally, resistance training should be conducted at low intensity without breath-holding. Individuals hypertension who experience chest problems, such as chest pain, should avoid resistance training (60). Thus, clinicians and practitioners should be aware of the latest research findings on the pivotal role of various exercise types in resting cardiovascular function parameters among patients with several lifestyle-related chronic diseases (61). Future research should explore the long-term effects of various exercise modalities on blood pressure regulation and vascular health in diverse populations with chronic diseases. Investigating the molecular mechanisms, such as nitric oxide production and VEGF expression, could provide deeper insights into the role of exercise in cardiovascular disease management. Randomized controlled trials with larger sample sizes and extended follow-up periods would strengthen the evidence for tailored exercise prescriptions.

CONCLUSION

The study compiles evidence on the effects of different types of exercise on BP. Aerobic and resistance training have been demonstrated to confer significant benefits for individuals with obesity and cardiovascular disease, leading to notable reductions in SBP and DBP. Combined aerobic and resistance training may provide the most effective strategy for managing BP in individuals with hypertension or cardiovascular disease. These findings highlight the importance of exercise therapy as a non-pharmacological treatment, potentially benefiting both normotensive and hypertensive individuals by regulating BP and reducing cardiovascular risk factors.

APPLICABLE REMARKS

- The study demonstrates that both aerobic and resistance training, when performed consistently, can effectively lower systolic and diastolic blood pressure in patients with chronic conditions, emphasizing the potential of exercise therapy in managing cardiovascular risk factors.
- Aerobic exercise, in particular, significantly reduces systemic vascular resistance, making it a viable non-pharmacological intervention for patients with metabolic syndrome and those at risk of developing cardiovascular diseases.
- Combining aerobic and circuit-based resistance training over 12 weeks was particularly beneficial in reducing mean arterial pressure, especially in individuals with pre- and stage-1 hypertension and type 2 diabetes.
- These findings reinforce the importance of integrating structured exercise programs into the treatment plans for hypertensive patients, offering a complementary approach to

traditional pharmacological treatments to manage blood pressure better and mitigate cardiovascular risks.

ACKNOWLEDGMENTS

This study was supported by the School of Medical Sciences, Universiti Sains Malaysia, grant number 304.PPSP.6315639.

AUTHORS' CONTRIBUTIONS

Study concept and design: Norsuhana Omar, Sameer Badri Al-mhanna, Alexios Batrakoulis, Hafeez Abiola Afolabi. Nouf H. Alkhamees. Acquisition of data: Sameer Badri Al-mhanna, Gerasimos V. Grivas, Nouf H. Alkhamees. Analysis and interpretation of data: Bishir Daku Abubakar, Khalid Mohamed, Norsuhana Omar. Drafting the manuscript: Sameer Badri Almhanna, Bishir Daku Abubakar. Critical revision of the manuscript for important intellectual content: Alexios Batrakoulis, Gerasimos V. Grivas, Yuni Astuti. Administrative, technical, and material support: Norsuhana Omar, Sameer Badri Al-mhanna, Alexios Batrakoulis, Hafeez Abiola Afolabi, Nouf H. Alkhamees. Study supervision: Yuni Astuti, Bishir Daku Abubakar, Alexios Batrakoulis.

CONFLICT OF INTEREST

The authors reported no potential conflict of interest.

FINANCIAL DISCLOSURE

The authors declare that they have no financial interests or relationships that could influence the content or outcomes of this study. No financial conflicts of interest are associated with this research.

FUNDING/SUPPORT

This research was supported by the School of Medical Sciences of Universiti Sains Malaysia. The funding body had no involvement in the design, execution, interpretation, or publication of this research.

ETHICAL CONSIDERATION

Not applicable.

ROLE OF THE SPONSOR

The sponsor had no role in the study design, data collection, data analysis, interpretation of results, manuscript preparation, or decision to submit the article for publication.

ARTIFICIAL INTELLIGENCE (AI) USE

Artificial intelligence tools were not utilized in this manuscript's conceptualization, data analysis, or writing, except for general-purpose language models used for proofreading or editing assistance, where applicable.

REFERENCES

- 1. Garovic VD, Hayman SR. Hypertension in pregnancy: an emerging risk factor for cardiovascular disease. Nature clinical practice Nephrology. 2007;3(11):613-22. [doi:10.1038/ncpneph0623] [PMid:17957198]
- 2. Mutalub YB, Abdulwahab M, Mohammed A, Yahkub AM, Al-Mhanna SB, Yusof W, et al. Gut microbiota modulation as a novel therapeutic strategy in cardiometabolic diseases. Foods. 2022;11(17):2575. [doi:10.3390/foods11172575] [PMid:36076760]
- 3. Tonelli M, Wiebe N, Guthrie B, James MT, Quan H, Fortin M, et al. Comorbidity as a driver of adverse outcomes in people with chronic kidney disease. Kidney international. 2015;88(4):859-66. [doi:10.1038/ki.2015.228] [PMid:26221754]
- 4. Al-Mhanna SB, Wan Ghazali WS, Maqsood A, Mohamed M, Ahmed N, Afolabi HA, et al. Physical activities pre-and post-COVID-19 vaccination and its implementations: A narrative review. SAGE Open Med. 2023;11:20503121231158981. [doi:10.1177/20503121231158981] [PMid:36909796]
- 5. Riyahi-Alam S, Derakhshan A, Mansournia M, Mansournia N, Almasi M, Azizi F, Hadaegh F. Blood pressure and cardiovascular morbidity risk in type 2 diabetes with hypertension over a decade of follow-up: evidence for J-shaped phenomenon. Journal of human hypertension. 2017;31(6):415-21. [doi:10.1038/jhh.2016.94] [PMid:28124685]
- 6. Cesena FH, Nary FC, Santos RD, Bittencourt MS. The contribution of the systolic and diastolic components for the diagnosis of arterial hypertension under the 2017 ACC/AHA Guideline and metabolic heterogeneity among individuals with Stage 1 hypertension. The Journal of Clinical Hypertension. 2020;22(7):1192-9. [doi:10.1111/jch.13907] [PMid:32558248]
- 7. Zhou B, Perel P, Mensah GA, Ezzati M. Global epidemiology, health burden and effective interventions for elevated blood pressure and hypertension. Nature Reviews Cardiology. 2021;18(11):785-802. [doi:10.1038/s41569-021-00559-8] [PMid:34050340]
- 8. Whelton SP, Chin A, Xin X, He J. Effect of aerobic exercise on blood pressure: a meta-analysis of randomized, controlled trials. Annals of internal medicine. 2002;136(7):493-503. [doi:10.7326/0003-4819-136-7-200204020-00006] [PMid:11926784]
- 9. Bundy JD, Li C, Stuchlik P, Bu X, Kelly TN, Mills KT, et al. Systolic blood pressure reduction and risk of cardiovascular disease and mortality: a systematic review and network meta-analysis. JAMA cardiology. 2017;2(7):775-81. [doi:10.1001/jamacardio.2017.1421] [PMid:28564682]
- 10.Al-Mhanna SB, Wan Ghazali WS, Mohamed M, Mohamed RN, Mirza MB, Basheer SN, et al. The Impact of COVID-19 on Physical Activity Patterns of Dental Students: A Multinational Survey. Healthcare (Basel). 2022;10(11). [doi:10.3390/healthcare10112140] [PMid:36360481]
- 11. Tanveer M, Asghar E, Tanveer U, Roy N, Zeba A, Al-Mhanna SB, Batrakoulis A. Community-Level Physical Activity Opportunities, Safe and Supportive Environment Factors, and Their Association with Overweight and Obesity Among School-Aged Children and Adolescents in Pakistan-A Cross-Sectional Study. Kurdish Studies. 2024;12(2):6425-32. [doi:10.3390/sports12090235] [PMid:39330712]
- 12.Blumenthal JA, Sherwood A, Gullette EC, Babyak M, Waugh R, Georgiades A, et al. Exercise and weight loss reduce blood pressure in men and women with mild hypertension: effects on cardiovascular, metabolic, and hemodynamic functioning. Archives of internal medicine. 2000;160(13):1947-58. [doi:10.1001/archinte.160.13.1947] [PMid:10888969]
- 13.Maciel AW, Pinto LM, Campos RC, Ferreira AC, Dias-Filho CA, Dias CJ, et al. Acute effects of resistance exercise with blood flow restriction in elderly women: A pilot study. Journal of aging and physical activity. 2020;29(3):361-71. [doi:10.1123/japa.2020-0137] [PMid:33373977]
- 14.Kayhan R, Bayrakdaroğlu S, Ceylan H, Eken Ö, Bayrakdaroğlu Y, Badicu G, et al. Effects of different rest intervals in high intensity interval training programs on VO₂max, body composition, and isokinetic strength and power. Journal of Men's Health. 2024;20(5). [doi:10.22514/jomh.2024.064]
- 15.Saco-Ledo G, Valenzuela PL, Ruiz-Hurtado G, Ruilope LM, Lucia A. Exercise reduces ambulatory blood pressure in patients with hypertension: a systematic review and meta-analysis of randomized

- controlled trials. Journal of the American Heart Association. 2020;9(24):e018487. [doi:10.1161/JAHA.120.018487] [PMid:33280503]
- 16.Jabbarzadeh Ganjeh B, Zeraattalab-Motlagh S, Jayedi A, Daneshvar M, Gohari Z, Norouziasl R, et al. Effects of aerobic exercise on blood pressure in patients with hypertension: a systematic review and dose-response meta-analysis of randomized trials. Hypertension Research. 2024;47(2):385-98. [doi:10.1038/s41440-023-01467-9] [PMid:37872373]
- 17. Edwards JJ, Deenmamode AHP, Griffiths M, Arnold O, Cooper NJ, Wiles JD, Driscoll JM. Exercise training and resting blood pressure: a large-scale pairwise and network meta-analysis of randomised controlled trials. British Journal of Sports Medicine. 2023;57(20):1317. [doi:10.1136/bjsports-2022-106503]
- 18.Lee J, Vasan RS, Xanthakis V. Association of blood pressure responses to submaximal exercise in midlife with the incidence of cardiovascular outcomes and all-cause mortality: the Framingham Heart Study. Journal of the American Heart Association. 2020;9(11):e015554. [doi:10.1161/JAHA.119.015554] [PMid:32431193]
- 19.Al-Mhanna SB, Rocha-Rodriguesc S, Mohamed M, Batrakoulis A, Aldhahi MI, Afolabi HA, et al. Effects of combined aerobic exercise and diet on cardiometabolic health in patients with obesity and type 2 diabetes: a systematic review and meta-analysis. BMC Sports Science, Medicine and Rehabilitation. 2023;15(1):165. [doi:10.1186/s13102-023-00766-5] [PMid:38049873]
- 20. Paluch AE, Boyer WR, Franklin BA, Laddu D, Lobelo F, Lee D-c, et al. Resistance exercise training in individuals with and without cardiovascular disease: 2023 update: a scientific statement from the American Heart Association. Circulation. 2024;149(3):e217-e31. [doi:10.1161/CIR.0000000000001189]
- 21. Sundell J. Resistance Training Is an Effective Tool against Metabolic and Frailty Syndromes. Adv Prev Med. 2011;2011:984683. [doi:10.4061/2011/984683] [PMid:21991450]
- 22.Banz WJ, Maher MA, Thompson WG, Bassett DR, Moore W, Ashraf M, et al. Effects of resistance versus aerobic training on coronary artery disease risk factors. Experimental biology and medicine. 2003;228(4):434-40. [doi:10.1177/153537020322800414] [PMid:12671188]
- 23.Grant S, Todd K, Aitchison T, Kelly P, Stoddart D. The effects of a 12-week group exercise programme on physiological and psychological variables and function in overweight women. Public Health. 2004;118(1):31-42. [doi:10.1016/S0033-3506(03)00131-8] [PMid:14643625]
- 24.Fagard RH, Cornelissen VA. Effect of exercise on blood pressure control in hypertensive patients. European Journal of Preventive Cardiology. 2007;14(1):12-7. [doi:10.1097/HJR.0b013e3280128bbb] [PMid:17301622]
- 25.Kokkinos PF, Narayan P, Colleran JA, Pittaras A, Notargiacomo A, Reda D, Papademetriou V. Effects of regular exercise on blood pressure and left ventricular hypertrophy in African-American men with severe hypertension. New England Journal of Medicine. 1995;333(22):1462-7. [doi:10.1056/NEJM199511303332204] [PMid:7477146]
- 26.Maeda S, Tanabe T, Otsuki T, Sugawara J, Iemitsu M, Miyauchi T, et al. Moderate regular exercise increases basal production of nitric oxide in elderly women. Hypertension Research. 2004;27(12):947-53. [doi:10.1291/hypres.27.947] [PMid:15894835]
- 27.Ruangthai R, Phoemsapthawee J. Combined exercise training improves blood pressure and antioxidant capacity in elderly individuals with hypertension. Journal of Exercise Science & Fitness. 2019;17(2):67-76. [doi:10.1016/j.jesf.2019.03.001] [PMid:30949214]
- 28.Collier S, Kanaley J, Carhart R, Frechette V, Tobin M, Hall A, et al. Effect of 4 weeks of aerobic or resistance exercise training on arterial stiffness, blood flow and blood pressure in pre-and stage-1 hypertensives. Journal of human hypertension. 2008;22(10):678-86. [doi:10.1038/jhh.2008.36] [PMid:18432253]
- 29. Katzmarzyk PT, Leon AS, Wilmore JH, Skinner JS, Rao D, Rankinen T, Bouchard C. Targeting the metabolic syndrome with exercise: evidence from the HERITAGE Family Study. Medicine & Science in Sports & Exercise. 2003;35(10):1703-9. [doi:10.1249/01.MSS.0000089337.73244.9B] [PMid:14523308]
- 30. Figueroa A, Park SY, Seo DY, Sanchez-Gonzalez MA, Baek YH. Combined resistance and endurance exercise training improves arterial stiffness, blood pressure, and muscle strength in postmenopausal women. Menopause. 2011;18(9):980-4. [doi:10.1097/gme.0b013e3182135442] [PMid:21540753]

- 31.Dimeo F, Pagonas N, Seibert F, Arndt R, Zidek W, Westhoff TH. Aerobic exercise reduces blood pressure in resistant hypertension. Hypertension. 2012;60(3):653-8. [doi:10.1161/HYPERTENSIONAHA.112.197780] [PMid:22802220]
- 32.Ho SS, Dhaliwal SS, Hills A, Pal S. Acute exercise improves postprandial cardiovascular risk factors in overweight and obese individuals. Atherosclerosis. 2011;214(1):178-84. [doi:10.1016/j.atherosclerosis.2010.10.015] [PMid:21081229]
- 33. Carpio-Rivera E, Moncada-Jiménez J, Salazar-Rojas W, Solera-Herrera A. Acute effects of exercise on blood pressure: a meta-analytic investigation. Arquivos brasileiros de cardiologia. 2016;106:422-33. [doi:10.5935/abc.20160064] [PMid:27168471]
- 34.Park S, Kim J, Lee J. Effects of exercise intervention on adults with both hypertension and type 2 diabetes mellitus: a systematic review and meta-analysis. Journal of Cardiovascular Nursing. 2021;36(1):23-33. [doi:10.1097/JCN.0000000000000651] [PMid:32011348]
- 36.Al-Mhanna SB, Batrakoulis A, Norhayati MN, Mohamed M, Drenowatz C, Irekeola AA, et al. Combined Aerobic and Resistance Training Improves Body Composition, Alters Cardiometabolic Risk, and Ameliorates Cancer-Related Indicators in Breast Cancer Patients and Survivors with Overweight/Obesity: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. J Sports Sci Med. 2024;23(2):366-95. [doi:10.52082/jssm.2024.366] [PMid:38841642]
- 37.Al-Mhanna SB, Batrakoulis A, Wan Ghazali WS, Mohamed M, Aldayel A, Alhussain MH, et al. Effects of combined aerobic and resistance training on glycemic control, blood pressure, inflammation, cardiorespiratory fitness and quality of life in patients with type 2 diabetes and overweight/obesity: a systematic review and meta-analysis. PeerJ. 2024;12:e17525. [doi:10.7717/peerj.17525] [PMid:38887616]
- 38.Batrakoulis A. Psychophysiological Adaptations to Yoga Practice in Overweight and Obese Individuals: A Topical Review. Diseases. 2022;10(4). [doi:10.3390/diseases10040107] [PMid:36412601]
- 39.Batrakoulis A. Psychophysiological Adaptations to Pilates Training in Overweight and Obese Individuals: A Topical Review. Diseases. 2022;10(4). [doi:10.3390/diseases10040071] [PMid:36278570]
- 40.Batrakoulis A. Role of Mind-Body Fitness in Obesity. Diseases. 2022;11(1). [doi:10.3390/diseases11010001] [PMid:36648866]
- 41.Kelley GA, Kelley KA, Vu Tran Z. Aerobic exercise and resting blood pressure: a meta-analytic review of randomized, controlled trials. Preventive cardiology. 2001;4(2):73-80. [doi:10.1111/j.1520-037X.2001.00529.x] [PMid:11828203]
- 42.Ross M, Kargl CK, Ferguson R, Gavin TP, Hellsten Y. Exercise-induced skeletal muscle angiogenesis: impact of age, sex, angiocrines and cellular mediators. Eur J Appl Physiol. 2023;123(7):1415-32. [doi:10.1007/s00421-022-05128-6] [PMid:36715739]
- 43.MacDonald HV, Johnson BT, Huedo-Medina TB, Livingston J, Forsyth KC, Kraemer WJ, et al. Dynamic resistance training as stand-alone antihypertensive lifestyle therapy: a meta-analysis. Journal of the American Heart Association. 2016;5(10):e003231. [doi:10.1161/JAHA.116.003231] [PMid:27680663]
- 44.Iellamo F, Caminiti G, Montano M, Manzi V, Franchini A, Mancuso A, Volterrani M. Prolonged post-exercise hypotension: effects of different exercise modalities and training statuses in elderly patients with hypertension. International journal of environmental research and public health. 2021;18(6):3229. [doi:10.3390/ijerph18063229] [PMid:33804738]
- 45. Cardoso Jr CG, Gomides RS, Queiroz ACC, Pinto LG, da Silveira Lobo F, Tinucci T, et al. Acute and chronic effects of aerobic and resistance exercise on ambulatory blood pressure. Clinics. 2010;65(3):317-25. [doi:10.1590/S1807-59322010000300013] [PMid:20360924]
- 46.Roberts CK, Hevener AL, Barnard RJ. Metabolic syndrome and insulin resistance: underlying causes and modification by exercise training. Comprehensive physiology. 2013;3(1):1. [doi:10.1002/cphy.c110062] [PMid:23720280]

- 47.Brown MD, Moore GE, Korytkowski MT, McCole SD, Hagberg JM. Improvement of insulin sensitivity by short-term exercise training in hypertensive African American women. Hypertension. 1997;30(6):1549-53. [doi:10.1161/01.HYP.30.6.1549] [PMid:9403581]
- 48. Huot M, Arsenault BJ, Gaudreault V, Poirier P, Pérusse L, Tremblay A, et al. Insulin resistance, low cardiorespiratory fitness, and increased exercise blood pressure: contribution of abdominal obesity. Hypertension. 2011;58(6):1036-42. [doi:10.1161/HYPERTENSIONAHA.111.180349] [PMid:22025379]
- 49.Larsen MK, Matchkov VV. Hypertension and physical exercise: The role of oxidative stress. Medicina. 2016;52(1):19-27. [doi:10.1016/j.medici.2016.01.005] [PMid:26987496]
- 50.Higashi Y, Yoshizumi M. Exercise and endothelial function: role of endothelium-derived nitric oxide and oxidative stress in healthy subjects and hypertensive patients. Pharmacology & therapeutics. 2004;102(1):87-96. [doi:10.1016/j.pharmthera.2004.02.003] [PMid:15056500]
- 51.Al-Mhanna SB, Batrakoulis A, Mohamed M, Alkhamees NH, Sheeha BB, Ibrahim ZM, et al. Home-based circuit training improves blood lipid profile, liver function, musculoskeletal fitness, and health-related quality of life in overweight/obese older adult patients with knee osteoarthritis and type 2 diabetes: a randomized controlled trial during the COVID-19 pandemic. BMC Sports Sci Med Rehabil. 2024;16(1):125. [doi:10.1186/s13102-024-00915-4] [PMid:38831437]
- 52.Al-Mhanna SB, Leão C, Wan Ghazali WS, Mohamed M, Batrakoulis A, Afolabi HA, et al. Impact of Exercise on High-Density Lipoprotein Cholesterol in Adults with Overweight and Obesity: A Narrative Review. Ann Appl Sport Sci. 2024;12(1). [doi:10.61186/aassjournal.1300]
- 53.Al-Mhanna SB, Wan Ghazali WS, Batrakoulis A, Alkhamees NH, Drenowatz C, Mohamed M, et al. Impact of Various Types of Exercise on Lipid Metabolism in Patients with Type 2 Diabetes and Concurrent Overweight/Obesity: A Narrative Review. Ann Appl Sport Sci. 2024;12(1). [doi:10.61186/aassjournal.1324]
- 54.García-Hermoso A, Alonso-Martinez AM, Ramírez-Vélez R, Izquierdo M. Effects of exercise intervention on health-related physical fitness and blood pressure in preschool children: A systematic review and meta-analysis of randomized controlled trials. Sports medicine. 2020;50:187-203. [doi:10.1007/s40279-019-01191-w]
- 55.Batrakoulis A, Jamurtas AZ, Draganidis D, Georgakouli K, Tsimeas P, Poulios A, et al. Hybrid Neuromuscular Training Improves Cardiometabolic Health and Alters Redox Status in Inactive Overweight and Obese Women: A Randomized Controlled Trial. Antioxidants (Basel). 2021;10(10). [doi:10.3390/antiox10101601] [PMid:34679738]
- 57.Batrakoulis A, Fatouros IG. Psychological Adaptations to High-Intensity Interval Training in Overweight and Obese Adults: A Topical Review. Sports (Basel). 2022;10(5). [doi:10.3390/sports10050064] [PMid:35622474]
- 58. Yang E, Sharma G, Ram V. American and European hypertension guidelines: finding common ground. American Journal of Cardiology. 2019;123(8):1378-82. [doi:10.1016/j.amjcard.2019.01.009] [PMid:30736966]
- 59. Franklin BA, Thompson PD, Al-Zaiti SS, Albert CM, Hivert M-F, Levine BD, et al. Exercise-related acute cardiovascular events and potential deleterious adaptations following long-term exercise training: placing the risks into perspective—an update: a scientific statement from the American Heart Association. Circulation. 2020;141(13):e705-e36. [doi:10.1161/CIR.00000000000000749] [PMid:32100573]
- 60.Sakamoto S. Prescription of exercise training for hypertensives. Hypertension Research. 2020;43(3):155-61. [doi:10.1038/s41440-019-0344-1] [PMid:31656311]
- 61.Batrakoulis A, Jamurtas AZ, Fatouros IG. Exercise and type II diabetes mellitus: A brief guide for exercise professionals. Strength Cond J. 2022;44(6):64-72. [doi:10.1519/SSC.0000000000000731]