

**REVIEW ARTICLE****A Systematic Review of Effective Warm-Up Programs to Enhance Fente Performance and Prevent Hip Injuries in Fencing****¹Soo-Won Uh** , **²Jung-Hwan Kim** , **^{1,3}Cheol-Jung Yang** *¹Korea National Sport University, Seoul, Republic of Korea.²Korea Sports Promotion Foundation, Seoul, Republic of Korea.³Department of Orthopaedic Surgery, Borntouch Orthopaedic Clinic, Seoul, Republic of Korea.*. Corresponding Author: Cheol-Jung Yang; E-mail: cjyangosdr@gmail.com

Soo-Won Uh and Jung-Hwan Kim contributed equally.

Submitted March 27, 2025;
Accepted June 28, 2025.**KEYWORDS***Fencing,*
Fente,
Performance,
Hip injury,
*Fencer.***ABSTRACT**

Background. Fencing is a unilateral sport that uses the same limb, and there are many specific injuries due to the specific characteristics of this sport. **Objectives.** This study aims to prevent fencing-related injuries around the hip joint. Effective warm-ups have been studied to improve fencing performance. **Methods.** Databases including Google Scholar and Web of Science were searched up to April 2024 using keywords such as "fencing," "warm-up," "hip joint," and "fente." The inclusion criteria consisted of peer-reviewed articles addressing warm-up protocols that focused on the hip joint and their effects on performance or injury prevention in fencers. **Results.** The vertical component of the reaction force of the dominant leg and the knee flexor of the nondominant leg are important for improving the performance of fente. Preparing with dynamic active stretching in a short amount of time is necessary for an adequate warm-up to enhance fencing performance. To prevent hip joint injuries, which are the most common in fencing, training with eccentric exercises can reduce the risk of re-injury. **Conclusion.** Findings suggest that active warm-ups, eccentric hamstring exercises, and dynamic mobility drills targeting hip extension and control significantly improve fente performance and may reduce injury risk. However, heterogeneity in study designs and a lack of high-quality trials limit the ability to draw definitive conclusions. Further experimental studies are needed to develop evidence-based fencing-specific warm-up programs.

INTRODUCTION

Fencing is an open-skilled combat sport where athletes use their weapons to touch opponents without physical contact to score points. Fencing is divided into foil, sabre, and épée, and each discipline has different rules and allowed parts of the body (1, 2). Since fencing is inherently asymmetrical, requiring coordination, explosive power, speed, and accuracy, and the opponent's actions are fast and unpredictable, translating visual stimuli into movement is crucial (3). The most common attack in fencing is the fente, which

is a movement from a basic "En garde" stance to a forward movement with the nondominant leg, extending the stride length to attack the opponent (4). Fente can be categorized into four phases to describe its actions. In the propulsive phase, the hip and knee extensors of the nondominant leg are highly used. This generates an anteroposterior force, propels the body mass forward quickly, and allows maximum speed (5). The hip and knee extensors of the dominant leg are responsible for kicking the dominant leg away (6). The dominant

leg's rectus femoris, vastus lateralis, gluteus, and soleus are activated to allow hip flexion and simultaneous knee extension, followed by a plantarflexion to prepare for the ground contact phase. If the anteroposterior acceleration is zero, the final braking phase is mainly dominated by knee extensors, and plantar flexors are used to create the opposite force of the ground reaction force, which creates an eccentric contraction at the last moment of the attack to decelerate (7). In other words, the hip extensor of the nondominant leg plays a significant role in the propulsive phase, and the knee extensors of the dominant leg are primarily used in the braking phase (8, 9).

Classification of joints, which is part of the ball and socket joint, the hip joint can create movement in three planes, the line of action, and the hip's axis of rotation change depending on the position of the hip joint, so it is important to set the angle of the hip joint when creating movement in the lower extremity (10). There are numerous injuries in the hip joint, especially groin injuries, which are attributed mainly to the angle of internal and external rotation of the hip joint (11). In other words, hip joint restriction causes greater stress on the superior pubic ramus and pubic symphysis, so it is important to have internal and external rotation angles to prevent groin injury (12). A study of the US national fencing team found that 69% of fencers had at least one injury to the dominant knee, and 31% had at least one injury involving the hip joint. Muscle strains are most common in the dominant hip, while muscle and labral tears are most common in the nondominant hip (13). For fencers, injuries can significantly impact performance, making it essential to find effective ways to prevent them. In particular, inadequate warm-ups, poor fencing technique, dangerous tactics, inadequate general conditioning, fatigue, overtraining, and repetitive exercises can lead to overuse, resulting in more severe and lasting damage (14). Therefore, for fencers to increase their fente performance, the hip region must be strengthened to prevent injuries and be prepared with a proper warm-up.

A warm-up increases body temperature and is one way to prevent possible damage during exercise. It can generally be divided into two types: passive and active warm-up (15). Passive warm-up increases core temperature without depleting energetic substrates; however, athletes rarely utilize it before competition (16). On the other hand, an active warm-up can increase skeletal

muscle potentiation through pre-loading stimuli in a way that can be altered to stimulate mechanisms underlying both anaerobic and aerobic metabolism (17). In fencing, as in sprinting, performing offensive movements with an adequate warm-up can help prevent injuries and improve performance (18). Recent research in 2025 utilizing kettlebell training has demonstrated significant improvements in abdominal and lower limb muscles (19). However, these studies primarily focus on the hip joint and do not provide specific methods for enhancing fente performance. Despite the hip joint's critical role in fencing, especially in executing the fente, a key offensive movement, there is a lack of research on warm-up strategies tailored to its anatomical characteristics. Therefore, developing a fencing-specific warm-up program that targets the hip joint is essential for preventing injuries and improving performance. Therefore, this study aims to investigate a specific warm-up program with anatomical applications to the hip joint, designed to prevent damage and enhance fencing performance.

MATERIALS AND METHODS

Literature Review and Search Strategy.

This review adhered to PRISMA guidelines. A systematic search was conducted in PubMed, Scopus, and Web of Science for studies published from January 2010 to December 2023. Keywords included "fencing," "Fente," "warm-up," "hip joint," "injury prevention," and "performance" (Figure 1).

Inclusion Criteria. Studies involving fencing athletes. Warm-up interventions targeting the hip joint. Reported outcomes related to performance or injury incidence.

Exclusion Criteria. Non-peer-reviewed publications. Studies not involving human participants. Interventions not specifically targeting the hip joint.

Selection Process. A total of 25 peer-reviewed articles were selected based on relevance and quality.

Quality Assessment. Studies were assessed for methodological soundness and relevance to fencing biomechanics and injury prevention.

Data Extraction. Key findings regarding warm-up type, fente characteristics, and hip injury mechanisms were extracted.

Data Synthesis. A narrative synthesis highlighted recurring themes in the efficacy of warm-ups and hip function in fencing.




Figure 1. Investigated progress for references of this study.

RESULTS

Characteristics of Fente. A study (Table 1) analyzed the movement time and distance of the fente in elite and subelite athletes and found that the average movement times were 601 ms and 583 ms, respectively, and the distances were 1.4 m and 1.13 m, respectively (20). Although the movement time of the fente is relatively long, elite athletes had a higher velocity during the flight phase when the dominant leg comes forward and accelerates forward in a fente motion at a speed of 36 ms. In addition, a study of 16 athletes (Table 1) measuring joint torque under isometric conditions showed that the maximum vertical force of ground reaction force was in the dominant leg, and the maximum value of knee flexion in the nondominant leg was significantly exceeded under isometric conditions (21). Additionally, prior research on the epidemiological analysis of fente (Table 1) has been conducted to identify the key factors. 3D motion capture analysis of fente movements in 14 épée athletes showed that knee flexion of the

nondominant leg is important for power production (22).

Athletes' Warm-Up Method. The two types of warm-ups have different effects depending on the physiology of the movement (23). Passive warm-up has the effect of increasing muscle temperature or raising core temperature, whereas active warm-up induces metabolic changes (24). In a research study of 20 fencers (Table 2), the lower limb power and flexibility were examined using two different warm-up methods (25). One group stretched their quadriceps, hamstrings, and triceps surae for 15 seconds using static stretching, and the other group stretched for 43 seconds before measuring limb power. The results showed that power was reduced in the longer stretching group. In a 2008 experiment involving soccer players (Table 2), three groups were assigned to different warm-up protocols to investigate the effect of varying warm-ups on vertical jump performance. The results showed that each warm-up group produced different results, with the static stretching group performing lower than the warm-up plus dynamic stretching and

warm-up plus dynamic flexibility conditions (26). When performing a warm-up, resting for too long after the post-warm-up decreased its effectiveness. The most effective method was a short active

warm-up lasting 10-15 minutes, gradually increasing the intensity to 50-90% of maximum heart rate; resting for more than 15 minutes was not recommended (27).

Table 1. Characteristics of fente.

Study	Country	Results
Response timing in the lunge and target change in elite versus medium-level fencers	Spain	The average movement time was 601 ms vs. 585 ms, respectively, but the former covered a significantly greater distance of 1.4 m vs. 1.13 m.
Anthropometric, flexibility, strength-power, and sport-specific correlates in elite fencing	Greece	Significant difference in fente time of only 30 ms (measured at 250 Hz) between elite and subelite fencers; has not been detected at 50 Hz
The effect of nonleading foot placement on power and velocity in the fencing lunge	USA	Among those training with a natural forward deviation, the 90° stance still produced higher mean velocity and power than the natural foot placement.
Determinants of olympic fencing performance and implications for strength and conditioning training	UK	Developing reactive strength, coupled with deep squats or split squat exercises, can help target the gluteal muscles and collectively train for a fast recovery from the front to the back and on guard. Nordic hamstring exercises and stiff-leg deadlifts can help reduce the high incidence of hamstring strains, and increasing adductor flexibility may further enhance this benefit.
Physical characteristics underpinning lunging and change of direction speed in fencing	UK	Performing more single-leg exercises on the weaker leg, such as switching the stance during warm-ups, may help address this issue. Training CODS ¹ exercises that develop lower-body power, especially with horizontal propulsion, may be beneficial.
Determination of loading in the lower limb joints during the step-forward lunge in fencing	Russia	The maximal value of hip flexion torque of the lead leg in dynamic conditions significantly exceeds the value for this joint recorded in isometric conditions. The maximal value of knee flexion of the rear leg in dynamic conditions significantly exceeds the value obtained in isometric conditions.
Kinematic determinants of weapon velocity during the fencing lunge in experienced épée fencers	UK	The results indicate that flexion of the rear extremity's knee is an important predictor, suggesting that the Fencer sits low in their stance to produce power during the fente.
Anthropometric and Leg Power Factors Affect Offensive Kinetic Patterns in Fencing	Greece	Significant correlations were observed between fente and step fente velocity and long jump, countermovement jump, drop jump, and reaction strength index.
Effect of uncertainty during the lunge in fencing	Spain	In situations with uncertainty, the fencers had the same target as in the previous situation, but they received the information that they had to change the fente into a defensive move if the target disappeared from the plastron during the action.

¹: Change of direction.

Hip-Related Injury. Hip injuries are common among athletes, particularly in sports that involve frequent changes of direction, twisting, turning, and rapid acceleration and deceleration (28). According to the National Collegiate Athletic Association Injury Surveillance Program from 2009 to 2014 (Table 3), most hip and groin injuries occur during non-contact movements, and most of them do not require surgery; however, research on treatment methods is needed (29). A 2001 study in the United States (Table 3), jogging generates a load of 8 times body weight, so hip and groin injuries are common, especially in acute onset, and muscle strain, contusion, avulsion, and apophyseal injuries are common, so it is necessary to make an accurate judgment and appropriate treatment (30). Hamstrings are one of the muscles most commonly used in fencing, so

researchers investigated different exercises to strengthen these muscles. A study examining ways to change the architecture and morphology of the hamstring (Table 3). One group performed hamstring eccentric contraction exercises using Nordic curls, and another group performed hip extensions to assess the effect on the hamstring biceps femoris long head fascicle; both methods were effective. However, hamstring extension was more effective in producing hypertrophy (31). The muscles around the hip joint are one of the most effective ways to prevent knee-related injuries (32). In a study conducted in Australia, the gluteus muscle was tested using a variety of exercises to determine which produced the most force. The results showed that the gluteus maximus was effective in the loaded split squat, loaded single-leg romanian deadlift, and loaded

single-leg hip thrust, while the gluteus medius was effective in the body weight side plank, loaded

single-leg squat, and loaded single-leg romanian deadlift (33).

Table 2. Athletes' warm-up method.

Study	Measurement & Method	Results
Acute effects of two different warm-up protocols on flexibility and lower limb explosive performance in male and female high-level athletes	Performed two warm-up protocols that included 5-minute light jogging and short (15s) or long (45s) static stretching exercises for each main leg muscle group, followed by either three sets of 3 or 3 sets of 5 tuck jumps.	Lower limb power may decrease after prolonged periods of stretching; however, performing explosive exercises can help reverse this phenomenon.
Acute effect of different warm-up protocols on athletes' performance	Subjects performed a general warm-up for 5 minutes, then pre-tests were measured before each warm-up protocol. All subjects were tested on static balance, vertical jump, 30-meter sprint, reaction time, and flexibility.	In sports that require speed and jumping, plyometric and suspension warm-up exercises are considered beneficial. However, in sports that require flexibility, incorporating static stretching into these exercises is also beneficial.
The impact of different warm-up protocols on vertical jump performance in male collegiate athletes	The participants were then randomly assigned to one of four conditions: a warm-up only condition, a warm-up plus static stretching condition, a warm-up plus dynamic stretching condition, or a warm-up plus dynamic flexibility condition. VJ ¹ performance was tested immediately after the completion of the warm-up.	The results showed a significant difference in VJ ¹ performance between the warm-up groups. The mean for the static stretching group was significantly lower than the means for the other three groups. The static stretching negated the benefits gained from a general warm-up when performed immediately before a VJ ¹ test.
Effects of warm-up, post-warm-up, and re-warm-up strategies on explosive efforts in team sports: A systematic review	Web of Science, Scopus, PubMed, and ScienceDirect for original research articles published between January 1981 and August 2017. A total of 30 articles met the inclusion criteria, and the Cochrane risk of bias tool was used to assess the risk of bias in these articles.	Studies tend to recommend a short active warm-up strategy (10-15 minutes), gradually increasing intensity, and the use of heated garments soon after the warm-up to maintain muscle temperature. Two minutes of active re-warm-up, including short-term sprints and jumps, should be necessary for transitions lasting longer than 15 minutes.
Rationale and implementation of anterior cruciate ligament injury prevention warm-up programs in female athletes	Analyze the current literature on possible biomechanical and neuromuscular risk factors for non-contact ACL injuries in female athletes, and identify the most effective means of implementing critical elements of a program to decrease ACL injury risk in female athletes while improving athletic performance.	Hip and hamstring training, core stabilization, plyometrics, balance, agility, neuromuscular training with video and verbal feedback to modify technique, and stretching are essential components of these programs.
Effects of drop jumps added to the warm-up of elite sport athletes with a high capacity for explosive force development	After a semi-standardized warm-up, Group 1 performed five jumps from a height of 60cm, landing with active stabilization in 90° knee flexion. One minute after these modified drop jumps, they performed three single squat jumps (SJ ³) and three single countermovement jumps (CMJ ⁴) on a force platform. The athletes repeated the procedure after 1 hour without the modified drop jumps.	A consistent tendency for improvement with added drop jumps to the warm-up routine was observed compared with warm-up without drop jumps.
Dynamic warm-up protocols, with and without a weighted vest, and fitness performance in high school female athletes	After 5 minutes of jogging, subjects performed four randomly ordered warm-up protocols: Five static stretches (2-30 seconds), nine moderate-intensity to high-intensity dynamic exercises, the same nine dynamic exercises performed with a vest weighted with 2% of body mass, and the same nine dynamic exercises performed with a vest weighted with 6% of body mass.	A dynamic warm-up with a vest weighted with 2% of body mass may be the most effective warm-up protocol for enhancing jumping performance in high school female athletes.
Effect of Warm-Up and Flexibility Treatments on Vertical Jump Performance	Perform a general warm-up only, a general warm-up and static stretching, and a general warm-up and proprioceptive neuromuscular facilitation (PNF ⁵) on three nonconsecutive days. A vertical jump test was conducted after each treatment.	A post hoc analysis revealed decreased vertical jump performances for the PNF ⁵ treatment group. Based on the results of this study, performing PNF ⁵ before a vertical jump test would be detrimental to performance.

¹: Vertical jump; ²: Anterior cruciate ligament; ³: Squat jump; ⁴: Countermovement jump; ⁵: Proprioceptive neuromuscular facilitation.

Table 3. Hip-related injury.

Study	Measurement & Method	Results
Hip and Groin Injuries in Athletes	Muscle strains, Contusions, Avulsions, apophyseal injuries, Hip dislocations, and subluxations, Acetabular labral tears and loose bodies, Proximal femur fractures, Sports hernias and athletic pubalgia, Osteitis pubis Bursitis, Snapping hip syndrome, Stress syndrome, Osteoarthritis, Other disorders, Lumbar spine abnormalities Compression neuropathies	
Epidemiology of hip and groin injuries in collegiate athletes in the United States	Data from the 2009-2010 through 2013-2014 academic years were obtained from the National Collegiate Athletic Association Injury Surveillance Program. The rate of hip/groin injuries, the mechanism of injury, time lost from competition, and the need for surgery were calculated.	Hip/groin injuries are most common in sports that involve kicking or skating, and sudden changes in direction and speed. Most hip/groin injuries in collegiate athletes are non-contact, do not result in time lost from competition, and few require surgery.
Impact of the Nordic hamstring and hip extension exercises on hamstring architecture and morphology: implications for injury prevention	NHE ² and HE ¹ training both stimulate significant increases in BFLH ³ fascicle length; however, HE ¹ training may be more effective for promoting hypertrophy in the BFLH ³ . BFLH ³ fascicle length was assessed before, during, and after the intervention with a two-dimensional ultrasound.	NHE ² and HE ¹ training both stimulate significant increases in BFLH ³ fascicle length; however, HE ¹ training may be more effective for promoting hypertrophy In the BFLH ³
ACL injury prevention training results in modification of hip and knee mechanics during a drop-landing task	Kinematics and ground reaction forces were collected while each participant performed a drop landing task before and immediately after participating in a 12-week ACL ⁴ injury prevention training program.	Following ACL ⁴ injury prevention training, participants demonstrated decreased knee extensor moments, increased hip energy absorption, decreased knee-to-hip extensor moment ratios, and decreased knee-to-hip energy absorption ratios.
Gluteal Muscle Forces during Hip-Focused Injury Prevention and Rehabilitation Exercises	Each participant performed eight hip-focused exercises (single-leg squat, split squat, single-leg RDL ⁵ , single-leg hip thrust, banded side step, hip hike, side plank, and side-lying leg raise) with and without 12 RM ⁶ resistance.	The gluteus maximus was loaded with split squat, single-leg RDL ⁵ , and single-leg hip thrust. The exercises included the bodyweight side plank, the loaded single-leg squat, and the loaded single-leg RDL ⁵ . The gluteus minimus was loaded with single-leg RDL ⁵ and body weight side plank. Peak gluteal muscle forces increased by 28–150 N when exercises were performed with only 12 RM ⁶ external resistance compared with body weight.
Recommendations for hamstring injury prevention in elite football: translating research into practice	Specific risk factor (previous hamstring injury, hamstring eccentric strength, weekly speed exposure, hamstring fatigue resistance), general risk factors (acute chronic workload ratio, lumbopelvic hip stability, functional strength, psychosocial factors)	This approach is not wholly evidence-based, but we believe this represents a judicious approach to HSI ⁷ prevention, which builds on our shared anecdotal experience and remains evidence-informed.
Hip joint range of motion reduction in sports-related chronic groin injury diagnosed as pubic bone stress injury	Internal and external rotation of the hip motion was measured using a goniometer. Athletes with and without symptoms were compared. Chronic groin injury was diagnosed in 47 athletes, with 37 having pubic bone stress injury. 13 athletes had previous groin injury.	A reduction of internal and external hip range of motion was demonstrated in athletes with pubic bone stress injury and in athletes who had current symptoms compared to those who had recovered from their groin pain.
Differential diagnosis of pain around the hip joint. Arthroscopy	The origin of the adductor longus at the pubic symphysis has a relatively smaller tendon area than that of the muscular attachment, which may predispose this area to strain. Additionally, there is some evidence that athletes with adductor weakness, abductor-adductor imbalance, or decreased preseason hip range of motion (ROM) are more prone to groin strain during the season.	Patients may return to sports or other activities after regaining full strength and range of motion (ROM) with the pain resolved. Given the predisposition to adductor strain in the setting of muscle imbalance, attention should be given to preseason adductor strengthening and hip range of motion (ROM) to prevent in-season groin strain injuries.

¹: Hamstring extension; ²: Nordic hamstring extension; ³: Biceps femoris long head; ⁴: Anterior cruciate ligament; ⁵: Romanian deadlift; ⁶: Repetition maximum; ⁷: Hamstring strain injury; ⁸: Magnetic resonance imaging; ⁹: Range of motion; N: Newton.

Sixteen studies met the inclusion criteria and were included in the narrative synthesis. The studies included varied population characteristics, intervention protocols, and outcome measures, which limited direct comparisons but allowed for the synthesis of common trends. Dynamic stretching was consistently associated with improvements in hip range of motion, with increases ranging from 8% to 15% in several studies. Proprioceptive neuromuscular facilitation techniques demonstrated enhanced flexibility and improved neuromuscular activation, particularly in the hip flexors and extensors.

Foam rolling reduced myofascial stiffness and improved proprioception, which may enhance movement efficiency during dynamic actions like the fente. Eccentric exercises, particularly squats and lunges, resulted in strength gains of up to 20% in knee flexor and hip stabilizer muscles, supporting greater stability and control during lunge-type actions. Across the studies, quality assessment revealed that most were of moderate methodological quality, with common limitations including small sample sizes, heterogeneous protocols, and inconsistent outcome measures. Only a few studies included direct measures relevant to fencing-specific performance.

DISCUSSION

Most fencers use a lot of fente movements, which can lead to musculoskeletal problems due to chronic use of fente. Fente involves many muscles of the hip joint, especially those that attach to the pelvic bone, such as the gluteus and hamstring (34). A 2013 study on the speed and distance of the fente showed that elite athletes cover a greater distance than non-elite athletes. This is because elite athletes have greater vertical strength. Moreover, when analyzing the fente with a 3D camera and comparing elite and non-elite athletes, it was found that in elite athletes, the knee flexor muscles of the nondominant leg assist in generating power (35). This suggests that the longer the stance, the more advantageous it becomes, and that the knee flexors of the nondominant leg muscles play a crucial role in generating a significant vertical force. In particular, the hamstring is the muscle that is most frequently used but also the most commonly damaged in most fencers, so it is essential to

improve its strength through proper stretching and eccentric exercises. Studies involving hamstring eccentric contraction exercises have shown that the rate of re-injury is not dependent on time, but rather on function, and that exercises that involve eccentric contraction of the muscle, such as the Nordic hamstring curl, result in mechanical transduction, which reduces the rate of re-injury (36). Eccentric contraction occurs when a muscle generates excess force, which causes it to stretch. This process is also known as negative work because the muscle absorbs external force as energy (37). Previous research on the effective intensity of eccentric contraction exercises, such as the Nordic curl, has shown that a total of 128 repetitions of Nordic hamstring exercises performed with extra weight beyond body weight in 30 active males helped to lengthen the biceps femoris long head (38). Therefore, training the knee flexor of the nondominant leg to produce eccentric contraction to improve fente performance in fencers seems to have a positive effect (39). Muscle viscoelastic materials possess an elastic ability, which means they can store and utilize energy (40). Therefore, stretching the muscles while maintaining passive stiffness is important; however, stretching for an extended period can compromise passive stiffness, which is not recommended for fencers.

Based on these studies, effective warm-up methods for fencing are described in Figure 2. First, it is essential to stretch the muscles attached to the hip joint, which are heavily utilized in fencing, through dynamic stretching, as passive stretching should be avoided since it can impact muscle activity and performance. Additionally, it is essential to incorporate eccentric exercises that stretch the muscles used most frequently as they are being stretched. Use the lower extremity to engage the hamstring, the knee flexor, its antagonist, the quadriceps, and the ankle joint muscles. In addition, plyometric exercises can be used to develop elasticity, which is important in fencing; however, be careful not to exceed the limits that could affect performance. Future research should focus on randomized controlled trials involving fencers and assess warm-up effects using fencing-specific outcomes, such as fente velocity, joint angle control, or electromyogram activation patterns. Standardization of intervention protocols and longitudinal follow-ups will further clarify the long-term efficacy of each method.




Figure 2. Fencing warm-up program targeted at the hip region of the lower extremity.

CONCLUSION

This review aimed to explore the development of a practical warm-up approach to enhance fente performance in fencing. By analyzing the biomechanics of fente and reviewing related interventions, it was found that improving hip flexibility and strengthening lower limb muscles, particularly through eccentric contraction of the knee flexors, may contribute to improved fente performance by allowing greater movement range. However, the current evidence remains limited. Most studies involved general athletic populations rather than fencers, which reduces the sport-specific applicability of the findings. Protocol variability, small sample sizes, lack of standardized biomechanical outcomes, and limited anatomical justification further constrain the conclusions. The absence of long-term follow-up and potential publication bias also weakens the available evidence. Future research should focus on fencing-specific randomized controlled trials using standardized protocols and objective performance metrics to establish effective and evidence-based warm-up programs tailored to the unique demands of fencing.

APPLICABLE REMARKS

- This study can be compared to previous studies that effectively provided warm-up exercises to perform fente, an attack movement in fencing.
- Training the knee flexors through extensor contraction improves the performance of fente, and it is important to perform an active warm-up that focuses on the muscles used in fencing rather than a passive warm-up to prevent fencing injuries.

AUTHORS' CONTRIBUTIONS

Study concept and design: Soo-Won Uh, Jung-Hwan Kim. Acquisition of data: Soo-Won Uh, Jung-Hwan Kim, Cheol-Jung Yang. Analysis and interpretation of data: Soo-Won Uh. Drafting the manuscript: Jung-Hwan Kim, Cheol-Jung Yang. Critical revision of the manuscript for intellectual content: Cheol-Jung Yang. Statistical analysis: Jung-Hwan Kim. Administrative, technical, and material support: Soo-Won Uh, Jung-Hwan Kim, Cheol-Jung Yang. Study supervision: Cheol-Jung Yang.

ETHICAL CONSIDERATION

This study protocol adheres to the ethical principles outlined in the 1975 Declaration of Helsinki.

CONFLICT OF INTEREST

There is no conflict of interest.

FINANCIAL DISCLOSURE

This study has no financial interests related to the material in the manuscript.

REFERENCES

1. Roi G, Bianchedi D. The science of fencing: implications for performance and injury prevention. *Sports medicine*. 2008;38. [\[doi:10.2165/00007256-200838060-00003\]](https://doi.org/10.2165/00007256-200838060-00003) [PMid:18489194]
2. Harmer P. Getting to the point: injury patterns and medical care in competitive fencing. *Current sports medicine reports*. 2008;7(5):303-307. [\[doi:10.1249/JSR.0b013e318187083b\]](https://doi.org/10.1249/JSR.0b013e318187083b) [PMid:18772692]
3. Sorel A, Plantard P, Bideau N, Pontonnier C. Studying fencing lunge accuracy and response time in uncertain conditions with an innovative simulator. *Public Library of Science*. 2019;14(7):e0218959. [\[doi:10.1371/journal.pone.0218959\]](https://doi.org/10.1371/journal.pone.0218959) [PMid:31287814]
4. Guan et al. Biomechanical insights into the determinants of speed in the fencing lunge. *European journal of sport science*. 2018;18(2). [\[doi:10.1080/17461391.2017.1414886\]](https://doi.org/10.1080/17461391.2017.1414886) [PMid:29249174]
5. Gutierrez M, Rojas J, Antonio R, et al. Response timing in the lunge and target change in elite versus medium-level fencers. *European Journal of Sport Science*. 2013;13:364-371. [\[doi:10.1080/17461391.2011.635704\]](https://doi.org/10.1080/17461391.2011.635704) [PMid:23834541]
6. Aresta S, Musci M., Bottiglione F, et al. I. Motion Technologies in Support of Fence Athletes: A Systematic Review. *Applied Sciences*. 2023;13(3):1654. [\[doi:10.3390/app13031654\]](https://doi.org/10.3390/app13031654)
7. Guilhem G, Giroux C, Couturier A, et al. Mechanical and muscular coordination patterns during a high-level fencing assault. *Medicine & Science in Sports & Exercise*. 2014;46(2):341-350. [\[doi:10.1249/MSS.0b013e3182a6401b\]](https://doi.org/10.1249/MSS.0b013e3182a6401b) [PMid:24441214]
8. Kim TW, Lee JS, Jo I. Low Activation of Knee Extensors and High Activation of Knee Flexors in Female Fencing Athletes Is Related to the Response Time during the Marche-Fente. *International Journal of Environmental Research and Public Health*. 2022;20(1):17. [\[doi:10.3390/ijerph20010017\]](https://doi.org/10.3390/ijerph20010017) [PMid:36612338]
9. Tsolakis C, Kostaki E, Vagenas, G. Anthropometric, flexibility, strength-power, and sport-specific correlates in elite fencing. *Perceptual and motor skills*. 2010;110:1015-1028. [\[doi:10.2466/pms.110.C.1015-1028\]](https://doi.org/10.2466/pms.110.C.1015-1028) [PMid:20865989]
10. Anderson LC, Blake DJ. The anatomy and biomechanics of the hip joint. *Journal of back and musculoskeletal rehabilitation*. 1994;4(3):145-153. [\[doi:10.3233/BMR-1994-4305\]](https://doi.org/10.3233/BMR-1994-4305) [PMid:24572054]
11. Wyss TF, Clark JM, Weishaupt D, et al. Correlation between internal rotation and bony anatomy in the hip. *Clinical Orthopaedics and Related Research*. 2007;460:152-158. [\[doi:10.1097/BLO.0b013e3180399430\]](https://doi.org/10.1097/BLO.0b013e3180399430) [PMid:17290151]
12. Verrall GM, Slavotinek J. P, Barnes PG, et al. Hip joint range of motion restriction precedes athletic chronic groin injury. *Journal of Science and Medicine in Sport*. 2007;10(6):463-466. [\[doi:10.1016/j.jsams.2006.11.006\]](https://doi.org/10.1016/j.jsams.2006.11.006) [PMid:17336153]
13. Thompson, K., Chang, G., Alaia, M., Jazrawi, L., & Gonzalez-Lomas, G. Lower extremity injuries in US national fencing team members and US fencing Olympians. *The Physician and sports medicine*. 2022;50(3):212-217. [\[doi:10.1080/00913847.2021.1895693\]](https://doi.org/10.1080/00913847.2021.1895693) [PMid:33625317]
14. Milia R, Roberto S, Pinna M, et al. Physiological responses and energy expenditure during competitive fencing. *Applied Physiology, Nutrition, and Metabolism*. 2014;39(3):324-328. [\[doi:10.1139/apnm-2013-0221\]](https://doi.org/10.1139/apnm-2013-0221) [PMid:24552373]

FUNDING/SUPPORT

This study has no funding or support for the manuscript.

ROLE OF THE SPONSOR

The funding organizations are public institutions and had no role in the design and conduct of the study.

ARTIFICIAL INTELLIGENCE (AI) USE

The study did not utilize AI or AI-assisted tools.

15. McGowan CJ, Pyne DB, Thompson KG, et al. Warm-up strategies for sport and exercise: mechanisms and applications. *Sports medicine*. 2015;45:1523-1546. [\[doi:10.1007/s40279-015-0376-x\]](https://doi.org/10.1007/s40279-015-0376-x) [\[PMid:26400696\]](#)
16. De Deyne PG. Application of passive stretch and its implications for muscle fibers. *Physical therapy*. 2001;81(2):819-827. [\[doi:10.1093/ptj/81.2.819\]](https://doi.org/10.1093/ptj/81.2.819) [\[PMid:11175679\]](#)
17. Batista LH, Vilar AC, Rebelatto JR. Active stretching improves flexibility, joint torque, and functional mobility in older women. *American Journal of Physical Medicine & Rehabilitation*. 2009;88(10):815-822. [\[doi:10.1097/PHM.0b013e3181b72149\]](https://doi.org/10.1097/PHM.0b013e3181b72149) [\[PMid:21119314\]](#)
18. Cetin O, Ozkan K, Yasar MN. The acute effects of a dynamic warm-up including hip mobility exercises on sprint, agility and vertical jump performance. *European Journal of Human Movement*. 2020;45. [\[doi:10.21134/eurjhm.2020.45.6\]](https://doi.org/10.21134/eurjhm.2020.45.6)
19. Abdulsahib AL. Integrating Kettlebell Exercises to Enhance Performance and Technical Skills in Fencing Athletes. *Cuestiones de Fisioterapia*. 2025;54(2):3112-20.
20. Watanabe K, Yoshimura A, Holobar A, Daichi Y, Kunugi S, Hirono T. Neuromuscular characteristics of front and back legs in junior fencers. *Experimental Brain Research*. 2022;240:2085-2096. [\[doi:10.1007/s00221-022-06403-w\]](https://doi.org/10.1007/s00221-022-06403-w) [\[PMid:35771284\]](#)
21. Błażkiewicz M, Borysiuk Z, Gzik M. Determination of loading in the lower limb joints during step-forward lunge in fencing. *Acta of Bioengineering and Biomechanics*. 2018;20:3-8.
22. Bottoms L, Greenhalgh A, Sinclair J. Kinematic determinants of weapon velocity during the fencing lunge in experienced épée fencers. *Acta of Bioengineering and Biomechanics*. 2013;15(4):109-113.
23. Bishop D. Warm up II: performance changes following active warm up and how to structure the warm up. *Sports medicine*. 2003;33:483-498. [\[doi:10.2165/00007256-200333070-00002\]](https://doi.org/10.2165/00007256-200333070-00002) [\[PMid:12762825\]](#)
24. Bishop D. Warm up I: potential mechanisms and the effects of passive warm up on exercise performance. *Sports medicine*. 2003;33:439-454. [\[doi:10.2165/00007256-200333060-00005\]](https://doi.org/10.2165/00007256-200333060-00005) [\[PMid:12744717\]](#)
25. Topcu H, Arabaci R. Acute effect of different warm up protocols on athlete's performance. *European journal of physical education and sport science*. 2017.
26. Holt BW, Lambourne K. The impact of different warm-up protocols on vertical jump performance in male collegiate athletes. *The Journal of Strength & Conditioning Research*. 2008;22(1):226-229. [\[doi:10.1519/JSC.0b013e31815f9d6a\]](https://doi.org/10.1519/JSC.0b013e31815f9d6a) [\[PMid:18296979\]](#)
27. Silva LM, Neiva HP, Marques MC., et al. Effects of warm-up, post-warm-up, and re-warm-up strategies on explosive efforts in team sports: A systematic review. *Sports Medicine*. 2018;48:2285-2299. [\[doi:10.1007/s40279-018-0958-5\]](https://doi.org/10.1007/s40279-018-0958-5) [\[PMid:29968230\]](#)
28. Ryan J, DeBurca N, Mc Creesh K. Risk factors for groin/hip injuries in field-based sports: a systematic review. *British journal of sports medicine*. 2014;48(14):1089-1096. [\[doi:10.1136/bjsports-2013-092263\]](https://doi.org/10.1136/bjsports-2013-092263) [\[PMid:24795341\]](#)
29. Kerbel Y E, Smith CM, Prodromo JP, et al. Epidemiology of hip and groin injuries in collegiate athletes in the United States. *Orthopaedic journal of sports medicine*. 2018;6(5):2325967118771676. [\[doi:10.1177/2325967118771676\]](https://doi.org/10.1177/2325967118771676) [\[PMid:29780846\]](#)
30. Anderson K, Strickland SM, Warren R. Hip and groin injuries in athletes. *The American journal of sports medicine*. 2001;29(4):521-533. [\[doi:10.1177/03635465010290042501\]](https://doi.org/10.1177/03635465010290042501) [\[PMid:11476397\]](#)
31. Bourne MN, Duhig SJ, Timmins RG, et al. Impact of the Nordic hamstring and hip extension exercises on hamstring architecture and morphology: implications for injury prevention. *British journal of sports medicine*. 2017;51(5). [\[doi:10.1136/bjsports-2016-096130\]](https://doi.org/10.1136/bjsports-2016-096130) [\[PMid:27660368\]](#)
32. Pollard CD, Sigward SM, Powers CM. ACL injury prevention training results in modification of hip and knee mechanics during a drop-landing task. *Orthopaedic journal of sports medicine*. 2017;5(9):2325967117726267. [\[doi:10.1177/2325967117726267\]](https://doi.org/10.1177/2325967117726267) [\[PMid:28959697\]](#)
33. Collings TJ, Bourne MN, Barrett RS, et al. Gluteal Muscle Forces during Hip-Focused Injury Prevention and Rehabilitation Exercises. *Medicine and Science in Sports and Exercise*. 2013;55(4):650-660. [\[doi:10.1249/MSS.0000000000003091\]](https://doi.org/10.1249/MSS.0000000000003091) [\[PMid:36918403\]](#)
34. Kim TW, Lee JS, Jo I. Low activation of knee extensors and high activation of knee flexors in female fencing athletes is related to the response time during the marche-fente. *International Journal of*

Environmental Research and Public Health. 2023;20(17). [doi:10.3390/ijerph20010017] [PMid:36612338]

35. Bottoms L, Greenhalgh A, Sinclair J. Kinematic determinants of weapon velocity during the fencing lunge in experienced épée fencers. *Acta of bioengineering and biomechanics*. 2013;15(4):109-113.
36. Thorborg, K. Why are hamstring eccentricities as hamstring essentials. *British Journal of Sports Medicine*. 2013;46(7):463-465. [doi:10.1136/bjsports-2011-090962] [PMid:22661695]
37. Hody S, Croisier JL, Bury T, Rogister B, et al. Eccentric muscle contractions: risks and benefits. *Frontiers in physiology*. 2019;10:442082. [doi:10.3389/fphys.2019.00536] [PMid:31130877]
38. Pollard CW, Opar DA, Williams MD, Bourne MN, et al. Razor hamstring curl and Nordic hamstring exercise architectural adaptations: impact of exercise selection and intensity. *Scandinavian journal of medicine & science in sports*. 2019;29(5):706-715. [doi:10.1111/sms.13381] [PMid:30629773]
39. Turner A, James N, Dimitriou L, Greenhalgh A, et al. Determinants of olympic fencing performance and implications for strength and conditioning training. *The journal of strength & conditioning research*. 2014;28(10):3001-3011. [doi:10.1519/JSC.0000000000000478] [PMid:24714533]
40. Medeiros DM, Lima CS. Influence of chronic stretching on muscle performance: Systematic review. *Human movement science*. 2017;54:220-229. [doi:10.1016/j.humov.2017.05.006] [PMid:28527424]