

**ORIGINAL ARTICLE**

Effects of Elastic Band Resistance Training and Dumbbell Resistance Training on Shoulder Pain and Range of Motion in Korean Traditional Dancers with Adhesive Capsulitis: A Randomized Controlled Trial

¹Hee-Jin Kim , **²Hea Kyung Choi** *¹Department of Dance, College of Arts & Physical Education, Sejong University, Seoul, Republic of Korea.²Department of Dance and Convergence Art, Sungshin Women's University, Seoul, Republic of Korea.*. Corresponding Author: Hea Kyung Choi; E-mail: queenndance@sungshin.ac.krSubmitted April 16, 2025;
Accepted June 27, 2025.**KEYWORDS***Dancer,
Shoulder Pain,
Rehabilitation,
Adhesive Capsulitis.***ABSTRACT**

Background. Shoulder pain is prevalent not only in the general population but also in dancers, who engage their shoulders intensely. **Objectives.** This study aimed to identify practical rehabilitation exercises for Korean traditional dancers suffering from adhesive capsulitis (AC), focusing on the comparative effects of elastic bands and dumbbells. **Methods.** Sixteen female participants diagnosed with shoulder pain were divided into two groups and underwent a 6-week training program involving either elastic bands or dumbbells. Measurements of shoulder joint range of motion (ROM) and pain intensity using the Numerical Rating Scale (NRS) were collected pre- and post-intervention. **Results.** Both groups demonstrated improvements in ROM and NRS, with large effect sizes; however, differences between groups were not statistically significant. **Conclusion.** Strength training using both elastic bands and dumbbells has a positive influence on shoulder rehabilitation. This suggests that resistance training, regardless of the tool used, is beneficial for recovery. Further research with larger samples and longer durations is recommended.

INTRODUCTION

In Korea, the condition is referred to as adhesive capsulitis (AC), which occurs when inflammation develops in the shoulder joint capsule, causing the capsule to thicken, contract, and adhere to surrounding tissues. This condition is commonly associated with individuals in their 50s, characterized by a natural recovery process, which has led to the colloquial term "frozen shoulder" (1). AC affects approximately 2-5% of the population, and according to national health big data on diseases of public interest, about 62% of patients are women as of 2021. Furthermore, it

has been found that approximately 20% of patients experience similar symptoms in the opposite shoulder (2). AC generally occurs within five years following shoulder recovery (3). Although it was initially common in individuals over 50, recent trends indicate that it is increasingly affecting younger age groups, including those in their 30s and 40s, particularly dancers who use their shoulders intensively (4).

Appropriate treatment for AC patients is essential to prevent permanent shoulder dysfunction (5). Among various treatment

options, non-surgical treatments are preferred, including medication, manual therapy, and exercise-based interventions. Medication, such as nonsteroidal anti-inflammatory drugs and corticosteroids, is typically used early to reduce inflammation and pain (6). Additionally, interventional procedures such as hydraulic distension, which involves injecting saline and local anesthetics to expand the joint capsule and break adhesions, are commonly used. Hydraulic distension, introduced by Andren and Lundberg in 1965, remains a widely utilized treatment for AC patients (7).

Various tools are used in exercise programs to improve physical strength and balance (8). Resistance bands are one of the most popular exercise tools due to their portability, ease of use, and lack of space constraints, as well as their widespread use across age groups (9). Resistance bands are effective in rehabilitation and fitness improvement due to their low-impact nature (10). Their ability to adjust resistance in accordance with changes in joint movement direction during exercise allows for the activation of multiple muscles in a single movement (11).

Despite differences in the direction and magnitude of resistance, resistance bands and other resistance tools, such as dumbbells, are often used interchangeably in various exercise programs (12). Ribeiro et al. (13) reported that resistance band exercises can effectively replace heavy weights for strength training, and Andersen et al. (14) found that resistance bands and dumbbells elicit similar levels of muscle activation, suggesting that both resistance tools can be used equivalently during exercise. However, other studies have reported that resistance band training produces higher muscle activation compared to dumbbells (15). These differing findings in the literature highlight the challenges in comparing the effectiveness of resistance bands and dumbbells (14), which may lead to confusion among users.

Additionally, research on suitable resistance exercise tools for patients with AC is limited. Although flexibility exercises are important for patients with adhesive capsulitis, they may lead to greater damage if they are only aimed at improving the range of motion (16, 17). Isometric exercises using resistance bands can safely increase the reduced range of motion in patients with adhesive capsulitis. Isometric exercises that increase muscle strength to maintain muscle

length can reduce additional damage and safely increase the range of motion (18). However, research on appropriate resistance exercise tools is limited. Due to this lack of research, patients with adhesive capsulitis may not benefit significantly from exercise even if they start treatment at the right time and with the proper method. Therefore, this study aimed to compare the effects of resistance bands and dumbbell-based strength training on shoulder range of motion, pain, and functional disability indices in patients with adhesive capsulitis.

MATERIALS AND METHODS

Study Design. This randomized controlled trial involved 16 women who visited a medical institution and were diagnosed with adhesive capsulitis within the past year. All of these women were dancers majoring in Korean dance and complained of shoulder pain. The participants were randomly assigned to two groups: a resistance band group ($n = 8$) and a dumbbell group ($n = 8$). The intervention was performed three times a week for six weeks, with each session lasting 25 minutes and consisting of 12 repetitions per set (Figure 1).

Participants. Inclusion criteria were female participants experiencing shoulder pain rated seven or higher on the NRS. Individuals with cardiac conditions, visual or cognitive impairments, or those who had undergone shoulder surgery within the previous 6 months were excluded.

Tools and Reagents. Inclusion criteria included female participants with AC who experienced shoulder pain with a pain score of 7 or greater on the Nearly Rated Shoulder Pain Rating Scale (NRS). Those with cardiac disease, visual or cognitive impairment, or who had undergone shoulder surgery within the previous 6 months were excluded.

Training Protocol. Both groups performed the same exercises using resistance bands or dumbbells, and additional supervised home exercise was performed weekly. Each session included joint mobility exercises (25 minutes) using the lowest level of resistance bands and muscle strengthening exercises (25 minutes) using 2 kg dumbbells (Tables 1, 2).

Data Collection. To check the mobility of the shoulder joint, the subject stands with his back to the wall, performs shoulder flexion and horizontal abduction movements, and measures

the angle between the arm and the wall. The anatomical posture for horizontal abduction is 0 degrees, and the angle is measured by how much the arm is raised. NRS scores were recorded both before and after the intervention. SPSS 26.0 was used for statistical analysis, and

an independent sample t-test was conducted to compare the pre- and post-test data of each group. A two-way repeated measures analysis of variance was used to confirm the difference between groups in NRS, with a significance level set at $p < 0.05$.




Figure 1. Indicates the research design.

Table 1. Shoulder articular mobilization.

Articular Mobilization	Grade	Time
<u>Anterior gliding</u>		
<u>Inferior gliding</u>		
<u>Posterior gliding</u>	I~III	25 min / once a week
<u>Superior gliding</u>		

Table 2. Exercise program.

	Workout	Rep/Set
Warming up	Wall slide – Flexion	
	Wall slide – Horizontal Abduction	
	Wall slide – Horizontal Adduction	10 reps / 1 set
	Quadruped Shoulder Tap	
Main Workout	Scaption	
	Lateral Rotation	
	Medial Rotation	10 reps / three sets
	Horizontal Abduction	
	Horizontal Adduction	
Cool down	Thoracic Rotation	10 reps / 1 set

Rep: Repetition.

RESULTS

The average age of the participants was 54.7 ± 6.3 years for the elastic band group and 53.3 ± 5.8 years for the dumbbell group. The average height was 161.7 ± 5.7 cm for the elastic band group and 161.2 ± 5.7 cm for the dumbbell group.

The results of ROM measurements showed an increase in numerical values; however, no

statistically significant interaction effects between time and group were found for each movement. Specifically, for flexion ($F=1.00$, $p=0.300$), abduction ($F=0.977$, $p=0.788$), lateral rotation ($F=0.001$, $p=0.955$), and medial rotation ($F=0.187$, $p=0.598$), there were no significant interaction effects between time and group (Table 3, Figure 2).




Figure 2. Changes in shoulder ROM pre- and post-intervention. ROM: Range of motion.

Table 3. Analysis results of ROM [°].

Variable	Group	Pre (Mean±SD)	Post (Mean±SD)	Source	F	p
Flexion	Elastic band group	117 ± 15.4	166.4 ± 7.5	Time	180.1	$<0.001^{***}$
	Dumbbell group	122.7 ± 21.4	159.8 ± 11.2	Group	0.587	0.346
Abduction	Elastic band group	84.5 ± 15.2	140.1 ± 17.1	Time	202.01	$<0.001^{***}$
	Dumbbell group	95.3 ± 18.9	147.7 ± 24.1	Group	0.850	0.344
Lateral Rotation	Elastic band group	45.7 ± 11.2	77.8 ± 4.9	Time	155.4	$<0.001^{***}$
	Dumbbell group	47.4 ± 15.0	74.4 ± 10.7	Group	0.000	0.985
Medial Rotation	Elastic band group	50.5 ± 9.2	69.2 ± 9.9	Time	87.51	$<0.001^{***}$
	Dumbbell group	53.7 ± 14.9	77 ± 11.9	Group	1.52	0.215
				Time × Group	0.001	0.955
					0.187	0.598

ROM: Range of motion; Values are expressed as Mean \pm SD; *: $p<0.05$; **: $p<0.01$; ***: $p<0.001$.

For the NRS in the two-way repeated measures ANOVA, although no significant differences between groups or significant interaction effects between time and group were found, a significant effect was observed between time points ($F = 357.11, p < 0.001$) (Table 4, Figure 3). The average

effect size for the elastic band group was $d=4.57, g=4.57$, while the average effect size for the dumbbell group was $d=4.00, g=4.00$. Both groups demonstrated a large effect between time points (Table 5), with the elastic band group showing a relatively larger effect size.




Figure 3. Pain level reduction after 6 weeks. NRS: Numerical rating scale.

Table 4. Analysis results of NRS.

Variable	Group	Pre (Mean±SD)	Post (Mean±SD)	Source	F	p
NRS	Elastic band group	7.7 ± 0.8	2.5 ± 1.4	Time	357.11	<0.001***
	Dumbbell group	7.9 ± 0.9	2.4 ± 1.7	Group	0.036	0.851
Time × Group						
Time × Group						

NRS: Numerical rating scale; Values are expressed as Mean ± SD; *: $p < 0.05$; **: $p < 0.01$; ***: $p < 0.001$.

Table 5. Effect size of the elastic band and dumbbell on NRS.

Variable	Group	Cohen's d	Hedges' g
NRS	Elastic band group	4.57	4.57
	Dumbbell group	4.00	4.00

NRS: Numerical rating scale.

DISCUSSION

Research on the effects of exercise in AC joint patients has been steadily reported, but studies on appropriate exercise methods or identifying resistance tools to enhance exercise efficiency are still lacking. While some studies are comparing elastic bands and dumbbells, the most commonly used resistance tools for strength training in the general population, there is a notable lack of research specifically targeting AC patients. As strengthening exercises are emphasized for AC patients, this study aims to compare the effects of

elastic bands and dumbbells to identify suitable resistance tools that can aid in shoulder recovery for AC patients.

The key result of this study is that there was no significant difference in shoulder ROM and NRS between the elastic band group and the dumbbell group, even when the same exercise program was applied to both groups. These results align with studies by Andersen et al. (14) and Lopes et al. (19), which suggested that there was no significant difference in shoulder recovery between elastic bands and dumbbells in AC

patients. Although the interaction effect between time and groups was not confirmed, both groups showed improvements in shoulder pain and ROM post-exercise compared to pre-exercise, supporting the positive effects of elastic band exercises on shoulder joints (20) and the positive influence of dumbbell exercises on shoulder recovery (21, 22), as mentioned in previous studies.

Based on the positive results observed in both groups, it can be concluded that strength training itself may have a significant impact on shoulder recovery, more so than the difference in resistance tools used. This supports the argument made by Escamilla et al. (23) regarding the necessity of strength exercises. Additionally, as suggested by Kim Yeon-soon (24) and Cho Han-su, Kim Won-moon (1), the combined application of exercises and joint mobilization likely contributed positively to pain recovery.

While previous studies have reported no significant difference between the effects of elastic bands and dumbbells on muscle activation, shoulder pain, and ROM recovery, this study further analyzed the extent of changes in ROM and NRS between pre- and post-assessment by calculating effect sizes. The results showed that the elastic band group had a greater effect on the recovery of shoulder ROM for flexion, external rotation, and internal rotation, while the dumbbell group had a more significant effect on abduction. This difference can be attributed to the nature of the resistance tools. While dumbbells rely on the moment arm and gravity, elastic bands provide variable resistance depending on their elasticity, and the direction of resistance can be adjusted based on the individual's chosen angle (25). The ease with which resistance and direction can be set using elastic bands may enhance exercise efficiency, allowing for more effective training without being restricted by the moment arm, which could explain why the elastic band group showed greater ROM recovery.

Furthermore, based on previous studies that observed greater improvements in pain, ROM, and strength in groups performing eccentric contractions with elastic bands (26) and both concentric and eccentric contractions (27), this study incorporated both types of muscle contractions during exercise, and the elastic band group showed greater improvement in ROM recovery.

Regarding the NRS effect size, the elastic band group showed more significant results than the dumbbell group (Table 5), which can be attributed to the nature of the NRS scale. NRS measures overall pain experienced by the participants, and since it is a comprehensive measure, the differences in the effects of resistance tools could vary depending on when the pain is felt. Additionally, the significant changes observed in the dumbbell group regarding the pain index are likely due to improved ROM recovery, shoulder strength, and stabilization, leading to reduced discomfort during daily activities.

Although all 16 participants showed reduced pain over the 6-week study period, functional disabilities, such as discomfort during daily movements, persisted longer than the pain. This suggests that the recovery of ROM was slower, highlighting the effectiveness of joint mobilization and exercises in reducing pain. Notably, the recovery of external and internal rotation was slower than other movements, with internal rotation showing the most limited improvement. The average recovery angle for internal rotation was 24.4°, significantly less than the 65° improvement observed in abduction, highlighting that recovery of movements in the transverse plane may require more time than the 6 weeks used in this study. Future research should focus on developing appropriate exercise methods for recovering internal rotation and other horizontal-plane movements in AC patients.

Due to the short duration of this study, it may not be sufficient to fully determine the differences in the effects of these two resistance tools or propose complete recovery methods for AC patients. Additionally, the study was limited to middle-aged women, and the different stages of AC were not considered, which may have influenced the results. Elastic bands also present challenges in determining resistance, as their resistance increases gradually and changes with angle, making it difficult to compare them with dumbbells that have fixed resistance. Future studies should focus on determining the appropriate weight settings for both elastic bands and dumbbells to ensure their effectiveness in shoulder recovery. Overall, a well-designed strength training program is likely to have a more positive impact on optimal recovery in AC patients than focusing solely on the differences between resistance tools.

CONCLUSION

This study demonstrated that both elastic bands and dumbbells effectively reduce shoulder pain and improve functional recovery in patients with AC. While no significant differences were found between groups, strength training overall showed meaningful improvements in ROM and pain. These results support the incorporation of resistance exercises as part of non-surgical interventions for shoulder rehabilitation.

APPLICABLE REMARKS

- Muscles move and feel sensations under the control of nerves. Therefore, when musculoskeletal pain occurs, applying interventions that allow the muscles to move smoothly helps relieve pain.
- This study confirmed the effect of reducing pain by using the shoulder joint muscles of subjects experiencing adhesive capsulitis.

ACKNOWLEDGMENTS

This research would not have been possible without the participation of individuals experiencing shoulder discomfort. I sincerely appreciate their involvement, which has contributed to the success of this study.

AUTHORS' CONTRIBUTIONS

Study concept and design: Hee-Jin Kim, Hea Kyung Choi. Acquisition of data: Hee-Jin Kim, Hea Kyung Choi. Analysis and interpretation of data: Hee-Jin Kim, Hea Kyung Choi. Drafting the

manuscript: Hee-Jin Kim, Hea Kyung Choi. Critical revision of the manuscript for important intellectual content: Hea Kyung Choi. Statistical analysis: Hea Kyung Choi. Administrative, technical, and material support: Hea Kyung Choi. Study supervision: Hea Kyung Choi.

CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

FINANCIAL DISCLOSURE

This study has no financial interests related to the material in the manuscript.

FUNDING/SUPPORT

This study has no funding or support for the manuscript.

ETHICAL CONSIDERATION

This study protocol adheres to the ethical principles outlined in the 1975 Declaration of Helsinki.

ROLE OF THE SPONSOR

The funding organizations are public institutions and had no role in the design and conduct of the study.

ARTIFICIAL INTELLIGENCE (AI) USE

The study did not utilize AI or AI-assisted tools.

REFERENCES

1. Cho HS, Kim WM. The effect of pain level, range of motion and myofunction on the combined treatment of adhesive capsulitis patients. *Journal of the Korean Society for Wellness*. 2021;16(1):303-309. [\[doi:10.21097/ksw.2021.02.16.1.303\]](https://doi.org/10.21097/ksw.2021.02.16.1.303)
2. Pandey V, Madi S. Clinical Guidelines in the Management of Frozen Shoulder: An Update! *Indian Journal of Orthopaedics*. 2021;55(2):299-309. [\[doi:10.1007/s43465-021-00351-3\]](https://doi.org/10.1007/s43465-021-00351-3) [\[PMid:33912325\]](https://pubmed.ncbi.nlm.nih.gov/33912325/)
3. Dias R, Cutts S, Massoud S. Frozen shoulder. *British Medical Journal*. 2005;331(7530):1453-6. [\[doi:10.1136/bmj.331.7530.1453\]](https://doi.org/10.1136/bmj.331.7530.1453) [\[PMid:16356983\]](https://pubmed.ncbi.nlm.nih.gov/16356983/)
4. Koo SW, Chae JR, Kim HJ. Effects of the upper extremity pattern and rhythmic stabilization technique of PNF on the ROM and VAS in Frozen shoulder patients. *The Korean Journal of Physical Education*. 2004;43(3):623-631.
5. Hand GC, Athanasou NA, Matthews T, Carr AJ. The pathology of frozen shoulder. *Journal of Bone and Joint Surgery Br*. 2007;89(7):928-32. [\[doi:10.1302/0301-620X.89B7.19097\]](https://doi.org/10.1302/0301-620X.89B7.19097) [\[PMid:17673588\]](https://pubmed.ncbi.nlm.nih.gov/17673588/)
6. Dudkiewicz I, Oran A, Salai M, Palti R, Pritsch M. Idiopathic adhesive capsulitis: long-term results of conservative treatment. *The Israel Medical Association*. 2004;6(9):524-6.
7. Hollmann L, Halaki M, Kamper SJ, Haber M, Ginn KA. Does muscle guarding play a role in range of motion loss in patients with frozen shoulder?. *Musculoskeletal Science and Practice*. 2018;37:64-8. [\[doi:10.1016/j.msksp.2018.07.001\]](https://doi.org/10.1016/j.msksp.2018.07.001) [\[PMid:29986193\]](https://pubmed.ncbi.nlm.nih.gov/29986193/)

8. Cucchi D, Marmotti A, De Giorgi S, Costa A, D'Apolito R, Conca M, Russo A, Saccomanno MF, de Girolamo L, SIGASCOT Research Committee. Risk factors for shoulder stiffness: current concepts. *Joints.* 2017;5(04):217-23. [[doi:10.1055/s-0037-1608951](https://doi.org/10.1055/s-0037-1608951)] [[PMid:29270559](#)]
9. Wessner B. An elastic band exercise programme improves functional fitness in older adults. *Evidence-Based Nursing.* 2016;19(2):64. [[doi:10.1136/eb-2015-102193](https://doi.org/10.1136/eb-2015-102193)] [[PMid:26684686](#)]
10. Park HJ. Effects of progressive resistance exercise using elastic bands on blood glucose, blood lipids, muscle strength, and muscular endurance in elderly patients with diabetes. Master's thesis, Graduate School Korea University. 2001.
11. Brill PA, Probst JC, Greenhouse DL, Schell B, Macera CA. Clinical feasibility of a free-weight strength-training program for older adults. *Journal of American Board of Family Practice.* 1998;11(6):445-51. [[doi:10.3122/jabfm.11.6.445](https://doi.org/10.3122/jabfm.11.6.445)] [[PMid:9875999](#)]
12. Rodrigues da Silva Barros, B., Dal'Ava Augusto, D., de Medeiros Neto, J. F., Michener, L. A., Silva, R. S., & Sousa, C. D. O. Isometric versus isotonic exercise in individuals with rotator cuff tendinopathy—Effects on shoulder pain, functioning, muscle strength, and electromyographic activity: A protocol for randomized clinical trial. *Plos one.* 2023;18(11):e0293457. [[doi:10.1371/journal.pone.0293457](https://doi.org/10.1371/journal.pone.0293457)] [[PMid:37956135](#)]
13. Ribeiro F, Teixeira F, Brochado G, Oliveira J. Impact of low cost strength training of dorsi- and plantar flexors on balance and functional mobility in institutionalized elderly people. *Geriatrics Gerontology International.* 2009;9(1):75-80. [[doi:10.1111/j.1447-0594.2008.00500.x](https://doi.org/10.1111/j.1447-0594.2008.00500.x)] [[PMid:19260983](#)]
14. Andersen LL, Andersen CH, Mortensen OS, Poulsen OM, Bjørnlund IB, Zebis MK. Muscle activation and perceived loading during rehabilitation exercises: comparison of dumbbells and elastic resistance. *Physical Therapy.* 2010;90(4):538-49. [[doi:10.2522/ptj.20090167](https://doi.org/10.2522/ptj.20090167)] [[PMid:20133444](#)]
15. Sundstrup E, Jakobsen MD, Andersen CH, Bandholm T, Thorborg K, Zebis MK, Andersen LL. Evaluation of elastic bands for lower extremity resistance training in adults with and without musculoskeletal pain. *Scandinavian Journal Medicine and Science in Sports.* 2014;24(5):e353-9. [[doi:10.1111/sms.12187](https://doi.org/10.1111/sms.12187)]
16. Alghadir AH, Anwer S, Iqbal A, Iqbal ZA. Test-retest reliability, validity, and minimum detectable change of visual analog, numerical rating, and verbal rating scales for measurement of osteoarthritic knee pain. *Journal of Pain Res.* 2018;11:851-856. [[doi:10.2147/JPR.S158847](https://doi.org/10.2147/JPR.S158847)] [[PMid:29731662](#)]
17. Godges JJ, Mattson-Bell M, Thorpe D, Shah D. The immediate effects of soft tissue mobilization with proprioceptive neuromuscular facilitation on glenohumeral external rotation and overhead reach. *Journal of Orthopaedic and Sports Physical Therapy.* 2003;33(12):713-8. [[doi:10.2519/jospt.2003.33.12.713](https://doi.org/10.2519/jospt.2003.33.12.713)] [[PMid:14743984](#)]
18. Eliason A, Harringe M, Engström B, Werner S. Guided exercises with or without joint mobilization or no treatment in patients with subacromial pain syndrome: A clinical trial. *Journal of Rehabilitation Medicine.* 2021;53(5):jrm00190. [[doi:10.2340/16501977-2806](https://doi.org/10.2340/16501977-2806)] [[PMid:33634829](#)]
19. Lopes JSS, Machado AF, Micheletti JK, de Almeida AC, Cavina AP, Pastre CM. Effects of training with elastic resistance versus conventional resistance on muscular strength: A systematic review and meta-analysis. *SAGE Open Med.* 2019;7:2050312119831116. [[doi:10.1177/2050312119831116](https://doi.org/10.1177/2050312119831116)] [[PMid:30815258](#)]
20. Lee IH, Moon SG, Han DU. Treatment effect by Thera-Band for two cases of frozen shoulder. *Journal of the Korean Physical Therapy Association.* 1997;4(2):453-460.
21. Park MC, Lee SY. Deltoid middle fiber of the isometric contraction according to elastic band and dumbbell on various shoulder abduction angle. *Journal of the Korean Society of Physical Medicine.* 2016;11(2):71-76. [[doi:10.13066/kspm.2016.11.2.71](https://doi.org/10.13066/kspm.2016.11.2.71)]
22. Lee JN, You GH, Shin YA. Comparative analysis of shoulder muscle activity during cast movements for clubbell and dumbbell exercises. *Journal of Coaching Development.* 2021;23(3):160-169. [[doi:10.47684/jcd.2021.09.23.3.160](https://doi.org/10.47684/jcd.2021.09.23.3.160)]
23. Escamilla RF, Yamashiro K, Paulos L, Andrews JR. Shoulder muscle activity and function in common shoulder rehabilitation exercises. *Sports Medicine.* 2009;39(8):663-85. [[doi:10.2165/00007256-200939080-00004](https://doi.org/10.2165/00007256-200939080-00004)] [[PMid:19769415](#)]

24. Kim YS. Physical Science : Effects of Compound Program of Therapeutic Exercise and Therapeutic Massage on Pain and ROM(Range of Motion) in Patient with Frozen Shoulder. Korea Sports Research. 2005;16(2):237-249.

25. Brill PA, Probst JC, Greenhouse DL, Schell B, Macera CA. Clinical feasibility of a free-weight strength-training program for older adults. Journal of American Board of Family Practice. 1998;11(6):445-51. [doi:10.3122/jabfm.11.6.445] [PMid:9875999]

26. Kim WM, Seo YG, Park YJ, Cho HS, Lee SA, Jeon SJ, Ji SM. Effects of different types of contraction exercises on shoulder function and muscle strength in patients with adhesive capsulitis. International Journal of Environmental Research and Public Health. 2021;18(24):13078. [doi:10.3390/ijerph182413078] [PMid:34948688]

27. Han DW, Ahn YR, Lee NJ, Lee EJ. Effects of combined resistance training program of concentric and eccentric contraction using Theraband on shoulder rotation torque max and peak power. Korean Journal of Sport Biomechanics. 2009;19(1):139-148. [doi:10.5103/KJSB.2009.19.1.139]